

Faslodex® (fulvestrant)

(Intramuscular)

Document Number: IC-0043

Last Review Date: 03/01/2022 Date of Origin: 01/01/2012

Dates Reviewed: 03/2012, 06/2012, 09/2012, 12/2012, 03/2013, 06/2013, 09/2013, 12/2013, 03/2014, 06/2014, 09/2014, 12/2014, 03/2015, 05/2015, 08/2015, 11/2015, 02/2016, 05/2016, 08/2016, 11/2016, 02/2017, 05/2017, 08/2017, 09/2017, 11/2017, 02/2018, 05/2018, 09/2018, 12/2018, 03/2019, 04/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020, 09/2020, 03/2021, 03/2022

I. Length of Authorization

Coverage will be provided for six months and may be renewed.

II. Dosing Limits

- A. Quantity Limit (max daily dose) [NDC Unit]:
- Faslodex 250 mg/5 mL single-dose prefilled syringe: 6 syringes first 29 days initially (loading doses), then 2 syringes per 28 days thereafter as maintenance
- B. Max Units (per dose and over time) [HCPCS Unit]:

Ovarian Cancer

Loading Dosing:

• 20 billable units on day 1 and 10 billable units on days 15 and 29

Maintenance Dosing:

• 10 billable units every 28 days

Endometrial Cancer

• 10 billable units every 28 days

Breast Cancer/Uterine Sarcoma

Loading Dosing:

• 20 billable units every 14 days for 3 doses

Maintenance Dosing:

• 20 billable units every 28 days

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

Patient is at least 18 years of age; AND

Breast Cancer † 1,2,5,8-11

• Patient is postmenopausal, premenopausal with ovarian ablation/suppression, or male with suppression of testicular steroidogenesis; AND



- Patient has advanced, metastatic, or recurrent unresectable invasive disease; AND
 - ➤ Patient has hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative disease with no visceral crisis; **AND**
 - Used as initial therapy; AND
 - Used as a single agent †; OR
 - Used in combination with a CDK 4/6 inhibitor (abemaciclib, palbociclib, or ribociclib) †; OR
 - Used in combination with a non-steroidal aromatase inhibitor (i.e., anastrozole or letrozole) ‡; OR
 - Used as subsequent therapy in patients; **AND**
 - Used as a single agent †; OR
 - Used in combination with a CDK 4/6 inhibitor (abemaciclib, palbociclib, or ribociclib) if a CDK 4/6 inhibitor was not previously used †; **OR**
 - Used in combination with everolimus ‡; **OR**
 - Used in combination with alpelisib in patients who have PIK3CA activating mutation positive disease ‡; OR
 - ➤ Patient has HR-positive, HER2-positive disease ‡; AND
 - Used as a single agent or in combination with trastuzumab; OR
- o Patient has recurrent unresectable or metastatic inflammatory disease ‡; AND
 - ➤ Patient has hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative disease with no visceral crisis; **AND**
 - Used as initial therapy; AND
 - Used as a single agent; OR
 - Used in combination with a non-steroidal aromatase inhibitor (i.e., anastrozole or letrozole); OR
 - Used in combination with a CDK 4/6 inhibitor (abemaciclib, palbociclib, or ribociclib); OR
 - Used as subsequent therapy; AND
 - Used as a single agent; **OR**
 - Used in combination with everolimus; OR
 - Used in combination with a CDK 4/6 inhibitor (abemaciclib, palbociclib, or ribociclib) if a CDK 4/6 inhibitor was not previously used; OR
 - Used in combination with alpelisib in patients who have PIK3CA activating mutation positive disease; OR
 - ➤ Patient has HR-positive, HER2-positive disease; AND
 - Used as a single agent or in combination with trastuzumab

Ovarian Cancer (epithelial, fallopian tube, or primary peritoneal cancer) ‡ 2,7,12



- Used as single agent therapy; AND
- Patient has recurrent low-grade serous carcinoma; AND
- Patient has previously been received treatment with an aromatase inhibitor (i.e., letrozole, anastrozole, exemestane)

Endometrial Adenocarcinoma (Uterine Neoplasms) ‡ 2,6,13

- Used as single agent therapy; AND
- Patient has grade 1 or 2 endometrioid histology; AND
- Used in patients with a small tumor volume or an indolent growth pace; AND
- Used as ONE of the following:
 - Adjuvant treatment for stage III-IV disease; OR
 - o Treatment for disseminated metastases or locoregional recurrence; OR
 - Primary treatment in patients with disease limited to the uterus that is not suitable for primary surgery; OR
 - o Primary treatment in patients undergoing both sequential brachytherapy and external beam radiation therapy (EBRT) for disease with cervical involvement that is not suitable for primary surgery; **OR**
 - o Primary treatment in patients undergoing sequential EBRT for locoregional extrauterine disease that is not suitable for primary surgery; **OR**
 - o Primary treatment in patients with distant metastatic disease

Uterine Sarcoma (Uterine Neoplasms) ‡ 2,13

- Used as single agent therapy; AND
- Used in patients with a small tumor volume or an indolent growth pace; AND
- Used for low-grade endometrial stromal sarcoma (ESS), adenosarcoma without sarcomatous overgrowth, or ER/PR positive uterine sarcoma; AND
 - Used following total hysterectomy for stage II-IV disease; OR
 - Used for metastatic or recurrent disease; OR
 - Used for disease that is not suitable for primary surgery

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); **Φ** Orphan Drug

IV. Renewal Criteria 1

Coverage can be renewed based upon the following criteria:

• Patient continues to meet the indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**



- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: bleeding abnormalities, severe injection site reactions, etc.

V. Dosage/Administration ^{1,6,7}

Indication	Dose	
Breast Cancer	Loading Dose:	
& Uterine	• 500 mg intramuscularly (IM) on Days 1, 15, 29	
Sarcoma	Maintenance Dose:	
	• 500 mg IM every 28 days	
	Loading Dose:	
Ovarian Cancer	• 500 mg intramuscularly (IM) on Day 1 and 250 mg IM on Days 15 and 29	
Ovarian Cancer	Maintenance Dose:	
	• 250 mg IM every 28 days	
Endometrial	250 mg by IM injection every 4 weeks for at least 8 weeks. Therapy should be	
Cancer	continued until evidence of progressive disease or adverse effects prevent further	
	treatment.	

VI. Billing Code/Availability Information

HCPCS Code:

- J9395 Injection, fulvestrant, 25 mg; 1 billable unit = 25 mg
- J9393 Injection, fulvestrant (teva) not therapeutically equivalent to J9395, 25 mg; 1 billable unit = 25 mg (*Effective 01/01/2023*)
- J9394 Injection, fulvestrant (fresenius kabi) not therapeutically equivalent to J9395, 25 mg; 1 billable unit = 25 mg (*Effective 01/01/2023*)

NDC:

Faslodex* 250 mg/5 mL single-dose prefilled syringe: 00310-0720-xx
*Available generically

VII. References

- 1. Faslodex [package insert]. Wilmington, DE; AstraZeneca Pharmaceuticals LP; January 2021. Accessed January 2022.
- 2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for fulvestrant. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most



- recent and complete version of the Compendium, go online to NCCN.org. Accessed January 2022.
- 3. Chia S, Gradishar W, Mauriac L, et al. Double-blind, randomized placebo-controlled trial of fulvestrant compared with exemestane after prior nonsteroidal aromatase inhibitor therapy in postmenopausal women with hormone-receptor positive, advanced breast cancer: results from EFECT. J Clin Oncol 2008; 26:1664-1670.
- 4. Mauriac L, Romieu G, Bines J. Activity of fulvestrant versus exemestane in advanced breast cancer patients with or without visceral metastases: data from the EFECT trial. Breast Cancer Res Treat 2009; 117:69-75.
- 5. Di Leo A, Jerusalem G, Petruzelka L, et al. Results of the CONFIRM phase III trial comparing fulvestrant 250 mg with fulvestrant 500 mg in postmenopausal women with estrogen receptor-positive advanced breast cancer. J Clin Oncol 2010; 28:4594-4600.
- 6. Covens AL, Filiaci V, Gersell D. Phase II study of fulvestrant in recurrent/metastatic endometrial carcinoma: a Gynecologic Oncology Group study. Gynecol Oncol. 2011 Feb;120(2):185-8. doi: 10.1016/j.ygyno.2010.10.015. Epub 2010 Nov 13.
- 7. Argenta PA, Thomas SG, Judson PL, et al. A phase II study of fulvestrant in the treatment of multiply-recurrent epithelial ovarian cancer. Gynecol Oncol. 2009 May;113(2):205-209.
- 8. Robertson JFR, Bondarenko IM, Trishkina E, et al. Fulvestrant 500 mg versus anastrozole 1 mg for hormone receptor-positive advanced breast cancer (FALCON): an international, randomised, double-blind, phase 3 trial. Lancet. 2016;388(10063):2997-3005. doi:10.1016/S0140-6736(16)32389-3.
- 9. Robertson JF, Osborne CK, Howell A, et al. Fulvestrant versus anastrozole for the treatment of advanced breast carcinoma in postmenopausal women: a prospective combined analysis of two multicenter trials. Cancer. 2003;98(2):229-238. doi:10.1002/cncr.11468.
- 10. Cristofanilli M, Turner NC, Bondarenko I, et al. Fulvestrant plus palbociclib versus fulvestrant plus placebo for treatment of hormone-receptor-positive, HER2-negative metastatic breast cancer that progressed on previous endocrine therapy (PALOMA-3): final analysis of the multicentre, double-blind, phase 3 randomised controlled trial [published correction appears in Lancet Oncol. 2016 Apr;17 (4):e136] [published correction appears in Lancet Oncol. 2016 Jul;17 (7):e270]. Lancet Oncol. 2016;17(4):425-439. doi:10.1016/S1470-2045(15)00613-0.
- 11. Sledge GW Jr, Toi M, Neven P, et al. MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy. J Clin Oncol. 2017;35(25):2875-2884. doi:10.1200/JCO.2017.73.7585.
- 12. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer Version 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network,



- Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed January 2022.
- 13. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Uterine Neoplasms Version 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed January 2022.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description	
C48.1	Malignant neoplasm of specified parts of peritoneum	
C48.2	Malignant neoplasm of peritoneum, unspecified	
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum	
C50.011	Malignant neoplasm of nipple and areola, right female breast	
C50.012	Malignant neoplasm of nipple and areola, left female breast	
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast	
C50.021	Malignant neoplasm of nipple and areola, right male breast	
C50.022	Malignant neoplasm of nipple and areola, left male breast	
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast	
C50.111	Malignant neoplasm of central portion of right female breast	
C50.112	Malignant neoplasm of central portion of left female breast	
C50.119	Malignant neoplasm of central portion of unspecified female breast	
C50.121	Malignant neoplasm of central portion of right male breast	
C50.122	Malignant neoplasm of central portion of left male breast	
C50.129	Malignant neoplasm of central portion of unspecified male breast	
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast	
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast	
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast	
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast	
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast	
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast	
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast	
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast	
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast	
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast	
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast	
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast	
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast	
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast	

ICD-10	ICD-10 Description	
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast	
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast	
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast	
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast	
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast	
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast	
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast	
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast	
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast	
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast	
C50.611	Malignant neoplasm of axillary tail of right female breast	
C50.612	Malignant neoplasm of axillary tail of left female breast	
C50.619	Malignant neoplasm of axillary tail of unspecified female breast	
C50.621	Malignant neoplasm of axillary tail of right male breast	
C50.622	Malignant neoplasm of axillary tail of left male breast	
C50.629	Malignant neoplasm of axillary tail of unspecified male breast	
C50.811	Malignant neoplasm of overlapping sites of right female breast	
C50.812	Malignant neoplasm of overlapping sites of left female breast	
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast	
C50.821	Malignant neoplasm of overlapping sites of right male breast	
C50.822	Malignant neoplasm of overlapping sites of left male breast	
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast	
C50.911	Malignant neoplasm of unspecified site of right female breast	
C50.912	Malignant neoplasm of unspecified site of left female breast	
C50.919	Malignant neoplasm of unspecified site of unspecified female breast	
C50.921	Malignant neoplasm of unspecified site of right male breast	
C50.922	Malignant neoplasm of unspecified site of left male breast	
C50.929	Malignant neoplasm of unspecified site of unspecified male breast	
C54.0	Malignant neoplasm of isthmus uteri	
C54.1	Malignant neoplasm of endometrium	
C54.2	Malignant neoplasm of myometrium	
C54.3	Malignant neoplasm of fundus uteri	
C54.8	Malignant neoplasm of overlapping sites of corpus uteri	
C54.9	Malignant neoplasm of corpus uteri, unspecified	
C55	Malignant neoplasm of uterus, part unspecified	
C56.1	Malignant neoplasm of right ovary	
C56.2	Malignant neoplasm of left ovary	
C56.3	Malignant neoplasm of bilateral ovaries	
C56.9	Malignant neoplasm of unspecified ovary	



ICD-10	ICD-10 Description	
C57.00	Malignant neoplasm of unspecified fallopian tube	
C57.01	Malignant neoplasm of right fallopian tube	
C57.02	Malignant neoplasm of left fallopian tube	
C57.10	Malignant neoplasm of unspecified broad ligament	
C57.11	Malignant neoplasm of right broad ligament	
C57.12	Malignant neoplasm of left broad ligament	
C57.20	Malignant neoplasm of unspecified round ligament	
C57.21	Malignant neoplasm of right round ligament	
C57.22	Malignant neoplasm of left round ligament	
C57.3	Malignant neoplasm of parametrium	
C57.4	Malignant neoplasm of uterine adnexa, unspecified	
C57.7	Malignant neoplasm of other specified female genital organs	
C57.8	Malignant neoplasm of overlapping sites of female genital organs	
C57.9	Malignant neoplasm of female genital organ, unspecified	
Z85.43	Personal history of malignant neoplasm of ovary	

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC		
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC		
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)		
6	MN, WI, IL	National Government Services, Inc. (NGS)		
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.		
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)		
N (9)	FL, PR, VI	First Coast Service Options, Inc.		
J (10)	TN, GA, AL	Palmetto GBA, LLC		
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC		



Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.		
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)		
15	KY, OH	CGS Administrators, LLC		

