

Adcetris® (brentuximab vedotin) (Intravenous)

Document Number: IC-0004

Last Review Date: 09/01/2020

Date of Origin: 02/28/2012

Dates Reviewed: 06/2012, 09/2012, 09/2012, 12/2012, 03/2013, 06/2013, 09/2013, 12/2013, 03/2014, 06/2014, 09/2014, 12/2014, 03/2015, 05/2015, 08/2015, 11/2015, 02/2016, 05/2016, 08/2016, 11/2016, 02/2017, 05/2017, 08/2017, 11/2017, 02/2018, 04/2018, 09/2018, 12/2018, 03/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020, 09/2020

I. Length of Authorization ^{1,7}

Coverage will be provided for six months and may be renewed.

- Treatment for cHL post-auto HSCT, Mycosis Fungoides (MF)/Sezary Syndrome (SS), and Primary Cutaneous CD30+ T-Cell Lymphoproliferative Disorders has a maximum of 16 cycles.
- Treatment of previously untreated Stage III or IV Classical Hodgkin Lymphoma (cHL) has a maximum of 12 doses.
- Treatment of previously untreated Systemic Anaplastic Large Cell Lymphoma (sALCL) or other CD30-expressing Peripheral T-Cell Lymphomas (PTCL) has a maximum of 8 doses.
- Treatment of Breast-Implant Associated Anaplastic Large Cell Lymphoma (ALCL) has a maximum of 6 cycles as adjuvant therapy.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- 50 mg vial: 4 vials per 21 days

B. Max Units (per dose and over time) [HCPCS Unit]:

- 200 billable units per 21 days

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

Universal Criteria ¹

- Patient has CD30-positive disease; **AND**
- Patient must not be receiving concomitant bleomycin; **AND**

Classical Hodgkin Lymphoma (cHL) † Φ 1,2,4,12-14

- Used as single agent therapy; **AND**
 - Used as consolidation/maintenance therapy post-autologous hematopoietic stem cell transplant (auto-HSCT) in patients at high risk* for relapse or progression † Φ ; **OR**
 - Patient has relapsed disease after failure of auto-HSCT or after failure of at least 2 prior multi-agent chemotherapy regimens in patients who are not auto-HSCT candidates †; **OR**
 - Used as second-line or subsequent systemic therapy (if not previously used) for relapsed or refractory disease Φ ; **OR**
 - Used as palliative therapy for relapsed or refractory disease in patients >60 years of age Φ ; **OR**
- Used in combination with bendamustine; **AND**
 - Used as second-line or subsequent systemic therapy (if not previously used) for relapsed or refractory disease Φ ; **OR**
- Used in combination with nivolumab; **AND**
 - Used as second-line or subsequent systemic therapy (if not previously used) for relapsed or refractory disease Φ ; **OR**
- Used in combination with dacarbazine; **AND**
 - Used as primary treatment in patients > 60 years of age with stage I-II unfavorable or stage III-IV disease Φ ; **OR**
- Used in combination with doxorubicin, vinblastine, and dacarbazine (AVD); **AND**
 - Used as initial therapy for previously untreated stage III or IV disease †; **OR**
 - Used as initial therapy for previously untreated stage I or II unfavorable disease in patients >60 years of age Φ

**High risk for relapse or progression may be defined as:*

- *Refractory disease, relapse within 12 months, or extranodal disease following frontline therapy OR 2 or more of the following: PET+ response at time of transplant, B symptoms, and/or >1 salvage/subsequent therapy regimen)*

T-Cell Lymphomas 1-3,15,16

- Peripheral T-Cell Lymphoma (PTCL)
 - Used as a single agent after failure of at least one prior chemotherapy regimen for one of the following:
 - Systemic Anaplastic Large Cell Lymphoma (sALCL) † Φ
 - Peripheral T-Cell Lymphoma (PTCL) not otherwise specified Φ Φ
 - Angioimmunoblastic T-cell Lymphoma (AITL) Φ Φ ; **OR**
 - Used in combination with cyclophosphamide, doxorubicin, and prednisone (CHP) as initial therapy for previously untreated:
 - Systemic Anaplastic Large Cell Lymphoma (sALCL) † Φ
 - Peripheral T-Cell Lymphoma (PTCL) not otherwise specified † Φ
 - Angioimmunoblastic T-cell Lymphoma (AITL) † Φ

- Enteropathy-associated T-cell lymphoma Φ , monomorphic epitheliotropic intestinal T-cell lymphoma, nodal peripheral T-cell lymphoma with TFH phenotype, or follicular T-cell lymphoma \ddagger
- Breast-Implant Associated Anaplastic Large Cell Lymphoma (ALCL) \ddagger ; **AND**
 - Used as adjuvant therapy as a single agent or in combination with cyclophosphamide, doxorubicin, and prednisone (CHP); **AND**
 - Patient has localized disease to the capsule, implant, or breast with either lymph node involvement or radiation therapy is not feasible; **OR**
 - Patient has extended disease (stage II-IV)
- Adult T-Cell Leukemia/Lymphoma \ddagger Φ
 - Used as a single agent for acute disease or lymphoma; **AND**
 - Used as subsequent therapy for non-responders to first-line therapy; **OR**
 - Used in combination with cyclophosphamide, doxorubicin, and prednisone (CHP); **AND**
 - Used as first-line therapy or continued treatment in responders to first-line therapy for acute disease or lymphoma; **OR**
 - Used as subsequent therapy for non-responders to first-line therapy for chronic or smoldering disease
- Extranodal NK/T-Cell Lymphoma (Nasal type) \ddagger Φ
 - Used as a single agent for relapsed or refractory disease; **AND**
 - Patient has previously received at least 2 different prior lines of therapy including an asparaginase based combination chemotherapy regimen
- Hepatosplenic Gamma-Delta T-Cell Lymphoma \ddagger
 - Used as single-agent therapy; **AND**
 - Used for refractory disease as subsequent therapy after progression on two primary treatment regimens; **OR**
 - Used in combination with cyclophosphamide, doxorubicin, and prednisone (CHP); **AND**
 - Used as primary treatment or as an alternate induction regimen if not used during primary treatment

Primary Cutaneous Lymphomas ^{1,2,17}

- Mycosis Fungoides (MF) \ddagger Φ /Sezary Syndrome (SS) \ddagger
 - Used as a single agent; **AND**
 - Used as primary therapy in patients without stage IA MF with B1 blood involvement; **OR**
 - Used as subsequent therapy
- Primary Cutaneous CD30+ T-Cell Lymphoproliferative Disorders \ddagger Φ
 - Used as a single agent; **AND**
 - Patient has primary cutaneous anaplastic large cell lymphoma (pcALCL) \ddagger ; **OR**
 - Patient has cutaneous ALCL with regional node (N1) (excludes systemic ALCL); **OR**

- Patient has lymphomatoid papulosis (LyP) with extensive lesions that is relapsed or refractory to treatment options (e.g., clinical trial, observation, retreatment with primary treatment, or treatment with alternative regimen); **OR**
- Used in combination with cyclophosphamide, doxorubicin, and prednisone (CHP); **AND**
 - Patient has cutaneous ALCL with regional node (N1) (excludes systemic ALCL)

B-Cell Lymphomas †^{2,11}

- Diffuse Large B-Cell Lymphoma (DLBCL) or High Grade B-Cell Lymphoma
 - Used as second-line or subsequent therapy for partial response, no response, relapsed, progressive, or refractory disease in non-candidates for transplant
- AIDS-related DLBCL, primary effusion lymphoma, or HHV8-positive DLBCL, not otherwise specified
 - Used as second-line or subsequent therapy for relapsed disease in non-candidates for transplant
- Monomorphic Post-Transplant Lymphoproliferative Disorders (PTLD)
 - Used as second-line or subsequent therapy for patients with partial response, persistent, or progressive disease after receiving chemoimmunotherapy as first-line treatment for B-cell type disease; **OR**
 - Used in combination with CHP (cyclophosphamide, doxorubicin, prednisone) for T-cell type disease
- Histologic transformation of Follicular Lymphoma or Nodal Marginal Zone Lymphoma to Diffuse Large B-Cell Lymphoma (DLBCL) in patients who have received multiple lines of therapy for transformed or indolent disease

† FDA Approved Indication(s); ‡ Compensia Recommended Indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria¹

Coverage can be renewed based on the following criteria:

- Patient continues to meet universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Disease response with treatment defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: progressive multifocal leukoencephalopathy, peripheral neuropathy, anaphylaxis and infusion reactions, hematologic toxicities (thrombocytopenia, neutropenia, and anemia), serious infections, opportunistic infections, tumor lysis syndrome, increased toxicity in patients with severe renal (CrCl < 30 ml/min) and hepatic impairment (Child-Pugh B or C), hepatotoxicity, pulmonary toxicity, serious dermatologic reactions, gastrointestinal complications, uncontrolled hyperglycemia, etc.

V. Dosage/Administration ^{1,7}

Indication	Dose
Previously Untreated Stage III or IV Classical Hodgkin Lymphoma	1.2 mg/kg (up to 120 mg) by intravenous infusion every 2 weeks until a maximum of 12 doses, disease progression, or unacceptable toxicity
cHL post-auto HSCT, MF/SS, Primary Cutaneous CD30+ T-Cell Lymphoproliferative Disorders	1.8 mg/kg (up to 180 mg) by intravenous infusion every 3 weeks until a maximum of 16 cycles, disease progression, or unacceptable toxicity
Previously Untreated sALCL or Other CD30-expressing PTCL	1.8 mg/kg (up to 180 mg) by intravenous infusion every 3 weeks with each cycle of chemotherapy for a maximum of 6 to 8 doses
Breast-Implant Associated Anaplastic Large Cell Lymphoma (ALCL)	1.8 mg/kg (up to 180 mg) by intravenous infusion every 3 weeks until a maximum of 6 cycles, disease progression, or unacceptable toxicity
All other indications	1.8 mg/kg (up to 180 mg) by intravenous infusion every 3 weeks until disease progression or unacceptable toxicity

VI. Billing Code/Availability Information

HCPSC Code:

- J9042 - Injection, brentuximab vedotin, 1 mg; 1 billable unit = 1 mg

NDC:

- Adcetris single-use vial; 50 mg powder for injection: 51144-0050-xx

VII. References

1. Adcetris [package insert]. Bothell, WA; Seattle Genetics, Inc; October 2019. Accessed August 2020.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for brentuximab vedotin. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2020.
3. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) T-Cell Lymphomas. Version 1.2020. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2020.

4. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hodgkin Lymphoma. Version 2.2020. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2020.
5. Duvoc M, Tetzlaff MT, Gangar P, et al. Results of a Phase II trial of brentuximab vedotin for CD20+ cutaneous T-cell lymphoma and lymphomatoid papulosis. *J Clin Oncol* 2015; 33:3759-65.
6. Horwitz SM, Advani RH, Bartlett NL, et al. Objective responses in relapsed T-cell lymphomas with single-agent brentuximab vedotin. *Blood* 2014;123:3095-3100.
7. Alderuccio, JP., Desai, A., Yepes, M.M., et al. Frontline brentuximab vedotin in breast implant-associated anaplastic large-cell lymphoma. *Clin Case Rep* 2018; 6(4): 634-637. doi:10.1002/ccr3.1382.
8. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. *J Oncol Pract.* 2018 Mar;14(3):e130-e136.
9. Hematology/Oncology Pharmacy Association (2019). *Intravenous Cancer Drug Waste Issue Brief*. Retrieved from http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug_Waste_2019.pdf
10. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. *BMJ.* 2016 Feb 29;352:i788.
11. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) B-Cell Lymphomas. Version 4.2020. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2020.
12. Connors JM, Jurczak W, Straus DJ, et al. Brentuximab Vedotin with Chemotherapy for Stage III or IV Hodgkin's Lymphoma [published correction appears in *N Engl J Med.* 2018 Mar 1;378(9):878]. *N Engl J Med.* 2018;378(4):331-344.
13. Moskowitz CH, Nademanee A, Masszi T, et al. Brentuximab vedotin as consolidation therapy after autologous stem-cell transplantation in patients with Hodgkin's lymphoma at risk of relapse or progression (AETHERA): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet.* 2015;385(9980):1853-1862.
14. Younes A, Gopal AK, Smith SE, et al. Results of a pivotal phase II study of brentuximab vedotin for patients with relapsed or refractory Hodgkin's lymphoma. *J Clin Oncol.* 2012;30(18):2183-2189.

15. Horwitz S, O'Connor OA, Pro B, et al. Brentuximab vedotin with chemotherapy for CD30-positive peripheral T-cell lymphoma (ECHELON-2): a global, double-blind, randomised, phase 3 trial. *Lancet*. 2019;393(10168):229-240.
16. Pro B, Advani R, Brice P, et al. Brentuximab vedotin (SGN-35) in patients with relapsed or refractory systemic anaplastic large-cell lymphoma: results of a phase II study. *J Clin Oncol*. 2012;30(18):2190-2196.
17. Prince HM, Kim YH, Horwitz SM, et al. Brentuximab vedotin or physician's choice in CD30-positive cutaneous T-cell lymphoma (ALCANZA): an international, open-label, randomised, phase 3, multicentre trial. *Lancet*. 2017;390(10094):555-566.
18. Palmetto GBA. Local Coverage Article (LCA): Billing and Coding: Chemotherapy (A56141). Centers for Medicare & Medicaid Services, Inc. Updated on 05/26/2020 with effective date 04/30/2020. Accessed August 2020.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C81.10	Nodular sclerosis Hodgkin lymphoma, unspecified site
C81.11	Nodular sclerosis Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.12	Nodular sclerosis Hodgkin lymphoma, intrathoracic lymph nodes
C81.13	Nodular sclerosis Hodgkin lymphoma, intra-abdominal lymph nodes
C81.14	Nodular sclerosis Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.15	Nodular sclerosis Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.16	Nodular sclerosis Hodgkin lymphoma, intrapelvic lymph nodes
C81.17	Nodular sclerosis Hodgkin lymphoma, spleen
C81.18	Nodular sclerosis Hodgkin lymphoma, lymph nodes of multiple sites
C81.19	Nodular sclerosis Hodgkin lymphoma, extranodal and solid organ sites
C81.20	Mixed cellularity Hodgkin lymphoma, unspecified site
C81.21	Mixed cellularity Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.22	Mixed cellularity Hodgkin lymphoma, intrathoracic lymph nodes
C81.23	Mixed cellularity Hodgkin lymphoma, intra-abdominal lymph nodes
C81.24	Mixed cellularity Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.25	Mixed cellularity Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.26	Mixed cellularity Hodgkin lymphoma, intrapelvic lymph nodes
C81.27	Mixed cellularity Hodgkin lymphoma, spleen
C81.28	Mixed cellularity Hodgkin lymphoma, lymph nodes of multiple sites
C81.29	Mixed cellularity Hodgkin lymphoma, extranodal and solid organ sites
C81.30	Lymphocyte depleted Hodgkin lymphoma, unspecified site
C81.31	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of head, face, and neck

ICD-10	ICD-10 Description
C81.32	Lymphocyte depleted Hodgkin lymphoma, intrathoracic lymph nodes
C81.33	Lymphocyte depleted Hodgkin lymphoma, intra-abdominal lymph nodes
C81.34	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.35	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.36	Lymphocyte depleted Hodgkin lymphoma, intrapelvic lymph nodes
C81.37	Lymphocyte depleted Hodgkin lymphoma, spleen
C81.38	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of multiple sites
C81.39	Lymphocyte depleted Hodgkin lymphoma, extranodal and solid organ sites
C81.40	Lymphocyte-rich Hodgkin lymphoma, unspecified site
C81.41	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.42	Lymphocyte-rich Hodgkin lymphoma, intrathoracic lymph nodes
C81.43	Lymphocyte-rich Hodgkin lymphoma, intra-abdominal lymph nodes
C81.44	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.45	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.46	Lymphocyte-rich Hodgkin lymphoma, intrapelvic lymph nodes
C81.47	Lymphocyte-rich Hodgkin lymphoma, spleen
C81.48	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of multiple sites
C81.49	Lymphocyte-rich Hodgkin lymphoma, extranodal and solid organ sites
C81.70	Other Hodgkin lymphoma unspecified site
C81.71	Other Hodgkin lymphoma lymph nodes of head, face, and neck
C81.72	Other Hodgkin lymphoma intrathoracic lymph nodes
C81.73	Other Hodgkin lymphoma intra-abdominal lymph nodes
C81.74	Other Hodgkin lymphoma lymph nodes of axilla and upper limb
C81.75	Other Hodgkin lymphoma lymph nodes of inguinal region and lower limb
C81.76	Other Hodgkin lymphoma intrapelvic lymph nodes
C81.77	Other Hodgkin lymphoma spleen
C81.78	Other Hodgkin lymphoma lymph nodes of multiple sites
C81.79	Other Hodgkin lymphoma extranodal and solid organ sites
C81.90	Hodgkin lymphoma, unspecified, unspecified site
C81.91	Hodgkin lymphoma, unspecified, lymph nodes of head, face and neck
C81.92	Hodgkin lymphoma, unspecified, intrathoracic lymph nodes
C81.93	Hodgkin lymphoma, unspecified, intra-abdominal lymph nodes
C81.94	Hodgkin lymphoma, unspecified, lymph nodes of axilla and upper limb
C81.95	Hodgkin lymphoma, unspecified, lymph nodes of inguinal region and lower limb

ICD-10	ICD-10 Description
C81.96	Hodgkin lymphoma, unspecified, intrapelvic lymph nodes
C81.97	Hodgkin lymphoma, unspecified, spleen
C81.98	Hodgkin lymphoma, unspecified, lymph nodes of multiple sites
C81.99	Hodgkin lymphoma, unspecified, extranodal and solid organ sites
C83.30	Diffuse large B-cell lymphoma unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck
C83.32	Diffuse large B-cell lymphoma intrathoracic lymph nodes
C83.33	Diffuse large B-cell lymphoma intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma extranodal and solid organ sites
C83.80	Other non-follicular lymphoma, unspecified site
C83.81	Other non-follicular lymphoma, lymph nodes of head, face and neck
C83.82	Other non-follicular lymphoma, intrathoracic lymph nodes
C83.83	Other non-follicular lymphoma, intra-abdominal lymph nodes
C83.84	Other non-follicular lymphoma, lymph nodes of axilla and upper limb
C83.85	Other non-follicular lymphoma, lymph nodes of inguinal region and lower limb
C83.86	Other non-follicular lymphoma, intrapelvic lymph nodes
C83.87	Other non-follicular lymphoma, spleen
C83.88	Other non-follicular lymphoma, lymph nodes of multiple sites
C83.89	Other non-follicular lymphoma, extranodal and solid organ sites
C83.90	Non-follicular (diffuse) lymphoma, unspecified site
C83.91	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of head, face, and neck
C83.92	Non-follicular (diffuse) lymphoma, unspecified intrathoracic lymph nodes
C83.93	Non-follicular (diffuse) lymphoma, unspecified intra-abdominal lymph nodes
C83.94	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of axilla and upper limb
C83.95	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of inguinal region and lower limb
C83.96	Non-follicular (diffuse) lymphoma, unspecified intrapelvic lymph nodes
C83.97	Non-follicular (diffuse) lymphoma, unspecified spleen
C83.98	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of multiple sites
C83.99	Non-follicular (diffuse) lymphoma, unspecified extranodal and solid organ sites

ICD-10	ICD-10 Description
C84.00	Mycosis fungoides, unspecified site
C84.01	Mycosis fungoides, lymph nodes of head, face and neck
C84.02	Mycosis fungoides, intrathoracic lymph nodes
C84.03	Mycosis fungoides, intra-abdominal lymph nodes
C84.04	Mycosis fungoides, lymph nodes of axilla and upper limb
C84.05	Mycosis fungoides, lymph nodes of inguinal region and lower limb
C84.06	Mycosis fungoides, intrapelvic lymph nodes
C84.07	Mycosis fungoides, spleen
C84.08	Mycosis fungoides, lymph nodes of multiple sites
C84.09	Mycosis fungoides, extranodal and solid organ sites
C84.10	Sézary disease, unspecified site
C84.11	Sézary disease, lymph nodes of head, face, and neck
C84.12	Sézary disease, intrathoracic lymph nodes
C84.13	Sézary disease, intra-abdominal lymph nodes
C84.14	Sézary disease, lymph nodes of axilla and upper limb
C84.15	Sézary disease, lymph nodes of inguinal region and lower limb
C84.16	Sézary disease, intrapelvic lymph nodes
C84.17	Sézary disease, spleen
C84.18	Sézary disease, lymph nodes of multiple sites
C84.19	Sézary disease, extranodal and solid organ sites
C84.40	Peripheral T-cell lymphoma, not classified, unspecified site
C84.41	Peripheral T-cell lymphoma, not classified, lymph nodes of head, face and neck
C84.42	Peripheral T-cell lymphoma, not classified, intrathoracic lymph nodes
C84.43	Peripheral T-cell lymphoma, not classified, intra-abdominal lymph nodes
C84.44	Peripheral T-cell lymphoma, not classified, lymph nodes of axilla and upper limb
C84.45	Peripheral T-cell lymphoma, not classified, lymph nodes of inguinal region of lower limb
C84.46	Peripheral T-cell lymphoma, not classified, intrapelvic lymph nodes
C84.47	Peripheral T-cell lymphoma, not classified, spleen
C84.48	Peripheral T-cell lymphoma, not classified, lymph nodes of multiple sites
C84.49	Peripheral T-cell lymphoma, not classified, extranodal and solid organ sites
C84.60	Anaplastic large cell lymphoma, ALK-positive, unspecified site
C84.61	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of head, face and neck
C84.62	Anaplastic large cell lymphoma, ALK-positive, intrathoracic lymph nodes
C84.63	Anaplastic large cell lymphoma, ALK-positive, intra-abdominal lymph nodes

ICD-10	ICD-10 Description
C84.64	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of axilla and upper limb
C84.65	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of inguinal region and lower limb
C84.66	Anaplastic large cell lymphoma, ALK-positive, intrapelvic lymph nodes
C84.67	Anaplastic large cell lymphoma, ALK-positive, spleen
C84.68	Anaplastic large cell lymphoma, ALK-positive, lymph nodes of multiple sites
C84.69	Anaplastic large cell lymphoma, ALK-positive, extranodal and solid organ sites
C84.70	Anaplastic large cell lymphoma, ALK-negative, unspecified site
C84.71	Anaplastic large cell lymphoma, ALK-negative, lymph nodes of head, face and neck
C84.72	Anaplastic large cell lymphoma, ALK-negative, intrathoracic lymph nodes
C84.73	Anaplastic large cell lymphoma, ALK-negative, intra-abdominal lymph nodes
C84.73	Anaplastic large cell lymphoma, ALK-negative, lymph nodes of axilla and upper limb
C84.75	Anaplastic large cell lymphoma, ALK-negative, lymph nodes of inguinal region and lower limb
C84.76	Anaplastic large cell lymphoma, ALK-negative, intrapelvic lymph nodes
C84.77	Anaplastic large cell lymphoma, ALK-negative, spleen
C84.78	Anaplastic large cell lymphoma, ALK-negative, lymph nodes of multiple sites
C84.79	Anaplastic large cell lymphoma, ALK-negative, extranodal and solid organ sites
C84.90	Mature T/NK-cell lymphomas, unspecified site
C84.91	Mature T/NK-cell lymphomas, lymph nodes of head, face, and neck
C84.92	Mature T/NK-cell lymphomas, intrathoracic lymph nodes
C84.93	Mature T/NK-cell lymphomas, intra-abdominal lymph nodes
C84.94	Mature T/NK-cell lymphomas, lymph nodes of axilla and upper limb
C84.95	Mature T/NK-cell lymphomas, lymph nodes of inguinal region and lower limb
C84.96	Mature T/NK-cell lymphomas, intrapelvic lymph nodes
C84.97	Mature T/NK-cell lymphomas, spleen
C84.98	Mature T/NK-cell lymphomas, lymph nodes of multiple sites
C84.99	Mature T/NK-cell lymphomas, extranodal and solid organ sites
C84.Z0	Other mature T/NK-cell lymphomas, unspecified site
C84.Z1	Other mature T/NK-cell lymphomas, lymph nodes of head, face, and neck
C84.Z2	Other mature T/NK-cell lymphomas, intrathoracic lymph nodes
C84.Z3	Other mature T/NK-cell lymphomas, intra-abdominal lymph nodes
C84.Z4	Other mature T/NK-cell lymphomas, lymph nodes of axilla and upper limb
C84.Z5	Other mature T/NK-cell lymphomas, lymph nodes of inguinal region and lower limb
C84.Z6	Other mature T/NK-cell lymphomas, intrapelvic lymph nodes
C84.Z7	Other mature T/NK-cell lymphomas, spleen

ICD-10	ICD-10 Description
C84.Z8	Other mature T/NK-cell lymphomas, lymph nodes of multiple sites
C84.Z9	Other mature T/NK-cell lymphomas, extranodal and solid organ sites
C85.10	Unspecified B-cell lymphoma unspecified site
C85.11	Unspecified B-cell lymphoma lymph nodes of head, face, and neck
C85.12	Unspecified B-cell lymphoma intrathoracic lymph nodes
C85.13	Unspecified B-cell lymphoma intra-abdominal lymph nodes
C85.14	Unspecified B-cell lymphoma lymph nodes of axilla and upper limb
C85.15	Unspecified B-cell lymphoma lymph nodes of inguinal region and lower limb
C85.16	Unspecified B-cell lymphoma intrapelvic lymph nodes
C85.17	Unspecified B-cell lymphoma spleen
C85.18	Unspecified B-cell lymphoma lymph nodes of multiple sites
C85.19	Unspecified B-cell lymphoma extranodal and solid organ sites
C85.20	Mediastinal (thymic) large B-cell lymphoma, unspecified site
C85.21	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck
C85.22	Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes
C85.23	Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes
C85.24	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb
C85.25	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.26	Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes
C85.27	Mediastinal (thymic) large B-cell lymphoma, spleen
C85.28	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites
C85.29	Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites
C85.80	Other specified types of non-Hodgkin lymphoma, unspecified site
C85.81	Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face and neck
C85.82	Other specified types of non-Hodgkin lymphoma, intrathoracic lymph nodes
C85.83	Other specified types of non-Hodgkin lymphoma, intra-abdominal lymph nodes
C85.84	Other specified types of non-Hodgkin lymphoma, lymph nodes of axilla and upper limb
C85.85	Other specified types of non-Hodgkin lymphoma, lymph nodes of inguinal region of lower limb
C85.86	Other specified types of non-Hodgkin lymphoma, intrapelvic lymph nodes
C85.87	Other specified types of non-Hodgkin lymphoma, spleen
C85.88	Other specified types of non-Hodgkin lymphoma, lymph nodes of multiple sites
C85.89	Other specified types of non-Hodgkin lymphoma, extranodal and solid organ sites
C86.0	Extranodal NK/T-cell lymphoma, nasal type
C86.1	Hepatosplenic T-cell lymphoma

ICD-10	ICD-10 Description
C86.2	Enteropathy-type (intestinal) T-cell lymphoma
C86.5	Angioimmunoblastic T-cell lymphoma
C86.6	Primary cutaneous CD30-positive T-cell proliferations
C91.50	Adult T-cell lymphoma/leukemia (HTLV-1-associated) not having achieved remission
C91.51	Adult T-cell lymphoma/leukemia (HTLV-1-associated) in remission
C91.52	Adult T-cell lymphoma/leukemia (HTLV-1-associated) in relapse
D47.Z1	Post-transplant lymphoproliferative disorder (PTLD)
Z85.71	Personal history of Hodgkin lymphoma

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

Jurisdiction(s): J&M	NCD/LCD/LCA Document (s): A56141
https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A56141&bc=gAAAAAAAAAAAA	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC