

Jakafi® (ruxolitinib) (Oral)

Document Number: IC-0073

Last Review Date 10/26/2020

Date of Origin: 03/01/2012

Dates Reviewed: 12/2012, 03/2013, 02/2014, 08/2014, 12/2014, 10/2015, 10/2016, 10/2017, 10/2018, 06/2019, 11/2020

I. Length of Authorization

Coverage will be provided for six months and may be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- 5 mg tablets: 2 tablets per day
- 10 mg tablets: 2 tablets per day
- 15 mg tablets: 2 tablets per day
- 20 mg tablets: 2 tablets per day
- 25 mg tablets: 2 tablets per day

B. Max Units (per dose and over time) [HCPCS Unit]:

Acute GVHD

- 20 mg per day

All other indications

- 50 mg per day

III. Initial Approval Criteria ¹

- Patient is at least 18 years of age (unless otherwise specified); AND

Universal Criteria ¹

- Patient does not have an active infection, including clinically important localized infections; AND
- Patient will avoid concomitant therapy with all of the following:
 - Coadministration with fluconazole, or if therapy is unavoidable, the patient will be monitored closely for adverse reaction and/or dose modifications will be implemented; AND

- Coadministration with strong CYP3A4 inhibitors (e.g., clarithromycin, ketoconazole, nefazodone, etc.), or if therapy is unavoidable, the patient will be monitored closely for adverse reaction and/or dose modifications will be implemented; **AND**

Myelofibrosis (MF) (including primary, post-polycythemia vera and post-essential thrombocythemia MF) † Φ^{1,4,13}

- Patient has symptomatic low- to intermediate-1-risk disease ‡ ; **OR**
- Patient has intermediate-2 or high-risk disease† ; **AND**
 - Starting platelet count (<30 days old) is $\geq 50 \times 10^9/L$; **OR**
- Patient has accelerated phase or blast phase/acute myeloid leukemia ‡

Polycythemia Vera † Φ^{1,4,13}

- Patient has had an inadequate response (or intolerance) to a 3 month or longer trial of hydroxyurea † or interferon therapy ‡; **AND**
 - Patient has symptomatic low risk disease with indications for cytoreductive therapy; **OR**
 - Patient has high risk disease

Essential Thrombocythemia ‡^{4,13}

- Patient has had an inadequate response or loss of response to hydroxyurea, interferon therapy, or anagrelide

Graft Versus Host Disease (GVHD)^{1,4,5,14,15}

- Used for disease related to allogeneic hematopoietic stem cell transplantation; **AND**
- Used in combination with systemic corticosteroids for steroid-refractory disease; **AND**
 - Patient has acute graft versus host disease (aGVHD) † Φ; **AND**
 - Patient is at least 12 years of age; **OR**
 - Patient has chronic graft versus host disease (cGVHD) ‡

Myelodysplastic/Myeloproliferative Overlap Neoplasms (MDS/MPN)^{4,7,9}

- Used in combination with a hypomethylating agent (e.g., decitabine, azacitidine, etc.) for chronic myelomonocytic leukemia (CMML)-2; **OR**
- Used as a single agent or in combination with a hypomethylating agent (e.g., decitabine, azacitidine, etc.) for BCR-ABL negative atypical chronic myeloid leukemia (aCML)

Myeloid/Lymphoid Neoplasms with Eosinophilia ‡^{4,10}

- Used in combination with ALL- or AML-type induction chemotherapy followed by allogeneic HCT (if eligible); **AND**
 - Patient has lymphoid, myeloid, or mixed lineage neoplasm; **AND**
 - Patient has JAK2 rearrangement in blast phase; **OR**

- Patient has myeloid or lymphoid neoplasms; **AND**
 - Patient has JAK2 rearrangement in chronic phase

Pediatric Acute Lymphoblastic Leukemia (ALL) †^{4,11,12}

- Patient is at least 1 year of age; **AND**
- Patient has Ph-like B-ALL; **AND**
 - Used as induction therapy in combination with Total Therapy XVII regimen [prednisone, vincristine, daunorubicin, pegaspargase, cyclophosphamide, cytarabine, 6-MP, intrathecal (IT) therapy (methotrexate OR cytarabine OR methotrexate, cytarabine, and corticosteroid)]; **AND**
 - Patient has disease with mutations associated with JAK-STAT pathway activation; **OR**
 - Used as consolidation therapy in combination with COG AALL1521 regimen (cyclophosphamide, cytarabine, 6-MP, vincristine, pegaspargase, IT methotrexate); **AND**
 - Patient has CRLF2+ or CRLF2- with JAK2 fusions, EPOR rearrangements, SH2B3 alterations, IL7R insertions/deletions; **OR**
 - Used as consolidation therapy in combination with the standard risk/high risk (SR/HR) arm of the Total Therapy XVII regimen [high-dose methotrexate, pegaspargase, 6-MP, intrathecal (IT) therapy (methotrexate OR cytarabine OR methotrexate, cytarabine, and corticosteroid)]; **AND**
 - Patient has with mutations associated with JAK-STAT pathway activation

† FDA Approved Indication(s); ‡ Compendia Approved Indication(s); **Φ** Orphan Drug

IV. Renewal Criteria¹

Authorizations can be renewed based on the following criteria:

- Patient continues to meet universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: serious infections (bacterial, mycobacterial, fungal, and viral), severe hematologic toxicity (neutropenia, thrombocytopenia, and anemia), non-melanoma skin cancer, lipid elevations (including total cholesterol, LDL, and triglycerides), etc.; **AND**
- **Myelofibrosis^{1,4,13}**
 - Treatment response with a decrease in spleen size or improvements in other myelofibrosis symptoms (such as fatigue, bone pain, frequent infections, fever, night sweats, easy bruising/bleeding)
- **Polycythemia Vera^{1,4,13}**

- Treatment response such as hematocrit control and/or spleen volume reduction
- **Essential Thrombocythemia** ^{4,13}
 - Disease response as evidenced by at least one of the following:
 - Decrease in spleen size or improvements in other myelofibrosis symptoms (such as fatigue, bone pain, frequent infections, fever, night sweats, easy bruising/bleeding, etc.)
 - Platelet count $\leq 400 \times 10^9/L$, WBC count $< 10 \times 10^9/L$, absence of leukoerythroblastosis
 - Absence of any signs of progressive disease or hemorrhagic or thrombotic events
- **aGVHD** ^{1,4,5}
 - Treatment response such as stabilization or improvement in disease; **AND**
 - In patients who have had a response and have discontinued therapeutic doses of corticosteroids, tapering of Jakafi should be considered
- **cGVHD** ^{4,14,15}
 - Treatment response as evidenced by stabilization or improvement in disease
- **Myelodysplastic/Myeloproliferative Overlap Neoplasms (MDS/MPN)** ^{4,7-9}
 - Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread
- **Myeloid/Lymphoid Neoplasms with Eosinophilia** ^{4,10}
 - Disease response as evidenced by at least one of the following:
 - Decrease in spleen size or improvements in other myelofibrosis symptoms (such as fatigue, bone pain, frequent infections, fever, night sweats, easy bruising/bleeding, etc.)
 - Stabilization or improvement as evidenced by a complete response [CR] (i.e. morphologic, cytogenetic or molecular complete response CR), complete hematologic response or a partial response by CBC, bone marrow cytogenetic analysis, QPCR, or FISH
- **Pediatric Acute Lymphoblastic Leukemia (ALL)** ^{4,11,12}
 - Treatment response or stabilization of disease as indicated by CBC, bone marrow cytogenetic analysis, QPCR, or FISH

V. Dosage/Administration ^{1,8,9,12,14,15}

Indication	Dose
Myelofibrosis & Myeloid/Lymphoid Neoplasms with Eosinophilia	<ul style="list-style-type: none"> ▪ Platelets $>200 \times 10^9/L$ – Starting dose is 20mg orally twice daily ▪ Platelets $100-200 \times 10^9/L$ – Starting dose is 15mg orally twice daily ▪ Platelets $50-100 \times 10^9/L$ – Starting dose is 5mg orally twice daily <p><i>Doses may be increased in 5 mg twice daily increments to a maximum of 25 mg twice daily</i></p>

Polycythemia Vera & Essential Thrombocythemia	<ul style="list-style-type: none"> ▪ 10 mg orally twice daily. <p><i>Doses may be titrated based on safety and efficacy up to a maximum of 25 mg twice daily.</i></p>
aGVHD	<ul style="list-style-type: none"> • Starting dose is 5 mg orally twice daily. Consider increasing the dose to 10 mg twice daily after at least 3 days of treatment if the ANC and platelet counts are not decreased by 50% or more relative to the first day of dosing with Jakafi. • Tapering of Jakafi may be considered after 6 months of treatment in patients with response who have discontinued therapeutic doses of corticosteroids. Taper Jakafi by one dose level approximately every 8 weeks (10 mg twice daily to 5 mg twice daily to 5 mg once daily). If acute GVHD signs or symptoms recur during or after the taper of Jakafi, consider retreatment.
cGVHD	<ul style="list-style-type: none"> • 5 mg orally twice daily <p><i>Doses may be titrated based on safety and efficacy up to a maximum of 10 mg twice daily.</i></p>
MDS/MPN	<ul style="list-style-type: none"> ▪ Platelets >200 X 10⁹/L – Starting dose is 20mg orally twice daily ▪ Platelets 100-200 X 10⁹/L –Starting dose is 15mg orally twice daily ▪ Platelets 50-100 X 10⁹/L –Starting dose is 5mg orally twice daily <p><i>Doses may be titrated based on safety and efficacy up to a maximum of 20 mg twice daily.</i></p>
Pediatric ALL	<ul style="list-style-type: none"> ▪ 40 mg/m² orally twice daily

VI. Billing Code/Availability Information

HCPCS Code:

- J8999 – Prescription drug, oral, chemotherapeutic, not otherwise specified

NDC:

- 5 mg tablets: 50881-0005-xx
- 10 mg tablets: 50881-0010-xx
- 15 mg tablets: 50881-0015-xx
- 20 mg tablets: 50881-0020-xx
- 25 mg tablets: 50881-0025-xx

VII. References

1. Jakafi [package insert]. Wilmington, DE; Incyte; January 2020. Accessed October 2020.
2. Tefferi A. Primary myelofibrosis: 2013 update on diagnosis, risk-stratification, and management. *Am J Hematol.* 2013 Feb; 88(2):141-50.
3. Reilly JT, McMullin MF, Beer PA, et al. Guideline for the diagnosis and management of myelofibrosis. *Br J Haematol* 2012; 158:453.
4. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) ruxolitinib oral. National Comprehensive Cancer Network, 2020. The

NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed October 2020.

5. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hematopoietic Cell Transplantation (HCT). Version 2.2020. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed October 2020.
6. Zeiser R, Burchert A, Lengerke C, et al. Ruxolitinib in corticosteroid-refractory graft-versus-host disease after allogeneic stem cell transplantation: a multicenter survey. *Leukemia*. 2015 Oct;29(10):2062-8. doi: 10.1038/leu.2015.212.
7. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Myelodysplastic Syndromes. Version 1.2021. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed October 2020.
8. Padron E, Dezern A, Andrade-Campos M, et al; Myelodysplastic Syndrome Clinical Research Consortium. A Multi-Institution Phase I Trial of Ruxolitinib in Patients with Chronic Myelomonocytic Leukemia (CMML). *Clin Cancer Res*. 2016 Aug 1;22(15):3746-54. doi: 10.1158/1078-0432.CCR-15-2781.
9. Assi R, Kantarjian HM, Garcia-Manero G, et al. A phase II trial of ruxolitinib in combination with azacytidine in myelodysplastic syndrome/myeloproliferative neoplasms. *Am J Hematol*. 2018 Feb;93(2):277-285. doi: 10.1002/ajh.24972.
10. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Fusion Genes Version 3.2021. National Comprehensive Cancer Network, 2020. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2020.
11. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Pediatric Acute Lymphoblastic Leukemia Version 1.2021. National Comprehensive Cancer Network, 2020. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2020.

12. Tasian SK, Assad A, Hunter DS, et al. A Phase 2 Study of Ruxolitinib with Chemotherapy in Children with Philadelphia Chromosome-like Acute Lymphoblastic Leukemia (INCB18424-269/AALL1521): Dose-Finding Results from the Part 1 Safety Phase. *Blood* 2018; 132 (Supplement 1): 555.
13. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Myeloproliferative Neoplasms Version 1.2020. National Comprehensive Cancer Network, 2020. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2020.
14. Modi B, Hernandez-Henderson M, Yang D, et al. Ruxolitinib as Salvage Therapy for Chronic Graft-versus-Host Disease. *Biol Blood Marrow Transplant*. 2019 Feb;25(2):265-269. doi: 10.1016/j.bbmt.2018.09.003.
15. Khoury HJ, Langston AA, Kota VK, et al. Ruxolitinib: a steroid sparing agent in chronic graft-versus-host disease. *Bone Marrow Transplant*. 2018 Jul;53(7):826-831. doi: 10.1038/s41409-017-0081-5.
16. Verstovsek S, Mesa RA, Gotlib J, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med*. 2012 Mar 1;366(9):799-807. doi: 10.1056/NEJMoa1110557.
17. Harrison C, Kiladjan JJ, Al-Ali HK, et al. JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med*. 2012 Mar 1;366(9):787-98. doi: 10.1056/NEJMoa1110556.
18. Vannucchi AM, Kiladjan JJ, Griesshammer M, et al. Ruxolitinib versus standard therapy for the treatment of polycythemia vera. *N Engl J Med*. 2015 Jan 29;372(5):426-35. doi: 10.1056/NEJMoa1409002.
19. Jagasia M, Perales MA, Schroeder MA, et al. Ruxolitinib for the treatment of steroid-refractory acute GVHD (REACH1): a multicenter, open-label phase 2 trial. *Blood*. 2020 May 14;135(20):1739-1749. doi: 10.1182/blood.2020004823.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C91.00	Acute lymphoblastic leukemia not having achieved remission
C92.20	Atypical chronic myeloid leukemia, BCR/ABL-negative, not having achieved remission
C92.22	Atypical chronic myeloid leukemia, BCR/ABL-negative, in relapse
C93.10	Chronic lymphocytic leukemia of B-cell type not having achieved remission
C93.12	Chronic lymphocytic leukemia of B-cell type in relapse
C94.40	Acute panmyelosis with myelofibrosis not having achieved remission
C94.41	Acute panmyelosis with myelofibrosis in remission
C94.42	Acute panmyelosis with myelofibrosis in relapse
C94.6	Myelodysplastic disease, not classified

C94.8	Other specified leukemias
C94.80	Other specified leukemias not having achieved remission
C94.81	Other specified leukemias, in remission
C94.82	Other specified leukemias, in relapse
C95.1	Chronic leukemia of unspecified cell type
C95.10	Chronic leukemia of unspecified cell type not having achieved remission
C95.11	Chronic leukemia of unspecified cell type, in remission
C95.12	Chronic leukemia of unspecified cell type, in relapse
C96.Z	Other specified malignant neoplasms of lymphoid, hematopoietic and related tissue
C96.9	Malignant neoplasm of lymphoid, hematopoietic and related tissue, unspecified
D45	Polycythemia vera
D47.1	Chronic myeloproliferative disease
D47.3	Essential (hemorrhagic) thrombocythemia
D47.4	Osteomyelofibrosis
D75.81	Myelofibrosis
D89.810	Acute graft-versus-host disease
D89.811	Chronic graft-versus-host disease
D89.812	Acute on chronic graft-versus-host disease
D89.813	Graft-versus-host disease, unspecified
T86.09	Other complications of bone marrow transplant
Z94.81	Bone marrow transplant status

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)

Medicare Part B Administrative Contractor (MAC) Jurisdictions

Jurisdiction	Applicable State/US Territory	Contractor
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC