



## **Remicade® (infliximab) (Intravenous)**

Document Number: SHP-0104

Last Review Date: 07/01/2019

Date of Origin: 07/20/2010

Dates Reviewed: 09/2010, 12/2012, 2/2011, 03/2011, 06/2011, 09/2011, 10/2011, 12/2011, 03/2012, 06/2012, 09/2012, 11/2012, 12/2012, 03/2013, 06/2013, 09/2013, 12/2013, 03/2014, 06/2014, 09/2014, 12/2014, 03/2015, 06/2015, 09/2015, 12/2015, 03/2016, 06/2016, 09/2016, 12/2016, 03/2017, 06/2017, 09/2017, 12/2017, 03/2018, 06/2018, 10/2018, 04/2019, 07/2019

### **I. Length of Authorization**

Coverage will be provided for 6 months and may be renewed.

Therapy for the management of Immune-Checkpoint Inhibitor related toxicity may not be renewed.

### **II. Dosing Limits**

#### **A. Quantity Limit (max daily dose) [Pharmacy Benefit]:**

**Loading Dose:**

Remicade 100 mg single-use vial: 6 vials at weeks 0, 2, 6 (18 vials total)

**Maintenance Dose:**

Remicade 100 mg single-use vial: 10 vials every 4 weeks

#### **B. Max Units (per dose and over time) [Medical Benefit]:**

**Loading Dose:**

**Rheumatoid Arthritis (RA)**

- 40 billable units at weeks 0, 2, 6

**All other indications (except management of immune checkpoint inhibitor toxicity)**

- 60 billable units at weeks 0, 2, 6

**Management of immune checkpoint inhibitor toxicity**

- 60 billable units at weeks 0,2

**Maintenance Dose:**

**Rheumatoid Arthritis (RA)**

- 100 billable units every 4 weeks

**Crohn's Disease/Ulcerative Colitis**

- 100 billable units every 8 weeks

**Ankylosing spondylitis**

- 60 billable units every 6 weeks

**All other indications (except management of immune checkpoint inhibitor toxicity)**

- 60 billable units every 8 weeks

### III. Initial Approval Criteria

- Renflexis is the preferred infliximab product. Patients must have failed, or have a contraindication, or intolerance to Renflexis prior to consideration of Remicade or Inflectra.
- Patients are allowed to continue treatment with Remicade or Inflectra that was authorized prior to January 1<sup>st</sup>, 2019 until the end of their Remicade/Inflectra authorization period. After the Remicade/Inflectra authorization expires, patients must switch to the preferred biosimilar, unless contraindicated.

Coverage is provided in the following conditions:

- Patient has been evaluated and screened for the presence of latent TB infection prior to initiating treatment; **AND**
- Patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment; **AND**
- Patient does not have an active infection, including clinically important localized infections; **AND**
- Must not be administered concurrently with live vaccines; **AND**
- Patient is not on concurrent treatment with another TNF-inhibitor, biologic response modifier or other non-biologic agent (i.e., apremilast tofacitinib, baricitinib); **AND**
- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**

#### Crohn's Disease †

- Must be prescribed by, or in consultation with, a specialist in gastroenterology; **AND**
- Adult patient (18 years or older); **AND**
- Documented moderate to severe disease; **AND**
- Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum (3) month trial of corticosteroids or immunomodulators (e.g., azathioprine, 6-mercaptopurine, or methotrexate)

#### Pediatric Crohn's Disease †

- Must be prescribed by, or in consultation with, a specialist in gastroenterology; **AND**
- Patient is at least 6 years of age; **AND**
- Documented moderate to severe disease; **AND**
- Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum (3) month trial of corticosteroids or immunomodulators (e.g., azathioprine, etc.)

#### Ulcerative Colitis †

- Must be prescribed by, or in consultation with, a specialist in gastroenterology; **AND**
- Adult patient (18 years or older); **AND**
- Documented moderate to severe disease; **AND**

---

#### REMICADE® (infliximab) Prior Auth Criteria

- Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum (3) month trial of corticosteroids or immunomodulators (e.g., azathioprine, 6-mercaptopurine, or methotrexate)

### Pediatric Ulcerative Colitis †

- Must be prescribed by, or in consultation with, a specialist in gastroenterology; **AND**
- Patient is at least 6 years of age; **AND**
- Documented moderate to severe disease; **AND**
- Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum (3) month trial of corticosteroids or immunomodulators (e.g., azathioprine, etc.)

### Fistulizing Crohn's Disease †

- Must be prescribed by, or in consultation with, a specialist in gastroenterology; **AND**
- Adult patient (18 years or older); **AND**
- Patient has at least one or more draining fistulas (i.e., enterovesical, enterocutaneous, enteroenteric, or enterovaginal fistulas) for at least 3 months

### Rheumatoid Arthritis (RA) †

- Must be prescribed by, or in consultation with, a specialist in rheumatology; **AND**
- Adult patient (18 years or older); **AND**
- Documented moderate to severe disease; **AND**
- Patient has had at least a 3 month trial and failed previous therapy with ONE oral disease modifying anti-rheumatic agent (DMARD) such as methotrexate, azathioprine, auranofin, hydroxychloroquine, penicillamine, sulfasalazine, or leflunomide; **AND**
- Used in combination with methotrexate (MTX) unless contraindicated; **AND**
- Previous failure with a preferred self-injectable TNF antagonist

### Psoriatic Arthritis †

- Must be prescribed by, or in consultation with, a specialist in dermatology or rheumatology; **AND**
- Adult patient (18 years or older); **AND**
- Documented moderate to severe active disease; **AND**
  - For patients with predominantly axial disease OR active enthesitis and/or dactylitis, an adequate trial and failure of at least TWO (2) non-steroidal anti-inflammatory agents (NSAIDs), unless use is contraindicated; **OR**
  - For patients with peripheral arthritis, a trial and failure of at least a 3 month trial of ONE oral disease-modifying anti-rheumatic agent (DMARD) such as methotrexate, azathioprine, sulfasalazine, or hydroxychloroquine

---

#### REMICADE® (infliximab) Prior Auth Criteria

## Ankylosing Spondylitis †

- Must be prescribed by, or in consultation with, a specialist in rheumatology; **AND**
- Adult patient (18 years or older); **AND**
- Documented active disease; **AND**
- Patient had an adequate trial and failure of at least TWO (2) non-steroidal anti-inflammatory agents (NSAIDs), unless use is contraindicated

## Plaque Psoriasis †

- Must be prescribed by, or in consultation with, a specialist in dermatology or rheumatology; **AND**
- Adult patient (18 years or older); **AND**
- Documented moderate to severe plaque psoriasis for at least 6 months with at least one of the following:
  - Involvement of at least 10% of body surface area (BSA); **OR**
  - Psoriasis Area and Severity Index (PASI) score of 10 or greater; **OR**
  - Incapacitation due to plaque location (i.e. head and neck, palms, soles or genitalia); **AND**
- Patient did not respond adequately (or is not a candidate) to a 3 month minimum trial of topical agents (i.e., anthralin, coal tar preparations, corticosteroids, emollients, immunosuppressives, keratolytics, retinoic acid derivatives, and/or vitamin D analogues); **AND**
- Patient did not respond adequately (or is not a candidate) to a 3 month minimum trial of at least one systemic agent (i.e., immunosuppressives, retinoic acid derivatives, and/or methotrexate); **AND**
- Patient did not respond adequately (or is not a candidate) to a 3 month minimum trial of phototherapy (i.e., psoralens with UVA light (PUVA) or UVB with coal tar or dithranol)

## Uveitis Associated with Behcet's Syndrome ‡

- Must be prescribed by, or in consultation with, a specialist in rheumatology or ophthalmology; **AND**
- Patient's disease is refractory to immunosuppressive therapy (e.g., corticosteroids, etc.); **AND**
- Patient had an inadequate response to a self-administered biologic therapy (e.g., adalimumab)

## Management of Immune Checkpoint Inhibitor related Toxicity ‡

- Patient has been receiving therapy with an immune checkpoint inhibitor (e.g. nivolumab, pembrolizumab, atezolizumab, avelumab, durvalumab, cemiplimab, ipilimumab, etc.); **AND**
- Patient has one of the following toxicities related to their immunotherapy:
  - Moderate (grade 2) to severe (grade 3-4) diarrhea or colitis

- Severe (grade 3-4) pneumonitis refractory to methylprednisolone after 48 hours of therapy
- Severe (grade 3) or life-threatening (grade 4) renal failure or elevated serum creatinine that is refractory to at least 1 week of therapy with corticosteroids
- Uveitis (grade 3-4) that is refractory to high-dose systemic corticosteroids
- Life-threatening (grade 4) myocarditis, pericarditis, arrhythmias or impaired ventricular function if no improvement within 24 hours of starting pulse-dose methylprednisolone
- Severe inflammatory arthritis as additional disease-modifying therapy refractory to high-dose corticosteroids after 14 days of treatment

† FDA Approved Indication(s); ‡ Compendia recommended indication(s)

#### IV. Renewal Criteria

- Patients are allowed to continue treatment with Remicade or Inflectra that was authorized prior to January 1<sup>st</sup>, 2019 until the end of their Remicade/Inflectra authorization period. After the Remicade/Inflectra authorization expires, patients must switch to the preferred biosimilar, unless contraindicated.

Coverage can be renewed based upon the following criteria:

- Patient continues to meet criteria identified in section III; AND
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: severe hypersensitivity reactions, malignancy, significant hematologic abnormalities, serious infections, cerebrovascular accidents, cardiotoxicity/heart failure, neurotoxicity, hepatotoxicity, lupus-like syndrome, demyelinating disease, etc.; AND
- Patient is receiving ongoing monitoring for presence of TB or other active infections; AND

#### Crohn's Disease

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as endoscopic activity, number of liquid stools, presence and severity of abdominal pain, presence of abdominal mass, body weight compared to IBW, hematocrit, presence of extra-intestinal complications, tapering or discontinuation of corticosteroid therapy, use of anti-diarrheal drugs, and/or an improvement on a disease activity scoring tool [e.g. an improvement on the Crohn's Disease Activity Index (CDAI) score or the Harvey-Bradshaw Index score.]

#### Pediatric Crohn's Disease

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as endoscopic activity, number of liquid stools, presence and severity of abdominal pain, presence of abdominal mass, body weight compared to IBW, hematocrit, presence of extra-intestinal complications, tapering or discontinuation of corticosteroid therapy, use of anti-diarrheal drugs and/or an improvement on a disease activity scoring tool [e.g. an improvement on the Pediatric Crohn's Disease Activity Index (PCDAI) score or the Harvey-Bradshaw Index score.]

#### Ulcerative Colitis

##### REMICADE® (infliximab) Prior Auth Criteria

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as stool frequency, rectal bleeding, and/or endoscopic activity, tapering or discontinuation of corticosteroid therapy, and/or an improvement on a disease activity scoring tool [e.g. an improvement on the Ulcerative Colitis Endoscopic Index of Severity (UCEIS) score or the Mayo Score].

### Pediatric Ulcerative Colitis

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as stool frequency, rectal bleeding, and/or endoscopic activity, tapering or discontinuation of corticosteroid therapy, and/or an improvement on a disease activity scoring tool [e.g. an improvement on the Pediatric Ulcerative Colitis Activity Index (PUCAI) score or the Mayo Score].

### Fistulizing Crohn's Disease

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as a reduction in number of enterocutaneous fistulas draining upon gentle compression, and/or an improvement on a disease activity scoring tool [e.g. an improvement on the Crohn's Disease Activity Index (CDAI) score or the Harvey-Bradshaw Index score].

### Psoriatic Arthritis

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts and/or an improvement on a disease activity scoring tool [e.g. defined as an improvement in at least 2 of the 4 Psoriatic Arthritis Response Criteria (PsARC), 1 of which must be joint tenderness or swelling score, with no worsening in any of the 4 criteria.]

### Rheumatoid Arthritis

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts, reduction of C-reactive protein, improvement of patient global assessment, and/or an improvement on a disease activity scoring tool [e.g. an improvement on a composite scoring index such as Disease Activity Score-28 (DAS28) of 1.2 points or more or a  $\geq 20\%$  improvement on the American College of Rheumatology-20 (ACR20) criteria].

### Ankylosing Spondylitis

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as total back pain, physical function, morning stiffness, and/or an improvement on a disease activity scoring tool [e.g.  $\geq 1.1$  improvement on the Ankylosing Spondylitis Disease Activity Score (ASDAS) or an improvement of  $\geq 2$  on the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI)].

### Plaque Psoriasis

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as redness, thickness, scaliness, and/or the amount of surface area involvement (a total BSA involvement  $\leq 1\%$ ), and/or an improvement on a disease activity scoring tool [e.g. a 75% reduction in the PASI score from when treatment started (PASI 75) or a 50% reduction in the PASI score (PASI 50) and a four-point reduction in the DLQI from when treatment started.]

### Uveitis Associated with Behcet's Syndrome

---

#### REMICADE® (infliximab) Prior Auth Criteria

- Disease response as indicated by an improvement in signs and symptoms compared to baseline [e.g. reduction in inflammation and/or lesions, dose reduction of oral glucocorticoids and/or immunosuppressive agents, improvement in vitreous haze, improvement in best corrected visual acuity (BCVA), disease stability and/or reduced rate of decline]

### **Management of Immune Checkpoint Inhibitor related Toxicity ‡**

- May not be renewed.

## **V. Dosage/Administration**

<b>Indication</b>	<b>Loading Doses</b>	<b>Maintenance Dosing</b>	<b>Maximum Dose &amp; Frequency</b>
Rheumatoid Arthritis	3 mg/kg at weeks 0, 2, & 6	3 mg/kg every 8 weeks thereafter	Up to 10 mg/kg every 4 weeks
Ankylosing Spondylitis	5 mg/kg at weeks 0, 2, & 6	5 mg/kg every 6 weeks thereafter	5 mg/kg every 6 weeks
Crohn's Disease & Ulcerative Colitis	5 mg/kg at weeks 0, 2, & 6	5 mg/kg every 8 weeks thereafter	Up to 10 mg/kg every 8 weeks
Psoriatic Arthritis, Plaque Psoriasis, Behçet's Uveitis	5 mg/kg at weeks 0, 2, & 6	5 mg/kg every 8 weeks thereafter	5 mg/kg every 8 weeks
Management of Immune Checkpoint Inhibitor Related Toxicity	5 mg/kg at weeks 0,2	N/A	N/A

- Dose escalation (up to the maximum dose and frequency specified above) may occur upon clinical review on a case by case basis provided that the patient has:
  - Shown an initial response to therapy; **AND**
  - Received the three loading doses at the dose **AND** interval specified above; **AND**
  - Received a minimum of one maintenance dose at the dose **AND** interval specified above; **AND**
  - Responded to therapy (by treatment week 16) with subsequent loss of response; **AND**
  - Dose escalation may either increase the dose **OR** decrease the interval provided it does not exceed the following limits:
    - Dose increase by no more than 2.5 mg/kg; **OR**
    - Interval decrease by no more than 2 weeks

Note:

- Prior to escalating doses a patient's neutralizing IFX-antibodies should be assessed and addition of a DMARD (e.g., thiopurine or MTX), if not already on such therapy, should be considered.
- Criteria for disease-specific response to therapy are noted in section IV. Patients with moderate to severe heart failure (NYHA Functional Class III/IV; LVEF ≤35%) should not receive doses in excess of 5 mg/kg.

## **VI. Billing Code/Availability Information**

JCode:

- J1745 – Injection, infliximab, excludes biosimilar, 10 mg; 1 billable unit = 10 mg

---

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

NDC:

- Remicade 100 mg single-use vial for injection: 57894-0030-xx

## VII. References

1. Remicade [package insert]. Horsham, PA; Janssen Biotech, Inc; June 2018. Accessed June 2019.
2. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res (Hoboken)*. 2015 Nov 6. doi: 10.1002/acr.22783.
3. Ward MM, Deodhar, A, Akl, EA, et al. American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network 2015 Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol*. 2015 Sep 24. doi: 10.1002/art.39298.
4. Menter A, Feldman SR, Weinstein GD, et al. A randomized comparison of continuous vs. intermittent infliximab maintenance regimens over 1 year in the treatment of moderate-to-severe plaque psoriasis. *J Am Acad Dermatol* 2007;56:31e1-15.
5. Niccoli L, Nannini C, Benucci M, Chindamo D, Cassarà E, Salvarani C, Cimino L, Gini G, Lenzetti I, Cantini F. Long-term efficacy of infliximab in refractory posterior uveitis of Behcet's disease: a 24-month follow-up study. *Rheumatology (Oxford)*. 2007 Jul;46(7):1161-4. Epub 2007 May 3.
6. Giardina A, Ferrante A, Ciccia F, et al. One year study of efficacy and safety of infliximab in the treatment of patients with ocular and neurological Behcet's disease refractory to standard immunosuppressive drugs. *Rheumatol Int* 2011;31:33–37.
7. Okada A, Goto H, Ohno S, et al. Multicenter study of infliximab for refractory uveoretinitis in Behcet disease. *Arch Ophthalmol* 2012;130(5):592-598.
8. Lichtenstein GR, Loftus EV, Isaacs KL, et al. American College of Gastroenterology. Clinical Guideline: Management of Crohn's disease in adults. *Am J Gastroenterol*. 2018;113:481-517.
9. Kornbluth, A, Sachar, DB; Practice Parameters Committee of the American College of Gastroenterology. Ulcerative colitis practice guidelines in adults: American College Of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol*. 2010 Mar;105(3):501-23.
10. Terdiman JP, Gruss CB, Heidelbaugh JJ, et al. American Gastroenterological Association Institute guideline on the use of thiopurines, methotrexate, and anti-TNF- $\alpha$  biologic drugs for the induction and maintenance of remission in inflammatory Crohn's disease. *Gastroenterology*. 2013 Dec;145(6):1459-63. doi: 10.1053/j.gastro.2013.10.047.
11. Hsu S, Papp KA, Lebwohl MG, et al. Consensus guidelines for the management of plaque psoriasis. *Arch Dermatol*. 2012 Jan;148(1):95-102.
12. Gottlieb A, Korman NJ, Gordon KB, Feldman SR, Lebwohl M, Koo JY, Van Voorhees AS, Elmets CA, Leonardi CL, Beutner KR, Bhushan R, Menter A. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. *J Am Acad Dermatol* 2008 May;58(5):851-64.

13. National Institute for Health and Clinical Excellence (NICE). Adalimumab, etanercept, infliximab, rituximab and abatacept for the treatment of rheumatoid arthritis after the failure of a TNF inhibitor. London (UK): National Institute for Health and Clinical Excellence (NICE); 2010 Aug. 73 p. (Technology appraisal guidance; no. 195)
14. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis*. 2015 Dec 7. pii: annrheumdis-2015-208337. doi: 10.1136/annrheumdis-2015-208337.
15. Smolen JS, Landewé R, Bijlsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis*. 2017 Mar 6. pii: annrheumdis-2016-210715.
16. Van Der Heijde D, Ramiro S, Landewe R, et al. 2016 update of the ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis* doi:10.1136/annrheumdis-2016-210770.
17. Harbord M, Eliakim R, Bettenworth D, et al. Third European Evidence-based Consensus on Diagnosis and Management of Ulcerative Colitis. Part 2: Current Management. *J Crohns Colitis*. 2017 Jan 28. doi: 10.1093/ecco-jcc/jx009.
18. Jabs DA, Rosenbaum JT, Foster CS, et al. Guidelines for the use of immunosuppressive drugs in patients with ocular inflammatory disorders: recommendations of an expert panel. *Am J Ophthalmol*. 2000 Oct;130(4):492-513.
19. Levy-Clarke G, Jabs DA, Read RW, et al. Expert panel recommendations for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders. *Ophthalmology*. 2014 Mar;121(3):785-96.e3. doi: 10.1016/j.ophtha.2013.09.048.
20. National Institute for Health and Care Excellence. NICE 2012. Crohn's Disease: Management. Published 10 October 2012. Clinical Guideline [CG152]. <https://www.nice.org.uk/guidance/cg152/resources/crohns-disease-management-pdf-35109627942085> .
21. Lewis JD, Chuai S, Nessel L, et al. Use of the Non-invasive Components of the Mayo Score to Assess Clinical Response in Ulcerative Colitis. *Inflamm Bowel Dis*. 2008 Dec; 14(12): 1660–1666. doi: 10.1002/ibd.20520
22. Paine ER. Colonoscopic evaluation in ulcerative colitis. *Gastroenterol Rep (Oxf)*. 2014 Aug; 2(3): 161–168.
23. Walsh AJ, Bryant RV, Travis SPL. Current best practice for disease activity assessment in IBD. *Nature Reviews Gastroenterology & Hepatology* 13, 567–579 (2016) doi:10.1038/nrgastro.2016.128
24. Kornbluth, A, Sachar, DB; Practice Parameters Committee of the American College of Gastroenterology. Ulcerative colitis practice guidelines in adults: American College Of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol*. 2010 Mar;105(3):501-23.
25. National Institute for Health and Care Excellence. NICE 2017. Certolizumab pegol and secukinumab for treating active psoriatic arthritis after inadequate response to DMARDs. Published 24 May 2017. Technology Appraisal Guidance [TA445]. <https://www.nice.org.uk/guidance/TA445/chapter/1-Recommendations>. Accessed August 2017.

26. National Institute for Health and Care Excellence. NICE 2009. Rheumatoid Arthritis in Adults: Management. Published 25 February 2009. Clinical Guideline [CG79]. <https://www.nice.org.uk/guidance/cg79/resources/rheumatoid-arthritis-in-adults-management-pdf-975636823525>.
27. National Institute for Health and Care Excellence. NICE 2010. Adalimumab, etanercept, infliximab, rituximab and abatacept for the treatment of rheumatoid arthritis after failure of a TNF inhibitor. Published 10 October 2012. Clinical Guideline [TA195]. <https://www.nice.org.uk/guidance/ta195/resources/adalimumab-etanercept-infliximab-rituximab-and-abatacept-for-the-treatment-of-rheumatoid-arthritis-after-the-failure-of-a-tnf-inhibitor-pdf-82598558287813>.
28. Ward MM, Guthrie LC, Alba MI. Rheumatoid Arthritis Response Criteria And Patient-Reported Improvement in Arthritis Activity: Is an ACR20 Response Meaningful to Patients". *Arthritis Rheumatol.* 2014 Sep; 66(9): 2339–2343. doi: 10.1002/art.38705
29. National Institute for Health and Care Excellence. NICE 2008. Infliximab for the treatment of adults with psoriasis. Published 23 January 2008. Technology Appraisal Guidance [TA134]. <https://www.nice.org.uk/guidance/ta134/resources/infliximab-for-the-treatment-of-adults-with-psoriasis-pdf-82598193811141>.
30. Smith CH, Jabbar-Lopez ZK, Yiu ZK, et al. British Association of Dermatologists guidelines for biologic therapy for psoriasis 2017. *Br J Dermatol.* 2017 Sep;177(3):628-636. doi: 10.1111/bjd.15665.
31. Jabs DA, Rosenbaum JT, Foster CS, et al. Guidelines for the use of immunosuppressive drugs in patients with ocular inflammatory disorders: recommendations of an expert panel. *Am J Ophthalmol.* 2000 Oct;130(4):492-513.
32. Levy-Clarke G, Jabs DA, Read RW, et al. Expert panel recommendations for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders. *Ophthalmology.* 2014 Mar;121(3):785-96.e3. doi: 10.1016/j.ophtha.2013.09.048.
33. Minor DR, Chin K, Sashani-Sabet M, et al. Infliximab in the treatment of anti-CTLA4 antibody (Ipilimumab) induced immune related colitis. *Cancer Biotherapy & Radiopharmaceuticals.* 2009 June; 24 (3). <https://doi.org/10.1089/cbr.2008.0607>
34. Villadolid J, Amin A. Immune checkpoint inhibitors in clinical practice: update on management of immune-related toxicities. *Translational Lung Cancer Research.* 2015;4(5):560-575. doi:10.3978/j.issn.2218-6751.2015.06.06.Appendix 1 – Covered Diagnosis Codes
35. Lichtenstein GR, Loftus EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol* 2018; 113:481–517; doi: 10.1038/ajg.2018.27
36. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis & Rheumatology.* Vol. 0, No. 0, Month 2018, pp 1–28 DOI 10.1002/art.40726
37. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019 Feb 13. pii: S0190-9622(18)33001-9. <https://doi.org/10.1016/j.jaad.2018.11.057>.
38. American Academy of Dermatology Work Group, Menter A, Korman NJ, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: section 6. Guidelines of care for

- the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011 Jul;65(1):137-74.
39. Wisconsin Physicians Service Insurance Corporation. Local Coverage Determination (LCD): Drugs and Biologics (Non-chemotherapy) (L34741). Centers for Medicare & Medicaid Services, Inc. Updated on 05/24/2018 with effective date 06/1/2018. Accessed March 2019.
  40. First Coast Service Options, Inc. Local Coverage Determination (LCD): Infliximab (Remicade™) (L33704). Centers for Medicare & Medicaid Services, Inc. Updated on 01/04/2019 with effective date 01/01/2019. Accessed March 2019.
  41. National Government Services, Inc. Local Coverage Article: Infliximab, Infliximab-dyyb, Infliximab-abda (e.g., Remicade™, Inflectra™, Renflexis) – Related to LCD L33394 (A52423). Centers for Medicare & Medicaid Services, Inc. Updated on 4/27/2018 with effective date 4/1/2018. Accessed March 2019.
  42. Palmetto GBA. Local Coverage Determination (LCD): Infliximab (L35677). Centers for Medicare & Medicaid Services, Inc. Updated on 02/01/2019 with effective date 02/07/2019. Accessed March 2019.
  43. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Infliximab. National Comprehensive Cancer Network, 2019. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2019.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
H30.891	Other chorioretinal inflammations, right eye
H30.892	Other chorioretinal inflammations, left eye
H30.893	Other chorioretinal inflammations, bilateral
H30.899	Other chorioretinal inflammations, unspecified eye
H30.90	Unspecified chorioretinal inflammation, unspecified eye
H30.91	Unspecified chorioretinal inflammation, right eye
H30.92	Unspecified chorioretinal inflammation, left eye
H30.93	Unspecified chorioretinal inflammation, bilateral
I30.8	Other forms of acute pericarditis
I30.9	Acute pericarditis, unspecified
I40.8	Other acute myocarditis
I40.9	Acute myocarditis, unspecified
I50.9	Heart failure, unspecified
J70.2	Acute drug-induced interstitial lung disorders

### REMICADE® (infliximab) Prior Auth Criteria

<b>ICD-10</b>	<b>ICD-10 Description</b>
J70.4	Drug-induced interstitial lung disorders, unspecified
K50.00	Crohn's disease of small intestine without complications
K50.011	Crohn's disease of small intestine with rectal bleeding
K50.012	Crohn's disease of small intestine with intestinal obstruction
K50.013	Crohn's disease of small intestine with fistula
K50.014	Crohn's disease of small intestine with abscess
K50.018	Crohn's disease of small intestine with other complication
K50.019	Crohn's disease of small intestine with unspecified complications
K50.10	Crohn's disease of large intestine without complications
K50.111	Crohn's disease of large intestine with rectal bleeding
K50.112	Crohn's disease of large intestine with intestinal obstruction
K50.113	Crohn's disease of large intestine with fistula
K50.114	Crohn's disease of large intestine with abscess
K50.118	Crohn's disease of large intestine with other complication
K50.119	Crohn's disease of large intestine with unspecified complications
K50.80	Crohn's disease of both small and large intestine without complications
K50.811	Crohn's disease of both small and large intestine with rectal bleeding
K50.812	Crohn's disease of both small and large intestine with intestinal obstruction
K50.813	Crohn's disease of both small and large intestine with fistula
K50.814	Crohn's disease of both small and large intestine with abscess
K50.818	Crohn's disease of both small and large intestine with other complication
K50.819	Crohn's disease of both small and large intestine with unspecified complications
K50.90	Crohn's disease, unspecified, without complications
K50.911	Crohn's disease, unspecified, with rectal bleeding
K50.912	Crohn's disease, unspecified, with intestinal obstruction
K50.913	Crohn's disease, unspecified, with fistula
K50.914	Crohn's disease, unspecified, with abscess
K50.918	Crohn's disease, unspecified, with other complication
K50.919	Crohn's disease, unspecified, with unspecified complications
K51.00	Ulcerative (chronic) pancolitis without complications
K51.011	Ulcerative (chronic) pancolitis with rectal bleeding
K51.012	Ulcerative (chronic) pancolitis with intestinal obstruction
K51.013	Ulcerative (chronic) pancolitis with fistula
K51.014	Ulcerative (chronic) pancolitis with abscess

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

<b>ICD-10</b>	<b>ICD-10 Description</b>
K51.018	Ulcerative (chronic) pancolitis with other complication
K51.019	Ulcerative (chronic) pancolitis with unspecified complications
K51.20	Ulcerative (chronic) proctitis without complications
K51.211	Ulcerative (chronic) proctitis with rectal bleeding
K51.212	Ulcerative (chronic) proctitis with intestinal obstruction
K51.213	Ulcerative (chronic) proctitis with fistula
K51.214	Ulcerative (chronic) proctitis with abscess
K51.218	Ulcerative (chronic) proctitis with other complication
K51.219	Ulcerative (chronic) proctitis with unspecified complications
K51.30	Ulcerative (chronic) rectosigmoiditis without complications
K51.311	Ulcerative (chronic) rectosigmoiditis with rectal bleeding
K51.312	Ulcerative (chronic) rectosigmoiditis with intestinal obstruction
K51.313	Ulcerative (chronic) rectosigmoiditis with fistula
K51.314	Ulcerative (chronic) rectosigmoiditis with abscess
K51.318	Ulcerative (chronic) rectosigmoiditis with other complication
K51.319	Ulcerative (chronic) rectosigmoiditis with unspecified complications
K51.50	Left sided colitis without complications
K51.511	Left sided colitis with rectal bleeding
K51.512	Left sided colitis with intestinal obstruction
K51.513	Left sided colitis with fistula
K51.514	Left sided colitis with abscess
K51.518	Left sided colitis with other complication
K51.519	Left sided colitis with unspecified complications
K51.80	Other ulcerative colitis without complications
K51.811	Other ulcerative colitis with rectal bleeding
K51.812	Other ulcerative colitis with intestinal obstruction
K51.813	Other ulcerative colitis with fistula
K51.814	Other ulcerative colitis with abscess
K51.818	Other ulcerative colitis with other complication
K51.819	Other ulcerative colitis with unspecified complications
K51.90	Ulcerative colitis, unspecified, without complications
K51.911	Ulcerative colitis, unspecified with rectal bleeding
K51.912	Ulcerative colitis, unspecified with intestinal obstruction
K51.913	Ulcerative colitis, unspecified with fistula

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

<b>ICD-10</b>	<b>ICD-10 Description</b>
K51.914	Ulcerative colitis, unspecified with abscess
K51.918	Ulcerative colitis, unspecified with other complication
K51.919	Ulcerative colitis, unspecified with unspecified complications
K52.1	Toxic gastroenteritis and colitis
L40.0	Psoriasis vulgaris
L40.50	Arthropathic psoriasis, unspecified
L40.51	Distal interphalangeal psoriatic arthropathy
L40.52	Psoriatic arthritis mutilans
L40.53	Psoriatic spondylitis
L40.59	Other psoriatic arthropathy
M05.10	Rheumatoid lung disease with rheumatoid arthritis of unspecified site
M05.111	Rheumatoid lung disease with rheumatoid arthritis of right shoulder
M05.112	Rheumatoid lung disease with rheumatoid arthritis of left shoulder
M05.119	Rheumatoid lung disease with rheumatoid arthritis of unspecified shoulder
M05.121	Rheumatoid lung disease with rheumatoid arthritis of right elbow
M05.122	Rheumatoid lung disease with rheumatoid arthritis of left elbow
M05.129	Rheumatoid lung disease with rheumatoid arthritis of unspecified elbow
M05.131	Rheumatoid lung disease with rheumatoid arthritis of right wrist
M05.132	Rheumatoid lung disease with rheumatoid arthritis of left wrist
M05.139	Rheumatoid lung disease with rheumatoid arthritis of unspecified wrist
M05.141	Rheumatoid lung disease with rheumatoid arthritis of right hand
M05.142	Rheumatoid lung disease with rheumatoid arthritis of left hand
M05.149	Rheumatoid lung disease with rheumatoid arthritis of unspecified hand
M05.151	Rheumatoid lung disease with rheumatoid arthritis of right hip
M05.152	Rheumatoid lung disease with rheumatoid arthritis of left hip
M05.159	Rheumatoid lung disease with rheumatoid arthritis of unspecified hip
M05.161	Rheumatoid lung disease with rheumatoid arthritis of right knee
M05.162	Rheumatoid lung disease with rheumatoid arthritis of left knee
M05.169	Rheumatoid lung disease with rheumatoid arthritis of unspecified knee
M05.171	Rheumatoid lung disease with rheumatoid arthritis of right ankle and foot
M05.172	Rheumatoid lung disease with rheumatoid arthritis of left ankle and foot
M05.179	Rheumatoid lung disease with rheumatoid arthritis of unspecified ankle and foot
M05.19	Rheumatoid lung disease with rheumatoid arthritis of multiple sites
M05.20	Rheumatoid vasculitis with rheumatoid arthritis of unspecified site

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

ICD-10	ICD-10 Description
M05.211	Rheumatoid vasculitis with rheumatoid arthritis of right shoulder
M05.212	Rheumatoid vasculitis with rheumatoid arthritis of left shoulder
M05.219	Rheumatoid vasculitis with rheumatoid arthritis of unspecified shoulder
M05.221	Rheumatoid vasculitis with rheumatoid arthritis of right elbow
M05.222	Rheumatoid vasculitis with rheumatoid arthritis of left elbow
M05.229	Rheumatoid vasculitis with rheumatoid arthritis of unspecified elbow
M05.231	Rheumatoid vasculitis with rheumatoid arthritis of right wrist
M05.232	Rheumatoid vasculitis with rheumatoid arthritis of left wrist
M05.239	Rheumatoid vasculitis with rheumatoid arthritis of unspecified wrist
M05.241	Rheumatoid vasculitis with rheumatoid arthritis of right hand
M05.242	Rheumatoid vasculitis with rheumatoid arthritis of left hand
M05.249	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hand
M05.251	Rheumatoid vasculitis with rheumatoid arthritis of right hip
M05.252	Rheumatoid vasculitis with rheumatoid arthritis of left hip
M05.259	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hip
M05.261	Rheumatoid vasculitis with rheumatoid arthritis of right knee
M05.262	Rheumatoid vasculitis with rheumatoid arthritis of left knee
M05.269	Rheumatoid vasculitis with rheumatoid arthritis of unspecified knee
M05.271	Rheumatoid vasculitis with rheumatoid arthritis of right ankle and foot
M05.272	Rheumatoid vasculitis with rheumatoid arthritis of left ankle and foot
M05.279	Rheumatoid vasculitis with rheumatoid arthritis of unspecified ankle and foot
M05.29	Rheumatoid vasculitis with rheumatoid arthritis of multiple sites
M05.30	Rheumatoid heart disease with rheumatoid arthritis of unspecified site
M05.311	Rheumatoid heart disease with rheumatoid arthritis of right shoulder
M05.312	Rheumatoid heart disease with rheumatoid arthritis of left shoulder
M05.319	Rheumatoid heart disease with rheumatoid arthritis of unspecified shoulder
M05.321	Rheumatoid heart disease with rheumatoid arthritis of right elbow
M05.322	Rheumatoid heart disease with rheumatoid arthritis of left elbow
M05.329	Rheumatoid heart disease with rheumatoid arthritis of unspecified elbow
M05.331	Rheumatoid heart disease with rheumatoid arthritis of right wrist
M05.332	Rheumatoid heart disease with rheumatoid arthritis of left wrist
M05.339	Rheumatoid heart disease with rheumatoid arthritis of unspecified wrist
M05.341	Rheumatoid heart disease with rheumatoid arthritis of right hand
M05.342	Rheumatoid heart disease with rheumatoid arthritis of left hand

#### REMICADE® (infliximab) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

ICD-10	ICD-10 Description
M05.349	Rheumatoid heart disease with rheumatoid arthritis of unspecified hand
M05.351	Rheumatoid heart disease with rheumatoid arthritis of right hip
M05.352	Rheumatoid heart disease with rheumatoid arthritis of left hip
M05.359	Rheumatoid heart disease with rheumatoid arthritis of unspecified hip
M05.361	Rheumatoid heart disease with rheumatoid arthritis of right knee
M05.362	Rheumatoid heart disease with rheumatoid arthritis of left knee
M05.369	Rheumatoid heart disease with rheumatoid arthritis of unspecified knee
M05.371	Rheumatoid heart disease with rheumatoid arthritis of right ankle and foot
M05.372	Rheumatoid heart disease with rheumatoid arthritis of left ankle and foot
M05.379	Rheumatoid heart disease with rheumatoid arthritis of unspecified ankle and foot
M05.39	Rheumatoid heart disease with rheumatoid arthritis of multiple sites
M05.40	Rheumatoid myopathy with rheumatoid arthritis of unspecified site
M05.411	Rheumatoid myopathy with rheumatoid arthritis of right shoulder
M05.412	Rheumatoid myopathy with rheumatoid arthritis of left shoulder
M05.419	Rheumatoid myopathy with rheumatoid arthritis of unspecified shoulder
M05.421	Rheumatoid myopathy with rheumatoid arthritis of right elbow
M05.422	Rheumatoid myopathy with rheumatoid arthritis of left elbow
M05.429	Rheumatoid myopathy with rheumatoid arthritis of unspecified elbow
M05.431	Rheumatoid myopathy with rheumatoid arthritis of right wrist
M05.432	Rheumatoid myopathy with rheumatoid arthritis of left wrist
M05.439	Rheumatoid myopathy with rheumatoid arthritis of unspecified wrist
M05.441	Rheumatoid myopathy with rheumatoid arthritis of right hand
M05.442	Rheumatoid myopathy with rheumatoid arthritis of left hand
M05.449	Rheumatoid myopathy with rheumatoid arthritis of unspecified hand
M05.451	Rheumatoid myopathy with rheumatoid arthritis of right hip
M05.452	Rheumatoid myopathy with rheumatoid arthritis of left hip
M05.459	Rheumatoid myopathy with rheumatoid arthritis of unspecified hip
M05.461	Rheumatoid myopathy with rheumatoid arthritis of right knee
M05.462	Rheumatoid myopathy with rheumatoid arthritis of left knee
M05.469	Rheumatoid myopathy with rheumatoid arthritis of unspecified knee
M05.471	Rheumatoid myopathy with rheumatoid arthritis of right ankle and foot
M05.472	Rheumatoid myopathy with rheumatoid arthritis of left ankle and foot
M05.479	Rheumatoid myopathy with rheumatoid arthritis of unspecified ankle and foot
M05.49	Rheumatoid myopathy with rheumatoid arthritis of multiple sites

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

<b>ICD-10</b>	<b>ICD-10 Description</b>
M05.50	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified site
M05.511	Rheumatoid polyneuropathy with rheumatoid arthritis of right shoulder
M05.512	Rheumatoid polyneuropathy with rheumatoid arthritis of left shoulder
M05.519	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified shoulder
M05.521	Rheumatoid polyneuropathy with rheumatoid arthritis of right elbow
M05.522	Rheumatoid polyneuropathy with rheumatoid arthritis of left elbow
M05.529	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified elbow
M05.531	Rheumatoid polyneuropathy with rheumatoid arthritis of right wrist
M05.532	Rheumatoid polyneuropathy with rheumatoid arthritis of left wrist
M05.539	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified wrist
M05.541	Rheumatoid polyneuropathy with rheumatoid arthritis of right hand
M05.542	Rheumatoid polyneuropathy with rheumatoid arthritis of left hand
M05.549	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified hand
M05.551	Rheumatoid polyneuropathy with rheumatoid arthritis of right hip
M05.552	Rheumatoid polyneuropathy with rheumatoid arthritis of left hip
M05.559	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified hip
M05.561	Rheumatoid polyneuropathy with rheumatoid arthritis of right knee
M05.562	Rheumatoid polyneuropathy with rheumatoid arthritis of left knee
M05.569	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified knee
M05.571	Rheumatoid polyneuropathy with rheumatoid arthritis of right ankle and foot
M05.572	Rheumatoid polyneuropathy with rheumatoid arthritis of left ankle and foot
M05.579	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified ankle and foot
M05.59	Rheumatoid polyneuropathy with rheumatoid arthritis of multiple sites
M05.60	Rheumatoid arthritis of unspecified site with involvement of other organs and systems
M05.611	Rheumatoid arthritis of right shoulder with involvement of other organs and systems
M05.612	Rheumatoid arthritis of left shoulder with involvement of other organs and systems
M05.619	Rheumatoid arthritis of unspecified shoulder with involvement of other organs and systems
M05.621	Rheumatoid arthritis of right elbow with involvement of other organs and systems
M05.622	Rheumatoid arthritis of left elbow with involvement of other organs and systems
M05.629	Rheumatoid arthritis of unspecified elbow with involvement of other organs and systems
M05.631	Rheumatoid arthritis of right wrist with involvement of other organs and systems
M05.632	Rheumatoid arthritis of left wrist with involvement of other organs and systems
M05.639	Rheumatoid arthritis of unspecified wrist with involvement of other organs and systems
M05.641	Rheumatoid arthritis of right hand with involvement of other organs and systems

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

<b>ICD-10</b>	<b>ICD-10 Description</b>
M05.642	Rheumatoid arthritis of left hand with involvement of other organs and systems
M05.649	Rheumatoid arthritis of unspecified hand with involvement of other organs and systems
M05.651	Rheumatoid arthritis of right hip with involvement of other organs and systems
M05.652	Rheumatoid arthritis of left hip with involvement of other organs and systems
M05.659	Rheumatoid arthritis of unspecified hip with involvement of other organs and systems
M05.661	Rheumatoid arthritis of right knee with involvement of other organs and systems
M05.662	Rheumatoid arthritis of left knee with involvement of other organs and systems
M05.669	Rheumatoid arthritis of unspecified knee with involvement of other organs and systems
M05.671	Rheumatoid arthritis of right ankle and foot with involvement of other organs and systems
M05.672	Rheumatoid arthritis of left ankle and foot with involvement of other organs and systems
M05.679	Rheumatoid arthritis of unspecified ankle and foot with involvement of other organs and systems
M05.69	Rheumatoid arthritis of multiple sites with involvement of other organs and systems
M05.70	Rheumatoid arthritis with rheumatoid factor of unspecified site without organ or systems involvement
M05.711	Rheumatoid arthritis with rheumatoid factor of right shoulder without organ or systems involvement
M05.712	Rheumatoid arthritis with rheumatoid factor of left shoulder without organ or systems involvement
M05.719	Rheumatoid arthritis with rheumatoid factor of unspecified shoulder without organ or systems involvement
M05.721	Rheumatoid arthritis with rheumatoid factor of right elbow without organ or systems involvement
M05.722	Rheumatoid arthritis with rheumatoid factor of left elbow without organ or systems involvement
M05.729	Rheumatoid arthritis with rheumatoid factor of unspecified elbow without organ or systems involvement
M05.731	Rheumatoid arthritis with rheumatoid factor of right wrist without organ or systems involvement
M05.732	Rheumatoid arthritis with rheumatoid factor of left wrist without organ or systems involvement
M05.739	Rheumatoid arthritis with rheumatoid factor of unspecified wrist without organ or systems involvement
M05.741	Rheumatoid arthritis with rheumatoid factor of right hand without organ or systems involvement
M05.742	Rheumatoid arthritis with rheumatoid factor of left hand without organ or systems involvement
M05.749	Rheumatoid arthritis with rheumatoid factor of unspecified hand without organ or systems involvement

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

ICD-10	ICD-10 Description
M05.751	Rheumatoid arthritis with rheumatoid factor of right hip without organ or systems involvement
M05.752	Rheumatoid arthritis with rheumatoid factor of left hip without organ or systems involvement
M05.759	Rheumatoid arthritis with rheumatoid factor of unspecified hip without organ or systems involvement
M05.761	Rheumatoid arthritis with rheumatoid factor of right knee without organ or systems involvement
M05.762	Rheumatoid arthritis with rheumatoid factor of left knee without organ or systems involvement
M05.769	Rheumatoid arthritis with rheumatoid factor of unspecified knee without organ or systems involvement
M05.771	Rheumatoid arthritis with rheumatoid factor of right ankle and foot without organ or systems involvement
M05.772	Rheumatoid arthritis with rheumatoid factor of left ankle and foot without organ or systems involvement
M05.779	Rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot without organ or systems involvement
M05.79	Rheumatoid arthritis with rheumatoid factor of multiple sites without organ or systems involvement
M05.80	Other rheumatoid arthritis with rheumatoid factor of unspecified site
M05.811	Other rheumatoid arthritis with rheumatoid factor of right shoulder
M05.812	Other rheumatoid arthritis with rheumatoid factor of left shoulder
M05.819	Other rheumatoid arthritis with rheumatoid factor of unspecified shoulder
M05.821	Other rheumatoid arthritis with rheumatoid factor of right elbow
M05.822	Other rheumatoid arthritis with rheumatoid factor of left elbow
M05.829	Other rheumatoid arthritis with rheumatoid factor of unspecified elbow
M05.831	Other rheumatoid arthritis with rheumatoid factor of right wrist
M05.832	Other rheumatoid arthritis with rheumatoid factor of left wrist
M05.839	Other rheumatoid arthritis with rheumatoid factor of unspecified wrist
M05.841	Other rheumatoid arthritis with rheumatoid factor of right hand
M05.842	Other rheumatoid arthritis with rheumatoid factor of left hand
M05.849	Other rheumatoid arthritis with rheumatoid factor of unspecified hand
M05.851	Other rheumatoid arthritis with rheumatoid factor of right hip
M05.852	Other rheumatoid arthritis with rheumatoid factor of left hip
M05.859	Other rheumatoid arthritis with rheumatoid factor of unspecified hip
M05.861	Other rheumatoid arthritis with rheumatoid factor of right knee
M05.862	Other rheumatoid arthritis with rheumatoid factor of left knee
M05.869	Other rheumatoid arthritis with rheumatoid factor of unspecified knee

#### REMICADE® (infliximab) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

<b>ICD-10</b>	<b>ICD-10 Description</b>
M05.871	Other rheumatoid arthritis with rheumatoid factor of right ankle and foot
M05.872	Other rheumatoid arthritis with rheumatoid factor of left ankle and foot
M05.879	Other rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot
M05.89	Other rheumatoid arthritis with rheumatoid factor of multiple sites
M05.9	Rheumatoid arthritis with rheumatoid factor, unspecified
M06.00	Rheumatoid arthritis without rheumatoid factor, unspecified site
M06.011	Rheumatoid arthritis without rheumatoid factor, right shoulder
M06.012	Rheumatoid arthritis without rheumatoid factor, left shoulder
M06.019	Rheumatoid arthritis without rheumatoid factor, unspecified shoulder
M06.021	Rheumatoid arthritis without rheumatoid factor, right elbow
M06.022	Rheumatoid arthritis without rheumatoid factor, left elbow
M06.029	Rheumatoid arthritis without rheumatoid factor, unspecified elbow
M06.031	Rheumatoid arthritis without rheumatoid factor, right wrist
M06.032	Rheumatoid arthritis without rheumatoid factor, left wrist
M06.039	Rheumatoid arthritis without rheumatoid factor, unspecified wrist
M06.041	Rheumatoid arthritis without rheumatoid factor, right hand
M06.042	Rheumatoid arthritis without rheumatoid factor, left hand
M06.049	Rheumatoid arthritis without rheumatoid factor, unspecified hand
M06.051	Rheumatoid arthritis without rheumatoid factor, right hip
M06.052	Rheumatoid arthritis without rheumatoid factor, left hip
M06.059	Rheumatoid arthritis without rheumatoid factor, unspecified hip
M06.061	Rheumatoid arthritis without rheumatoid factor, right knee
M06.062	Rheumatoid arthritis without rheumatoid factor, left knee
M06.069	Rheumatoid arthritis without rheumatoid factor, unspecified knee
M06.071	Rheumatoid arthritis without rheumatoid factor, right ankle and foot
M06.072	Rheumatoid arthritis without rheumatoid factor, left ankle and foot
M06.079	Rheumatoid arthritis without rheumatoid factor, unspecified ankle and foot
M06.08	Rheumatoid arthritis without rheumatoid factor, vertebrae
M06.09	Rheumatoid arthritis without rheumatoid factor, multiple sites
M06.4	Inflammatory polyarthropathy
M06.80	Other specified rheumatoid arthritis, unspecified site
M06.811	Other specified rheumatoid arthritis, right shoulder
M06.812	Other specified rheumatoid arthritis, left shoulder
M06.819	Other specified rheumatoid arthritis, unspecified shoulder

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

<b>ICD-10</b>	<b>ICD-10 Description</b>
M06.821	Other specified rheumatoid arthritis, right elbow
M06.822	Other specified rheumatoid arthritis, left elbow
M06.829	Other specified rheumatoid arthritis, unspecified elbow
M06.831	Other specified rheumatoid arthritis, right wrist
M06.832	Other specified rheumatoid arthritis, left wrist
M06.839	Other specified rheumatoid arthritis, unspecified wrist
M06.841	Other specified rheumatoid arthritis, right hand
M06.842	Other specified rheumatoid arthritis, left hand
M06.849	Other specified rheumatoid arthritis, unspecified hand
M06.851	Other specified rheumatoid arthritis, right hip
M06.852	Other specified rheumatoid arthritis, left hip
M06.859	Other specified rheumatoid arthritis, unspecified hip
M06.861	Other specified rheumatoid arthritis, right knee
M06.862	Other specified rheumatoid arthritis, left knee
M06.869	Other specified rheumatoid arthritis, unspecified knee
M06.871	Other specified rheumatoid arthritis, right ankle and foot
M06.872	Other specified rheumatoid arthritis, left ankle and foot
M06.879	Other specified rheumatoid arthritis, unspecified ankle and foot
M06.88	Other specified rheumatoid arthritis, vertebrae
M06.89	Other specified rheumatoid arthritis, multiple sites
M06.9	Rheumatoid arthritis, unspecified
M35.2	Behçet's disease
M45.0	Ankylosing spondylitis of multiple sites in spine
M45.1	Ankylosing spondylitis of occipito-atlanto-axial region
M45.2	Ankylosing spondylitis of cervical region
M45.3	Ankylosing spondylitis of cervicothoracic region
M45.4	Ankylosing spondylitis of thoracic region
M45.5	Ankylosing spondylitis of thoracolumbar region
M45.6	Ankylosing spondylitis lumbar region
M45.7	Ankylosing spondylitis of lumbosacral region
M45.8	Ankylosing spondylitis sacral and sacrococcygeal region
M45.9	Ankylosing spondylitis of unspecified sites in spine
N17.8	Other acute kidney failure
N17.9	Acute kidney failure, unspecified

**REMICADE® (infliximab) Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

ICD-10	ICD-10 Description
R19.7	Diarrhea, unspecified

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

### Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD):

Jurisdiction(s): 5, 8	NCD/LCD Document (s): L34741  <a href="https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L34741&amp;bc=gAAAAAAA==">https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L34741&amp;bc=gAAAAAAA==</a>
-----------------------	---

Jurisdiction(s): N, 9	NCD/LCD Document (s): L33704  <a href="https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L33704&amp;bc=gAAAAAAA==">https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L33704&amp;bc=gAAAAAAA==</a>
-----------------------	---

Jurisdiction(s): M, 11	NCD/LCD Document (s): L35677  <a href="https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L35677&amp;bc=gAAAAAAA==">https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L35677&amp;bc=gAAAAAAA==</a>
------------------------	---

Jurisdiction(s): 6, K	NCD/LCD Document (s): A52423  <a href="https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A52423&amp;bc=gAAAAAAA==">https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A52423&amp;bc=gAAAAAAA==</a>
-----------------------	---

### Medicare Part B Administrative Contractor (MAC) Jurisdictions

Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC

#### REMICADE® (infliximab) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2019, Magellan Rx Management

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

#### REMICADE® (infliximab) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2019, Magellan Rx Management