



Bavencio[®] (avelumab) (Intravenous)

Document Number: IC-0295

Last Review Date: 06/01/2023 Date of Origin: 04/25/2017 Dates Reviewed: 04/2017, 05/2017, 08/2017, 11/2017, 02/2018, 05/2018, 09/2018, 11/2018, 03/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020, 08/2020, 12/2020, 03/2021, 06/2021, 09/2021, 12/2021, 03/2022, 06/2022, 09/2022, 12/2022, 03/2023, 06/2023

I. Length of Authorization

Coverage will be provided for 6 months and may be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Bavencio 200 mg/10 mL single-dose vial: 4 vials per 14 days
- B. Max Units (per dose and over time) [HCPCS Unit]:
 - 80 billable units (800 mg) every 14 days (all indications)

III. Initial Approval Criteria¹

Coverage is provided in the following conditions:

• Patient is at least 18 years of age, unless otherwise indicated; AND

Universal Criteria

• Patient has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy (e.g., nivolumab, pembrolizumab, dostarlimab, atezolizumab, durvalumab, cemiplimab, nivolumab/relatlimab-rmbw, etc.), unless otherwise specified; **AND**

Merkel Cell Carcinoma (MCC) $\dagger \ddagger \Phi$ 1,2,4,5

- Patient is at least 12 years of age; **AND**
- Used as single-agent therapy; AND
- Patient has metastatic disease

Urothelial Carcinoma (Bladder Cancer) † ‡ 1,4,6,8,16

- Used as single-agent therapy; AND
 - Patient has one of the following diagnoses:

Proprietary & Confidential © 2023 Magellan Health, Inc.



- Locally advanced or metastatic urothelial carcinoma **†**
- Muscle invasive bladder cancer with local recurrence or persistent disease in a preserved bladder
- Metastatic or local bladder cancer recurrence post cystectomy
- Metastatic upper genitourinary (GU) tract tumors
- Metastatic urothelial carcinoma of the prostate
- Recurrent or metastatic primary carcinoma of the urethra (excluding recurrence of stage T3-4 disease or palpable inguinal lymph nodes),
 AND
- Used for disease that progressed during or following platinum-containing chemotherapy*; OR
- Used as second-line treatment after chemotherapy other than a platinum; **OR**
- Used as first-line maintenance treatment; AND
 - Patient has locally advanced or metastatic urothelial carcinoma (inclusive of bladder, upper GU tract, urethra, and/or prostate cancer); AND
 - Patient has not progressed with first-line platinum-containing chemotherapy

* Note: 6,7,17,20

- If patient was progression free for > 12 months after platinum therapy, consider re-treatment with platinum-based therapy if the patient is still platinum eligible (see below for cisplatin- or platinum-ineligible comorbidities).

- Cisplatin-ineligible comorbidities may include the following: CrCl < 60 mL/min, $PS \ge 2$, hearing loss of ≥ 25 decibels (dB) at two contiguous frequencies, grade ≥ 2 peripheral neuropathy, or NYHA class ≥ 3 . Carboplatin may be substituted for cisplatin particularly in those patients with a CrCl < 60 mL/min or a PS of 2.
- Platinum-ineligible comorbidities may include the following: CrCl < 30 mL/min, $PS \ge 3$, $grade \ge 2$ peripheral neuropathy, or NYHA class > 3, etc.

Renal Cell Carcinoma (RCC) † ‡ 1,4,9,14

- Used in combination with axitinib; AND
- Used as first-line therapy; **AND**
- Used for the treatment of advanced, relapsed, or stage IV disease and clear cell histology

Gestational Trophoblastic Neoplasia ‡ 4,13,15

- Used as single-agent therapy for multiagent chemotherapy-resistant disease; AND
 - Patient has intermediate placental site trophoblastic tumor (PSTT) or epithelioid trophoblastic tumor (ETT); AND
 - Patient has recurrent or progressive disease; **OR**
 - Patient has high-risk disease (i.e., prognostic score \geq 7 or FIGO stage IV disease)

Endometrial Carcinoma (Uterine Neoplasms) ‡ 4,18

• Used as single-agent therapy; AND



- Patient has recurrent disease; AND
- Used as subsequent therapy for microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) tumors
- FDA Approved Indication(s); Compendia Recommended Indication(s); Orphan Drug

IV. Renewal Criteria¹

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe or life-threatening infusion-related reactions, hepatotoxicity, severe immunemediated adverse reactions (e.g., pneumonitis, hepatitis, colitis, endocrinopathies, nephritis with renal dysfunction, dermatitis/dermatologic adverse reactions, etc.), major adverse cardiovascular events (MACE) when used in combination with axitinib, complications of allogeneic hematopoietic stem cell transplantation (HSCT), etc.

V. Dosage/Administration ^{1,13,18}

Indication	Dose		
All Indications	Administer 800 mg intravenously every 14 days, until disease progression or unacceptable toxicity		
Dosing should be calculated using actual body weight and not flat dosing (as applicable) based on the following:			
$\underline{Weight > 60 \ kg}$			
• Standard dose 800 mg IV every 2 weeks			

Weight is ≤ 60 kg:

• Use 600 mg IV every 2 weeks

Note: This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide. Patient-specific variables should be taken into account.

VI. Billing Code/Availability Information

HCPCS Code:

• J9023 – Injection, avelumab, 10 mg; 1 billable unit = 10 mg

NDC:

• Bavencio 200 mg/10 mL single-dose vial: 44087-3535-xx



VII. References

- 1. Bavencio [package insert]. Rockland, MA; EMD Serono, Inc; July 2022. Accessed May 2023.
- Kaufman HL, Russell J, Hamid O, et al. Avelumab in patients with chemotherapyrefractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial. Lancet Oncol. 2016 Oct;17(10):1374-1385.
- 3. Novakovic AM, Wilkins JJ, Dai H, et al. Changing body weight-based dosing to a flat dose for avelumab in metastatic Merkel cell and advanced urothelial carcinoma. Clin Pharmacol Ther. 2019 Sep 25.
- 4. Referenced with permission from the NCCN Drugs and Biologics Compendium (NCCN Compendium®) avelumab. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2023.
- 5. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium[®]) Merkel Cell Carcinoma. Version 1.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium[®] is a derivative work of the NCCN Guidelines[®]. NATIONAL COMPREHENSIVE CANCER NETWORK[®], NCCN[®], and NCCN GUIDELINES[®] are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2023.
- 6. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium[®]) Bladder Cancer. Version 2.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium[®] is a derivative work of the NCCN Guidelines[®]. NATIONAL COMPREHENSIVE CANCER NETWORK[®], NCCN[®], and NCCN GUIDELINES[®] are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2023.
- 7. Gupta S, Sonpavde G, Grivas P, et al. Defining "platinum-ineligible" patients with metastatic urothelial cancer (mUC). J Clin Oncol. 2019 Mar 1;37(7_suppl):451.
- Patel MR, Ellerton J, Infante JR, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-label, phase 1 trial. Lancet Oncol. 2018 Jan;19(1):51-64. Doi: 10.1016/S1470-2045(17)30900-2. Epub 2017 Dec 5.
- Motzer RJ, Penkov K, Haanen J, et al. Avelumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. N Engl J Med. 2019 Mar 21;380(12):1103-1115. Doi: 10.1056/NEJMoa1816047. Epub 2019 Feb 16.



- 10. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. J Oncol Pract. 2018 Mar;14(3):e130-e136.
- 11. Hematology/Oncology Pharmacy Association (2019). Intravenous Cancer Drug Waste Issue Brief. Retrieved from http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug_Waste_2019.pdf
- 12. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. BMJ. 2016 Feb 29;352:i788.
- 13. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Gestational Trophoblastic Neoplasia. Version 1.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2023.
- 14. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Kidney Cancer. Version 4.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2023.
- 15. You B, Bolze PA, Lotz JP, et al. Avelumab in Patients With Gestational Trophoblastic Tumors With Resistance to Single-Agent Chemotherapy: Cohort A of the TROPHIMMUN Phase II Trial. J Clin Oncol. 2020 Sep 20;38(27):3129-3137. Doi: 10.1200/JCO.20.00803. Epub 2020 Jul 27.
- Powles T, Park SH, Voog E, et al. Avelumab Maintenance Therapy for Advanced or Metastatic Urothelial Carcinoma. N Engl J Med. 2020 Sep 24;383(13):1218-1230. Doi: 10.1056/NEJMoa2002788. Epub 2020 Sep 18. PMID: 32945632.
- 17. Bellmunt, J. (2023). Treatment of metastatic urothelial cancer of the bladder and urinary tract. In Lerner SP, Shah S (Eds.), *UptoDate*. Last updated March 15, 2023. Accessed May 4, 2023. Available from <u>https://www.uptodate.com/contents/treatment-of-metastatic-urothelial-cancer-of-the-bladder-and-urinary-tract?search=cisplatin%20ineligible&source=search_result&selectedTitle=1~150&usage_ty_pe=default&display_rank=1.
 </u>
- 18. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Uterine Neoplasms. Version 2.2023. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network,



Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2023.

- Galsky MD, Balar AV, Black PC, et al. Society for Immunotherapy of Cancer (SITC) clinical practice guideline on immunotherapy for the treatment of urothelial cancer. Journal for ImmunoTherapy of Cancer 2021;9:e002552. doi: 10.1136/jitc-2021-002552.
- 20. Gupta S, Bellmunt J, Plimack ER, et al. Defining "platinum-ineligible" patients with metastatic urothelial cancer (mUC). J Clin Oncol. 2022 June 1;40(16_suppl):4577.

ICD-10	ICD-10 Description	
C4A.0	Merkel cell carcinoma of lip	
C4A.10	Merkel cell carcinoma of eyelid, including canthus	
C4A.111	Merkel cell carcinoma of right upper eyelid, including canthus	
C4A.112	Merkel cell carcinoma of right lower eyelid, including canthus	
C4A.121	Merkel cell carcinoma of left upper eyelid, including canthus	
C4A.122	Merkel cell carcinoma of left lower eyelid, including canthus	
C4A.20	Merkel cell carcinoma of unspecified ear and external auricular canal	
C4A.21	Merkel cell carcinoma of right ear and external auricular canal	
C4A.22	Merkel cell carcinoma of left ear and external auricular canal	
C4A.30	Merkel cell carcinoma of unspecified part of face	
C4A.31	Merkel cell carcinoma of nose	
C4A.39	Merkel cell carcinoma of other parts of face	
C4A.4	Merkel cell carcinoma of scalp and neck	
C4A.51	Merkel cell carcinoma of anal skin	
C4A.52	Merkel cell carcinoma of skin of breast	
C4A.59	Merkel cell carcinoma of other part of trunk	
C4A.60	Merkel cell carcinoma of unspecified upper limb, including shoulder	
C4A.61	Merkel cell carcinoma of right upper limb, including shoulder	
C4A.62	Merkel cell carcinoma of left upper limb, including shoulder	
C4A.70	Merkel cell carcinoma of unspecified lower limb, including hip	
C4A.71	Merkel cell carcinoma of right lower limb, including hip	
C4A.72	Merkel cell carcinoma of left lower limb, including hip	
C4A.8	Merkel cell carcinoma of overlapping sites	
C4A.9	Merkel cell carcinoma, unspecified	
C54.0	Malignant neoplasm of isthmus uteri	
C54.1	Malignant neoplasm of endometrium	
C54.2	Malignant neoplasm of myometrium	
C54.3	Malignant neoplasm of fundus uteri	
C54.8	Malignant neoplasm of overlapping sites of corpus uteri	

Appendix 1 – Covered Diagnosis Codes

BAVENCIO® (avelumab) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval. ©2023, Magellan Rx Management

ICD-10	ICD-10 Description	
C54.9	Malignant neoplasm of corpus uteri, unspecified	
C55	Malignant neoplasm of uterus, part unspecified	
C58	Malignant neoplasm of placenta	
C61	Malignant neoplasm of prostate	
C64.1	Malignant neoplasm of right kidney, except renal pelvis	
C64.2	Malignant neoplasm of left kidney, except renal pelvis	
C64.9	Malignant neoplasm of unspecified kidney, except renal pelvis	
C65.1	Malignant neoplasm of right renal pelvis	
C65.2	Malignant neoplasm of left renal pelvis	
C65.9	Malignant neoplasm of unspecified renal pelvis	
C66.1	Malignant neoplasm of right ureter	
C66.2	Malignant neoplasm of left ureter	
C66.9	Malignant neoplasm of unspecified ureter	
C67.0	Malignant neoplasm of trigone of bladder	
C67.1	Malignant neoplasm of dome of bladder	
C67.2	Malignant neoplasm of lateral wall of bladder	
C67.3	Malignant neoplasm of anterior wall of bladder	
C67.4	Malignant neoplasm of posterior wall of bladder	
C67.5	Malignant neoplasm of bladder neck	
C67.6	Malignant neoplasm of ureteric orifice	
C67.7	Malignant neoplasm of urachus	
C67.8	Malignant neoplasm of overlapping sites of bladder	
C67.9	Malignant neoplasm of bladder, unspecified	
C68.0	Malignant neoplasm of urethra	
C7B.1	Secondary Merkel cell carcinoma	
D09.0	Carcinoma in situ of bladder	
D39.2	Neoplasm of uncertain behavior of placenta	
001.9	Hydatidiform mole, unspecified	
Z85.42	Personal history of malignant neoplasm of other parts of uterus	
Z85.51	Personal history of malignant neoplasm of bladder	
Z85.59	Personal history of malignant neoplasm of other urinary tract organ	
Z85.821	Personal history of Merkel cell carcinoma	

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at:



<u>https://www.cms.gov/medicare-coverage-database/search.aspx</u>. Additional indications may be covered at the discretion of the health plan.

	Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor			
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC			
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC			
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)			
6	MN, WI, IL	National Government Services, Inc. (NGS)			
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.			
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)			
N (9)	FL, PR, VI	First Coast Service Options, Inc.			
J (10)	TN, GA, AL	Palmetto GBA, LLC			
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC			
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.			
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)			
15	KY, OH	CGS Administrators, LLC			

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

