

## Blincyto® (blinatumomab) (Intravenous)

Document Number: IC-0225

Last Review Date: 06/02/2020

Date of Origin: 12/16/2014

Dates Reviewed: 12/2014, 09/2015, 02/2016, 05/2016, 08/2016, 11/2016, 02/2017, 05/2017, 08/2017, 11/2017, 02/2018, 04/2018, 09/2018, 12/2018, 03/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020

### I. Length of Authorization

- Relapsed or Refractory B-Cell Precursor Acute Lymphocytic Leukemia (ALL)
  - Initial coverage will be provided for 30 weeks for a total of five cycles (2 cycles of induction followed by 3 cycles of consolidation)
  - Continued coverage will be provided every 24 weeks for a maximum of two additional authorizations (4 cycles of continued therapy)
- MRD+ B-Cell Precursor Acute Lymphocytic Leukemia (ALL)
  - Initial coverage will be provided for 24 weeks for a total of four cycles (1 cycle of induction followed by 3 cycles of consolidation)
  - Continued coverage may not be renewed

### II. Dosing Limits

#### A. Quantity Limit (max daily dose) [NDC Unit]:

- Blincyto 35 mcg powder for injection: 28 vials per 42 day supply

#### B. Max Units (per dose and over time) [HCPCS Unit]:

- Relapsed or Refractory B-Cell Precursor Acute Lymphocytic Leukemia (ALL)
  - Cycle 1 – 5 (Induction/Consolidation)
    - 980 billable units per 42 days
  - Cycle 6 – 9 (Continued Therapy)
    - 980 billable units per 84 days
- MRD+ B-Cell Precursor Acute Lymphocytic Leukemia (ALL)
  - Cycle 1 – 4 (Induction/Consolidation)
    - 980 billable units per 42 days

### III. Initial Approval Criteria

Coverage is provided in the following conditions:

#### Universal Criteria <sup>1</sup>

- Used as single agent therapy; **AND**
- Patient has not received a live vaccine within 2 weeks before initiating therapy and will not receive a live vaccine concurrently with Blincyto; **AND**

#### B-Cell Precursor Acute Lymphocytic Leukemia (ALL) † $\Phi$ <sup>1,2,3,4,5,8</sup>

- Patient is at least 1 month old; **AND**
  - Patient has relapsed or refractory disease (Philadelphia chromosome [Ph]-positive patients must be TKI intolerant/refractory); **OR**
  - Used in patients with minimal residual disease positive (MRD+) as consolidation therapy following a complete response/remission to induction therapy; **OR**
  - Used in patients that are MRD+ after consolidation therapy ‡; **OR**
  - Used in patients that are Ph-positive with less than complete response or high-risk genetics (e.g., KMT2Ar, low hypodiploidy, TCF3-PBX1, IKZF1 deletion, etc.) ‡

† FDA Approved Indication(s); ‡ Compendium Recommended Indication(s);  $\Phi$  Orphan Drug

### IV. Renewal Criteria <sup>1,2</sup>

Coverage can be renewed based upon the following criteria:

- Patient continues to meet universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: Cytokine Release Syndrome (CRS), neurological toxicities, serious infections, pancreatitis, tumor lysis syndrome, neutropenia/febrile neutropenia, elevation of LFTs, leukoencephalopathy, etc.; **AND**
- Treatment response or stabilization of disease as indicated by CBC, bone marrow cytogenic analysis, QPCR, or FISH; **AND**
  - Patient has not exceeded a total of 4 cycles of continued therapy or 9 total cycles of Blincyto for the treatment of relapsed or refractory disease; **OR**
  - Continued therapy for use in the treatment of MRD+ ALL may not be renewed.

### V. Dosage/Administration

Indication	Dose
	<u>Relapsed/Refractory Disease*</u> <ul style="list-style-type: none"><li>➤ Weight greater than or equal to 45 kg</li></ul>

<p>Acute Lymphoblastic Leukemia</p>	<ul style="list-style-type: none"> <li>- <u>Cycle 1 (induction):</u> <ul style="list-style-type: none"> <li>• 9 mcg daily x 7 days, then 28 mcg daily x 21 days in a 42 day cycle</li> </ul> </li> <li>- <u>Cycles 2-5 (induction/consolidation):</u> <ul style="list-style-type: none"> <li>• 28 mcg daily x 28 days in a 42 day cycle.</li> </ul> </li> <li>- <u>Cycles 6-9 (continued therapy):</u> <ul style="list-style-type: none"> <li>• 28 mcg daily x 28 days in an 84 day cycle</li> </ul> </li> </ul> <p>➤ Weight less than 45 kg</p> <ul style="list-style-type: none"> <li>- <u>Cycle 1 (induction) :</u> <ul style="list-style-type: none"> <li>• 5 mcg/m<sup>2</sup>/day (not to exceed 9 mcg/day) x 7 days, then 15 mcg/m<sup>2</sup>/day (not to exceed 28 mcg/day) x 21 days in a 42 day cycle</li> </ul> </li> <li>- <u>Cycles 2-5 (induction/consolidation):</u> <ul style="list-style-type: none"> <li>• 15 mcg/m<sup>2</sup>/day (not to exceed 28 mcg/day) x 28 days in a 42 day cycle</li> </ul> </li> <li>- <u>Cycles 6-9 (continued therapy):</u> <ul style="list-style-type: none"> <li>• 15 mcg/m<sup>2</sup>/day (not to exceed 28 mcg/day) x 28 days in an 84 day cycle</li> </ul> </li> </ul> <p><i>*Up to 9 total cycles of therapy.</i></p>
	<p><b><u>MRD+ Disease*</u></b></p> <p>➤ Weight greater than or equal to 45 kg</p> <ul style="list-style-type: none"> <li>- <u>Cycle 1 (induction) :</u> <ul style="list-style-type: none"> <li>• 28 mcg daily x 28 days in a 42-day cycle</li> </ul> </li> <li>- <u>Cycles 2-4 (consolidation):</u> <ul style="list-style-type: none"> <li>• 28 mcg daily x 28 days in a 42 day cycle</li> </ul> </li> </ul> <p>➤ Weight less than 45 kg</p> <ul style="list-style-type: none"> <li>- <u>Cycle 1 (induction) :</u> <ul style="list-style-type: none"> <li>• 15 mcg/m<sup>2</sup>/day (not to exceed 28 mcg/day) x 28 days in a 42 day cycle</li> </ul> </li> <li>- <u>Cycles 2-4 (consolidation):</u> <ul style="list-style-type: none"> <li>• 15 mcg/m<sup>2</sup>/day (not to exceed 28 mcg/day) x 28 days in a 42 day cycle</li> </ul> </li> </ul> <p><i>*Up to 4 total cycles of therapy.</i></p>

## VI. Billing Code/Availability Information

### HCPCS Code:

- J9039 - Injection, blinatumomab, 1 microgram; 1 billable unit = 1 microgram

### NDC:

- Blincyto 35 mcg single-use powder for injection: 55513-0160-xx

## VII. References

1. Blincyto [package insert]. Thousand Oaks, CA; Amgen, March 2020. Accessed April 2020.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) blinatumomab. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2020.

3. Jen EY, Xu Q, Schetter A, Przepiorka D, et al. FDA Approval: Blinatumomab for Patients with B-cell Precursor Acute Lymphoblastic Leukemia in Morphologic Remission with Minimal Residual Disease. *Clin Cancer Res.* 2019 Jan 15;25(2):473-477. doi: 10.1158/1078-0432.CCR-18-2337. Epub 2018 Sep 25.
4. Kantarjian H, Stein A, Gökbuget N, et al. Blinatumomab versus Chemotherapy for Advanced Acute Lymphoblastic Leukemia. *N Engl J Med.* 2017 Mar 2;376(9):836-847. doi: 10.1056/NEJMoa1609783.
5. Martinelli G, Boissel N, Chevallier P, et al. Complete Hematologic and Molecular Response in Adult Patients With Relapsed/Refractory Philadelphia Chromosome-Positive B-Precursor Acute Lymphoblastic Leukemia Following Treatment With Blinatumomab: Results From a Phase II, Single-Arm, Multicenter Study. *J Clin Oncol.* 2017 Jun 1;35(16):1795-1802. doi: 10.1200/JCO.2016.69.3531. Epub 2017 Mar 29. Erratum in: *J Clin Oncol.* 2017 Aug 10;35(23):2722. *J Clin Oncol.* 2017 Aug 20;35(24):2856.
6. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Pediatric Acute Lymphoblastic Leukemia 2.2020. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2020.
7. Palmetto GBA. Local Coverage Article (LCA): Billing and Coding for Chemotherapy (A56141). Centers for Medicare & Medicaid Services, Inc. Updated on 03/24/2020 with effective date 04/30/2020. Accessed April 2020.
8. CGS Administrators, LLC. Local Coverage Article (LCA): Billing and Coding: Blinatumomab (Blinicyto) – J9039 (A56886). Centers for Medicare & Medicaid Services, Inc. Updated on 09/20/2019 with effective date 09/26/2019. Accessed April 2020.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C83.50	Lymphoblastic (diffuse) lymphoma unspecified site
C83.51	Lymphoblastic (diffuse) lymphoma lymph nodes of head, face, and neck
C83.52	Lymphoblastic (diffuse) lymphoma intrathoracic lymph nodes
C83.53	Lymphoblastic (diffuse) lymphoma intra-abdominal lymph nodes
C83.54	Lymphoblastic (diffuse) lymphoma lymph nodes of axilla and upper limb
C83.55	Lymphoblastic (diffuse) lymphoma lymph nodes of inguinal region and lower limb
C83.56	Lymphoblastic (diffuse) lymphoma intrapelvic lymph nodes
C83.57	Lymphoblastic (diffuse) lymphoma spleen
C83.58	Lymphoblastic (diffuse) lymphoma lymph nodes of multiple sites
C83.59	Lymphoblastic (diffuse) lymphoma extranodal and solid organ sites

C91.00	Acute lymphoblastic leukemia not having achieved remission
C91.01	Acute lymphoblastic leukemia, in remission
C91.02	Acute lymphoblastic leukemia, in relapse

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

<b>Jurisdiction(s):</b> J&M	<b>NCD/LCD/LCA Document (s):</b> A56141
<a href="https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A56141&amp;bc=gAAAAAAAAAAAA">https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A56141&amp;bc=gAAAAAAAAAAAA</a>	

<b>Jurisdiction(s):</b> 15	<b>NCD/LCD/LCA Document (s):</b> A56886
<a href="https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A56886&amp;bc=gAAAAAAAAAAAA">https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A56886&amp;bc=gAAAAAAAAAAAA</a>	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC