



# Imlygic<sup>®</sup> (talimogene laherparepvec) Intralesional

Document Number: IC-0274

Last Review Date: 05/04/2023 Date of Origin: 04/26/2016 Dates Reviewed: 04/25/2017, 04/2018, 05/2019, 05/2020, 05/2021, 05/2022, 05/2023

#### I. Length of Authorization

Coverage will be provided for 6 months and may be renewed

#### II. Dosing Limits

#### A. Quantity Limit (max daily dose) [NDC unit]:

- Imlygic 10<sup>6</sup> (1 million) PFU per mL single-use vial: 4 mL one time only
- Imlygic 10<sup>8</sup> (100 million) PFU per mL single-use vial: 4 mL three weeks after initial treatment followed by 4 mL every two weeks thereafter

#### B. Max Units (per dose and over time) [HCPCS Unit]:

Initial treatment:4 billable unitsSecond treatment:400 billable units occurring 3 weeks after initial treatmentAll subsequent treatments:400 billable units occurring 2 weeks after previous treatment

#### III. Initial Approval Criteria<sup>1,2</sup>

Coverage is provided in the following conditions:

• Patient is at least 18 years of age; AND

#### Universal Criteria

- Patient is not pregnant (*Note: Women of childbearing potential should be advised to use an effective method of contraception to prevent pregnancy during treatment*); **AND**
- Patient is not immunocompromised (i.e., patients with a history of primary or acquired immunodeficient states, leukemia, lymphoma, AIDS or other clinical manifestations of infection with human immunodeficiency viruses, and those on immunosuppressive therapy); **AND**
- Treatment will be administered via intralesional injection; AND



#### Cutaneous Melanoma † ‡ $\Phi$

- Used for unresectable recurrent disease **†**; **OR**
- Used as primary treatment for unresectable or borderline resectable stage III disease with clinically positive node(s); **OR**
- Used for oligometastatic disease with accessible lesions; OR
- Used for widely disseminated distant metastatic disease with limited extracranial lesions; **OR**
- Patient has limited resectable or unresectable disease; AND
  - $\circ$  Used for stage III disease with clinical satellite/in-transit metastases; OR
  - Used for local satellite/in-transit recurrence

FDA Approved Indication(s); Compendia Recommended Indication(s); Orphan Drug

#### IV. Renewal Criteria<sup>1,2</sup>

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Patient continues to have injectable lesions; AND
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: herpetic infection, injection site complications (e.g., necrosis, ulceration, cellulitis, systemic bacterial infection, etc.), immune-mediated events, plasmacytoma at injection site, obstructive airway disorder, etc.

#### V. Dosage/Administration<sup>1</sup>

Indication	Dose	
Melanoma	<ul> <li>Initial Treatment</li> <li>Imlygic 10<sup>6</sup> (1 million) PFU per mL</li> <li>Inject largest lesion(s) first</li> <li>Prioritize injection of remaining lesion(s) based on lesion size until maximum injection volume is reached or until all injectable lesion(s) have been treated</li> <li>Second Treatment</li> <li>Imlygic 10<sup>8</sup> (100 million) PFU per mL</li> <li>Administer 3 weeks after initial treatment</li> <li>Inject any new lesion(s) (lesions that have developed since initial treatment)</li> </ul>	
	first.	



- Prioritize injection of remaining lesion(s) based on lesion size until maximum injection volume is reached or until all injectable lesion(s) have been treated.
   All subsequent treatments (including reinitiation)
   Imlygic 10<sup>8</sup> (100 million) PFU per mL
   Administer 2 weeks after previous treatment
  - Inject any new lesion(s) (lesions that have developed since previous treatment) first.
  - Prioritize injection of remaining lesion(s) based on lesion size until maximum injection volume is reached or until all injectable lesion(s) have been treated.

The total injection volume for each treatment visit should not exceed 4 mL for all injected lesions combined. It may not be possible to inject all lesions at each treatment visit or over the full course of treatment. Previously injected and/or uninjected lesion(s) may be injected at subsequent treatment visits.

Lesion size (longest dimension)	Intralesional Injection Volume
> 5 cm	up to 4 mL
> 2.5 cm to 5 cm	up to 2 mL
> 1.5 cm to 2.5 cm	up to 1 mL
> 0.5 cm to 1.5 cm	up to 0.5 mL
$\leq 0.5~{ m cm}$	up to 0.1 mL

# VI. Billing Code/Availability Information

## HCPCS Code:

• J9325 – Injection, talimogene laherparepvec, per 1 million plaque forming units; 1 billable unit = 10<sup>6</sup> (1 million) PFU

## NDC(s):

- Imlygic 10<sup>6</sup> (1 million) PFU per mL single-use vial: 55513-0078-xx
- Imlygic 10<sup>8</sup> (100 million) PFU per mL single-use vial: 55513-0079-xx

# VII. References

- 1. Imlygic [package insert]. Thousand Oaks, CA; Amgen Inc.; February 2023. Accessed March 2023.
- 2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for talimogene laherparepvec. National Comprehensive Cancer Network, 2023. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2023.
- Andtbacka RHI, Kaufman HL, Collichio F, et al. Talimogene laherparepvec improves durable response rate in patients with advanced melanoma. J Clin Oncol. 2015;33:2780-2788.



4. Andtbacka RHI, Kaufman HL, Collichio F, et al. Talimogene laherparepvec improves durable response rate in patients with advanced melanoma. J Clin Oncol. 2015;33 (suppl Clinical Study Protocol):doi:10.1200/JCO.2014.58.3377.

ICD-10	ICD-10 Description	
C43.0	Malignant melanoma of lip	
C43.111	Malignant melanoma of right upper eyelid, including canthus	
C43.112	Malignant melanoma of right lower eyelid, including canthus	
C43.121	Malignant melanoma of left upper eyelid, including canthus	
C43.122	Malignant melanoma of left lower eyelid, including canthus	
C43.20	Malignant melanoma of unspecified ear and external auricular canal	
C43.21	Malignant melanoma of right ear and external auricular canal	
C43.22	Malignant melanoma of left ear and external auricular canal	
C43.30	Malignant melanoma of unspecified part of face	
C43.31	Malignant melanoma of nose	
C43.39	Malignant melanoma of other parts of face	
C43.4	Malignant melanoma of scalp and neck	
C43.51	Malignant melanoma of anal skin	
C43.52	Malignant melanoma of skin of breast	
C43.59	Malignant melanoma of other part of trunk	
C43.60	Malignant melanoma of unspecified upper limb, including shoulder	
C43.61	Malignant melanoma of right upper limb, including shoulder	
C43.62	Malignant melanoma of left upper limb, including shoulder	
C43.70	Malignant melanoma of unspecified lower limb, including hip	
C43.71	Malignant melanoma of right lower limb, including hip	
C43.72	Malignant melanoma of left lower limb, including hip	
C43.8	Malignant melanoma of overlapping sites of skin	
C43.9	Malignant melanoma of skin, unspecified	
Z85.820	Personal history of malignant melanoma of skin	

#### Appendix 1 – Covered Diagnosis Codes

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <u>https://www.cms.gov/medicare-coverage-database/search.aspx</u>. Additional indications may be covered at the discretion of the health plan.



Medicare Part B Administrative Contractor (MAC) Jurisdictions				
Jurisdiction	Applicable State/US Territory	Contractor		
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC		
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC		
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)		
6	MN, WI, IL	National Government Services, Inc. (NGS)		
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.		
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)		
N (9)	FL, PR, VI	First Coast Service Options, Inc.		
J (10)	TN, GA, AL	Palmetto GBA, LLC		
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC		
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.		
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)		
15	КҮ, ОН	CGS Administrators, LLC		

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD): N/A

