

Hemophilia Products – Factor IX: Alphanine SD, Alprolix, Bebulin, BeneFIX, Idelvion, Ixinity, Mononine, Profilnine, Rebinyn, and Rixubis

Document Number: IC-0339

Last Review Date: 11/26/2018

Date of Origin: 12/16/2014

Dates Reviewed: 12/2014, 4/2015, 5/2015, 09/2015, 12/2015, 03/2016, 06/2016, 12/2016, 06/2017, 09/2017, 11/2017, 11/2018

I. Length of Authorization

- Initial and renewal authorization periods vary by specific covered indication
- Unless otherwise specified, the initial authorization will be provided for 3 months and may be renewed.

The cumulative amount of medication(s) the patient has on-hand will be taken into account when authorizing. The authorization will allow up to 5 doses on-hand for the treatment of acute bleeding episodes as needed for the duration of the authorization.

II. Dosing Limits

A. Quantity Limit (max daily dose) [Pharmacy Benefit]:

N/A

B. Max Units (per dose and over time) [Medical Benefit]:

Alprolix: 23,000 billable units per 28 day supply
Alphanine: 36,800 billable units per 28 day supply
BeneFIX: 36,800 billable units per 28 day supply
Bebulin: 36,800 billable units per 28 day supply
Idelvion: 25,300 billable units per 28 day supply
Rixubis: 73,600 billable units per 28 day supply
Ixinity: 36,800 billable units per 28 day supply
Profilnine: 36,800 billable units per 28 day supply
Mononine: 36,800 billable units per 28 day supply

III. Initial Approval Criteria

Hemophilia Management Program

Requirements for half-life study and inhibitor tests are a part of the hemophilia management program. This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide.

A. AlphaNine SD, Alprolix, Bebulin, BeneFIX, Profilnine SD, Mononine, Rixubis, IXINITY, Idelvion and Rebinyn

Coverage is provided in the following conditions:

Hemophilia B (congenital factor IX deficiency aka Christmas disease) †

- Diagnosis of congenital factor IX deficiency has been confirmed by blood coagulation testing; **AND**
- Used as treatment in at least one of the following:
 - Control and prevention of acute bleeding episodes (episodic treatment of acute hemorrhage); **OR**
 - Perioperative management; **OR**
 - Authorization is valid for 1 month
 - Routine prophylaxis to prevent or reduce the frequency of bleeding episodes (excluding Rebinyn); **AND**
 - Patient must have severe hemophilia B (factor IX level of <1%); **OR**
 - Patient has at least two documented episodes of spontaneous bleeding into joints; **AND**
- Therapy **NOT** used for induction of immune tolerance in patients with Hemophilia B for **ONLY** the following products:
 - Alprolix
 - Rixubis
 - Ixinity
 - Idelvion
 - Rebinyn

Hemophilia Management Program

- If the request is for prophylaxis and the requested dose exceeds dosing limits under part II, a half-life study should be performed to determine the appropriate dose and dosing interval.
- If the request is for Alprolix, Idelvion, or Rebinyn, a half-life study should be performed to determine the appropriate dose and dosing interval.

- For Alprolix, 50 IU/kg every 7 days is the preferred dosing regimen. To obtain 100 IU every 10 days, a half-life study must be submitted showing a significant clinical benefit over 50 IU/kg every 7 days.
- Prior to switching to Alprolix, Idelvion, or Rebinyn, a half-life study should also be performed on current non- EHL factor IX product to ensure that a clinical benefit will be achieved.
- For members with a BMI \geq 30, a half-life study should be performed to determine the appropriate dose and dosing interval.
- For minimally treated patients (< 50 exposure days to factor products) previously receiving a different factor product, inhibitor testing is required at baseline, then at every comprehensive care visit (yearly for the mild and moderate patients, semi-annually for the severe patients)

† FDA Approved Indication(s)

IV. Dispensing Requirements for Rendering Providers (Hemophilia Management Program)

- Prescriptions cannot be filled without an expressed need from the patient, caregiver or prescribing practitioner. Auto-filling is not allowed.
- Monthly, rendering provider must submit for authorization of dispensing quantity before delivering factor product. Information submitted must include:
 - Original prescription information, requested amount to be dispensed, vial sizes available to be ordered from the manufacturer, and patient clinical history (including patient product inventory and bleed history)
 - Factor dose should not exceed +1% of the prescribed dose and a maximum of three vials may be dispensed per dose. If unable to provide factor dosing within the required threshold, below the required threshold, the lowest possible dose able to be achieved above +1% should be dispensed. Prescribed dose should not be increased to meet assay management requirements.
- The cumulative amount of medication(s) the patient has on-hand should be taken into account when dispensing factor product. Patients should not have more than 5 extra doses on-hand for the treatment of acute bleeding episodes.
- Dispensing requirements for renderings providers are a part of the hemophilia management program. This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide.

V. Renewal Criteria

Coverage can be renewed based upon the following criteria:

- Patient continues to meet criteria identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: symptoms of allergic-anaphylactic reactions (anaphylaxis, dyspnea, rash);

thromboembolic events (thromboembolism, pulmonary embolism); and development of neutralizing antibodies (inhibitors); **AND**

- Any increases in dose must be supported by an acceptable clinical rationale (i.e. weight gain, half-life study results, increase in breakthrough bleeding when patient is fully adherent to therapy, etc.).
- The cumulative amount of medication(s) the patient has on-hand will be taken into account when authorizing. The authorization will allow up to 5 doses on-hand for the treatment of acute bleeding episodes as needed for the duration of the authorization.
- Patient meets the disease specific criteria below:

Treatment of acute bleeding episodes/Treatment of Spontaneous and trauma-induced bleeding episodes/On-demand treatment of bleeding episodes

- Renewals will be approved for a 3 month authorization period

Prevention of acute bleeding episodes/Routine prophylaxis to prevent or reduce the frequency of bleeding episode

- Renewals will be approved for a 6 month authorization period

VI. Dosage/Administration

Alprolix

Indication	Dose
Control and prevention of bleeding episodes Hemophilia B	<p>One unit per kilogram body weight increases the circulating Factor IX level by 1% (IU/dL). Estimate the required dose or the expected in vivo peak increase in Factor IX level expressed as IU/dL (or % of normal) using the following: IU/dL (or % of normal) = [Total Dose (IU)/Body Weight (kg)] x Recovery (IU/dL per IU/kg)</p> <p><u>Minor and Moderate</u></p> <p>Circulating Factor IX required (% of normal) = 30-60 IU/dL -Repeat every 48 hours as needed</p> <p><u>Major</u></p> <p>Circulating Factor IX required (% of normal) = 80-100 IU/dL - Consider repeat dose after 6-10 hours, then every 24 hours for 3 days, then every 48 hours until healing achieved.</p>
Perioperative management Hemophilia B	<p><u>Minor</u></p> <p>Circulating Factor IX required (% of normal) = 50-80 IU/dL -Repeat every 24-48 hours as needed, until bleeding stops and healing is achieved.</p> <p><u>Major</u></p> <p>Circulating Factor IX required (% of normal) = 60-100 IU/dL (initial level) - Consider repeat dose after 6-10 hours, then every 24 hours for 3 days, then every 48 hours until bleeding stops and healing achieved.</p>
Routine prophylaxis Hemophilia B	50 IU/kg once weekly or 100 IU/kg once every 10 days. Adjust dosing regimen based on individual response.

AlphaNine SD

Indication	Dose
Control and prevention of bleeding episodes Hemophilia B	<p>One unit per kilogram body weight increases the circulating Factor IX level by 1% (IU/dL). Number of Factor IX IU required = body wt (kg) x Desired increase in Plasma Factor IX(percent) x 1.0 IU/kg</p> <p><u>Minor</u></p> <p>Circulating Factor IX required (20 – 30 % of normal) = 20-30 IU/kg -Repeat every 12 hours as needed for 1-2 days</p> <p><u>Moderate</u></p> <p>Circulating Factor IX required (25 - 50% of normal) = 25-50 IU/kg -Repeat every 12 hours as needed for 2-7 days</p> <p><u>Major</u></p> <p>Circulating Factor IX required (50% of normal) = 50-100 IU/kg - Consider repeat dose after 12 hours as needed for 3-5 days. Following this treatment period, FIX levels should be maintained at 20% (20 IU FIX/kg/twice daily) until healing has been achieved. Major hemorrhages may require treatment for up to 10 days</p>
Routine prophylaxis Hemophilia B §	25-40 IU/kg two times weekly or 15-30 IU/kg two times weekly. Adjust dosing regimen based on individual response.
Perioperative management Hemophilia B	Prior to surgery, FIX should be brought to 50-100% of normal (50-100 IU/kg repeat every 12 hours). For the next 7 to 10 days, or until healing has been achieved, the patient should be maintained at 50-100%FIX levels (50-100 IU/kg every 12 hours).

BeneFIX

Indication	Dose
Control and prevention of bleeding episodes Hemophilia B and Perioperative management of Hemophilia B	<p>One unit per kilogram body weight increases the circulating Factor IX level by 1% (IU/dL). ADULT: Number of Factor IX IU required = body wt (kg) x Desired increase in Plasma Factor IX(percent) x 1.3 IU/kg; CHILD (<15 years) Number of Factor IX IU required = body wt (kg) x Desired increase in Plasma Factor IX(percent) x 1.4 IU/kg</p> <p><u>Minor</u></p> <p>Circulating Factor IX required (% of normal) = 20-30 IU/dL -Repeat every 12-24 hours as needed for 1-2 days</p> <p><u>Moderate</u></p> <p>Circulating Factor IX required (% of normal) = 25-50 IU/dL -Repeat every 12-24 hours as needed for 2-7 days</p> <p><u>Major</u></p> <p>Circulating Factor IX required (% of normal) = 50-100 IU/dL - Consider repeat dose after 12-24 hours as needed for 7- 10 days.</p>
Routine prophylaxis Hemophilia B §	25-40 IU/kg two times weekly or 15-30 IU/kg two times weekly. Adjust dosing regimen based on individual response.

Bebulin

Indication	Dose
Control and prevention of bleeding episodes Hemophilia B	<p>One unit per kilogram body weight increases the circulating Factor IX level by 1% (IU/dL). Number of Factor IX IU required = body wt (kg) x Desired increase in Plasma Factor IX(percent) x 1.2 IU/kg</p> <p><u>Minor</u></p> <p>Circulating Factor IX required (% of normal) (20%)= 25-35 IU/dL -Repeat every 24 hours as needed until adequate wound healing</p> <p><u>Moderate</u></p> <p>Circulating Factor IX required (% of normal) (40%)= 50-65 IU/dL -Repeat every 24 hours as needed for 2 days or until adequate wound healing</p> <p><u>Major</u></p> <p>Circulating Factor IX required (% of normal)(>60%) = 75-90 IU/dL - Consider repeat dose after 24 hours as needed for 2-3 days or until adequate wound healing.</p>
Routine prophylaxis Hemophilia B §	25-40 IU/kg two times weekly or 15-30 IU/kg two times weekly. Adjust dosing regimen based on individual response.
Perioperative management Hemophilia B	<p><u>Minor</u></p> <p>Circulating Factor IX required (% of normal) (40-60%)= 50-75 IU/dL given 1 hour prior to surgery, repeat every 12 hours, and continue replacement therapy over 1 to 2 weeks postop until adequate wound healing is achieved.</p> <p><u>Major</u></p> <p>Circulating Factor IX required (% of normal) (>60%)= 75-90 IU/dL given 1 hour prior to surgery, repeat every 12 hours, and continue replacement therapy over for up to 2 weeks postop. If treatment is required beyond 2 weeks post-up, then dosing interval can be adjusted to every 24 hours and continued until adequate wound healing is achieved.</p>

Idelvion

Indication	Dose
Control and prevention of bleeding episodes	<ul style="list-style-type: none"> One IU of IDELVION per kg body weight is expected to increase the circulating activity of Factor IX as follows: <ul style="list-style-type: none"> Adolescents and adults: 1.3 IU/dL per IU/kg Pediatrics (<12 years): 1 IU/dL per IU/kg Administer intravenously. Do not exceed infusion rate of 10 mL per minute. Dosage and duration of treatment with IDELVION depends on the severity of the Factor IX deficiency, the location and extent of bleeding, and the patient's clinical condition, age and recovery of Factor IX. Determine the initial dose using the following formula: <ul style="list-style-type: none"> Required Dose (IU) = Body Weight (kg) x Desired Factor IX rise (% of normal or IU/dL) x (reciprocal of recovery (IU/kg per IU/dL)) Adjust dose based on the patient's clinical condition and response. <p><u>Minor/Moderate</u></p> <p>Desired peak Factor IX Level (% of normal or IU/dL): 30-60, dosed every 48-72 hours for at least 1 day until healing is achieved</p> <p><u>Major</u></p>

	Desired peak Factor IX Level (% of normal or IU/dL): 60-100, dosed every 48-72 hours for 7-14 days until bleeding stops. Maintenance dose is weekly.
Perioperative management Hemophilia B	<p><u>Minor</u></p> <p>Desired peak Factor IX Level (% of normal or IU/dL): 50-80, dosed every 48-72 hours for at least 1 day until healing is achieved</p> <p><u>Major</u></p> <p>Desired peak Factor IX Level (% of normal or IU/dL): 60-100, dosed every 48-72 hours for 7-14 days until bleeding stops. Repeat dose every 48-72 hours for the first week or until healing is achieved. Maintenance dose is once or twice weekly.</p>
Routine prophylaxis Hemophilia B	<p><u>Patients ≥12 years of age:</u> 25-40 IU/kg body weight every 7 days. Patients who are well-controlled on this regimen may be switched to a 14-day interval at 50-75 IU/kg body weight.</p> <p><u>Patients <12 years of age:</u> 40-55 IU/kg body weight every 7 days.</p>

Ixinity

Indication	Dose
Control and prevention of bleeding episodes Congenital Hemophilia B	<p>One IU per kg body weight increases the circulating activity of factor IX by 0.698 IU/dL</p> <p><u>Initial dose:</u> Required factor IX units (IU) = body weight (kg) x desired factor IX increase (% of normal of IU/dL) x reciprocal of observed recovery (IU/kg per IU/dL)</p> <p><u>Maintenance dose:</u> Depends upon the type of bleed or surgery, clinical response, and the severity of the underlying factor IX deficiency</p> <p><u>Minor</u> Desired peak Factor IX Level (% of normal or IU/dL): 30-60, dosed every 24 hours on days 1-3 until healing is achieved</p> <p><u>Moderate</u> Desired peak Factor IX Level (% of normal or IU/dL): 40-60, dosed every 24 hours on days 2-7 until healing is achieved</p> <p><u>Major or Life threatening</u> Desired peak Factor IX Level (% of normal or IU/dL): 60-100, dosed every 12 – 24 hours on days 2-14 until healing is achieved</p>
Perioperative management Congenital Hemophilia B	<p><u>Minor</u></p> <p>Pre-op: Desired peak Factor IX Level (% of normal or IU/dL) 50-80 Post-op: Desired peak Factor IX Level (% of normal or IU/dL) 30-80, dosed every 24 hours on days 1-5, depending on type of procedure</p> <p><u>Major</u></p> <p>Pre-op: Desired peak Factor IX Level (% of normal or IU/dL) 60-80 Post-op: Desired peak Factor IX Level (% of normal or IU/dL) 40-60, dosed every 8 – 24 hours on days 1-3, or 30-50 dosed every 8 – 24 hours on days 4-6, or 20-40 dosed every 8 -24 hours on days 7-14</p>

Mononine

Indication	Dose
Control and prevention of bleeding episodes and perioperative management Hemophilia B	<p>One unit per kilogram body weight increases the circulating Factor IX level by 1% (IU/dL). Estimate the required dose with the following formula: Number of Factor IX IU required (IU) = Body Weight (in kg) x desired Factor IX increase (% or IU/dL normal) x 1.0 IU/kg [per IU/dL]</p> <p><u>Minor Spontaneous Hemorrhage Prophylaxis</u></p> <p>Circulating Factor IX required (% of normal)(15-25%) = up to 20-30IU/kg for one dose. Repeat in 24 hours if necessary.</p> <p><u>Major Trauma or Surgery</u></p> <p>Circulating Factor IX required (% of normal)(25-50%) = up to 75 IU/kg Dosed every 18-30 hours depending on T_{1/2} and measured Factor IX levels. Continue for up to 10 days depending upon nature of insult.</p>

Profilnine SD

Indication	Dose
Control and prevention of bleeding episodes Hemophilia B	<p>One unit per kilogram body weight increases the circulating Factor IX level by 1% (IU/dL). Number of Factor IX IU required = body wt (kg) x Desired increase in Plasma Factor IX(percent) x 1.0 IU/kg</p> <p><u>Mild to Moderate</u></p> <p>Single dose of product sufficient to raise plasma factor IX levels to 20 to 30 percent of normal. 20-30 IU/kg every 16-24 hours until hemorrhage stops and healing is achieved. For minor, may repeat for 1-2 days, for moderate, may repeat for 2-7 days.</p> <p><u>Major</u></p> <p>Single dose of product sufficient to raise plasma factor IX levels to 30 to 50 percent of normal. Daily infusions are generally required.</p>
Routine prophylaxis Hemophilia B §	25-40 IU/kg two times weekly or 15-30 IU/kg two times weekly. Adjust dosing regimen based on individual response.
Perioperative management Hemophilia B	Surgery associated with bleeding in factor IX deficient patients requires factor IX levels of 30 to 50 percent. 30-50 IU/kg every 16-24 hours for 7-10 days until healing is achieved. For dental extractions, the factor IX level should be raised to 50 percent immediately prior to procedure; additional factor IX complex may be given if bleeding recurs.

Rebinyn

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia B	<p><u>Minor and Moderate</u></p> <p>40 IU/kg of actual body weight. A single dose should be sufficient for minor and moderate bleeds. Additional doses of 40 IU/kg can be given.</p> <p><u>Major</u></p> <p>80 IU/kg of actual body weight. Additional doses of 40 IU/kg can be given.</p>

Perioperative management of bleeding	<u>Minor</u> Pre-op: 40 IU/kg of actual body weight (single pre-op dose should be sufficient) Post-op: Additional doses can be given if required
Congenital Hemophilia B	<u>Major</u> Pre-op: 80 IU/kg of actual body weight Peri/Post-op: 40 IU/kg of actual body weight. As clinically needed for the perioperative management of bleeding, repeated doses of 40 IU/kg (in 1-3 day intervals) within the first week after major surgery may be administered. Due to the long half-life the frequency of dosing in the post-surgical setting may be extended to once weekly after the first week until bleeding stops and healing is achieved.

Rixubis

Indication	Dose
Control and prevention of bleeding episodes Hemophilia B	One IU per kilogram body weight increases the circulating activity of Factor IX by 0.7 IU/dL for patients <12 years of age and 0.9 IU/dL for patients ≥ 12 years of age. Initial dose = body wt (kg) x desired factor IX increase (percent of normal or IU/dL) x reciprocal of observed recovery (IU/kg per IU/dL) <u>Minor</u> Circulating Factor IX level required (% or IU/dL) = 20-30 every 12 - 24 hours for at least 1 day, until healing is achieved <u>Moderate</u> Circulating Factor IX level required (% or IU/dL) = 25-50 every 12 - 24 hours for 2 - 7 days, until bleeding stops and healing is achieved <u>Major</u> Circulating Factor IX level required (% or IU/dL) = 50-100 every 12 - 24 hours for 7 - 10 days, until bleeding stops and healing is achieved
Routine prophylaxis Hemophilia B	Dosing for previously treated patients (PTPs): <u>Patients <12 years of age</u> 60 – 80 IU/kg twice weekly <u>Patients > 12 years of age</u> 40 – 60 IU/kg twice weekly Adjust the dose based on the individual patient’s age, bleeding pattern, and physical activity.
Perioperative management Hemophilia B	<u>Minor</u> Circulating Factor IX level required (% or IU/dL) = 30-60 every 24 hours for at least 1 day, until healing is achieved <u>Major</u> Circulating Factor IX level required (% or IU/dL) = 80-100 every 8 - 24 hours for 7 - 10 days, until bleeding stops and healing is achieved

§ Utrecht and/or Malmö protocols used as basis for dosing³⁶

VII. Billing Code/Availability Information

Jcode & NDC:

Drug	Manufacturer	J-Code	1 Billable Unit Equiv.	Vial Size	NDC
AlphaNine SD	Grifols Biologicals Inc	J7193	1 IU	500 units	68516-3601-02 68516-3607-02
				1000 units	68516-3602-02 68516-3608-02
				1500 units	68516-3603-02 68516-3609-02
Mononine	CSL Behring LLC	J7193	1 IU	1000 units	00053-6233-02
Alprolix	Biogen Idec, Inc	J7201	1 IU	250 units	64406-0966-01
				500 units	64406-0911-01
				1000 units	64406-0922-01
				2000 units	64406-0933-01
				3000 units	64406-0944-01
				4000 units	64406-0977-01
Bebulin	Baxalta US Inc	J7194	1 IU	Unassigned size	64193-0445-02
Profilnine SD	Grifols Biologicals Inc	J7194	1 IU	500 units	68516-3201-01 68516-3207-01
				1000 units	68516-3202-02 68516-3208-02
				1500 units	68516-3203-02 68516-3209-02
BeneFIX	Wyeth Biopharma	J7195	1 IU	250 units	58394-0633-03
				500 units	58394-0634-03
				1000 units	58394-0635-03
				2000 units	58394-0636-03

Hemophilia Products – Factor IX Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2018, Magellan Rx Management

				3000 units	58394-0637-03
Ixinity	Cangene Corp	J7195	1 IU	250 units	70504-0287-05
				500 units	70504-0282-05
				1000 units	70504-0283-05
				1500 units	70504-0284-05
				2000 units	70504-0288-05
				3000 units	70504-0289-05
				Rixubis	Baxalta US Inc
500 units	00944-3028-02				
1000 units	00944-3030-02				
2000 units	00944-3032-02				
3000 units	00944-3034-02				
Idelvion	Novozymes Biopharma A/S	J7202	1 IU	250 units	69911-0864-02
				500 units	69911-0865-02
				1000 units	69911-0866-02
				2000 units	69911-0867-02
Rebinyn	Novo Nordisk Inc	J7203	1 IU N/A	500 units	00169-7905-01
				1000 units	00169-7901-01
				2000 units	00169-7902-01

VIII. References

1. AlphaNine SD [package insert]. Los Angeles, CA; Grifols Biologicals Inc.; January 2013. Accessed June 2017.
2. Alprolix [package insert]. Cambridge, MA; Biogen Idec, Inc.; April 2017. Accessed June 2017.
3. Bebulin [package insert]. Westlake Village, CA; Baxalta US Inc. September 2015. Accessed June 2016.
4. BeneFIX [package insert]. Philadelphia, PA; Wyeth Biopharma; August 2015. Accessed June 2017.
5. Ixinity [package insert]. Winnipeg, Manitoba, Canada. Cangene Corporation; August 2016. Accessed June 2017.
6. Mononine [package insert]. Kankakee, IL; CSL Behring LLC; April 2016. Accessed June 2017.

7. Profilnine [package insert]. Los Angeles, CA; Grifols Biologicals Inc.; May 2014. Accessed June 2017.
8. Rebinyn [package insert]. Plainsboro, NJ; Novo Nordisk Inc.; June 2017. Accessed June 2017
9. Rixubis [package insert]. Westlake Village, CA; Baxalta US Inc.; March 2016; Accessed June 2017.
10. Idelvion [package insert]. Bagsvaerd, Denmark; Novozymes Biopharma A/S; February 2017. Accessed June 2017.
11. MASAC RECOMMENDATIONS CONCERNING PRODUCTS LICENSED FOR THE TREATMENT OF HEMOPHILIA AND OTHER BLEEDING DISORDERS. 2016 National Hemophilia Foundation. MASAC Document #249; October 2016. Available at: <http://www.hemophilia.org>. Accessed June 2017.
12. Guidelines for the Management of Hemophilia. 2nd Edition. World Federation of Hemophilia. 2013. Available at: <https://www1.wfh.org/publication/files/pdf-1472.pdf>. Accessed June 2017.
13. First Coast Service Options, Inc. Local Coverage Determination (LCD): Hemophilia Clotting Factors (L33684). Centers for Medicare & Medicaid Services, Inc. Updated on 01/03/2017 with effective date 01/01/2017. Accessed June 2017.
14. Novitas Solutions, Inc. Local Coverage Determination (LCD): Hemophilia Clotting Factors (L35111). Centers for Medicare & Medicaid Services, Inc. Updated on 01/06/2017 with effective date 01/01/2017. Accessed June 2017.
15. Annual Review of Factor Replacement Products. Oklahoma Health Care Authority Review Board. Updated April 2016. Access June 2016.
16. Graham A1, Jaworski K. Pharmacokinetic analysis of anti-hemophilic factor in the obese patient. Haemophilia. 2014 Mar;20(2):226-9.
17. Croteau SE1, Neufeld EJ. Transition considerations for extended half-life factor products. Haemophilia. 2015 May;21(3):285-8.
18. Mingot-Castellano, et al. Application of Pharmacokinetics Programs in Optimization of Haemostatic Treatment in Severe Hemophilia a Patients: Changes in Consumption, Clinical Outcomes and Quality of Life. Blood. 2014 December; 124 (21).
19. MASAC RECOMMENDATION CONCERNING PROPHYLAXIS. 2016 National Hemophilia Foundation. MASAC Document #241; February 2016. Available at: <http://www.hemophilia.org>. Accessed August 2017.

Appendix 1 – Covered Diagnosis Codes

AlphaNine SD, Alprolix, Bebulin, BeneFIX, Profilnine SD, Mononine, Rixubis, Ixinity, Idelvion, and Rebinyn

ICD-10	ICD-10 Description
D67	Hereditary factor IX deficiency

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD):

Jurisdiction(s): H,L	NCD/LCD Document (s): L35111
https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L35111&bc=gAAAAAAAAAAAAA==	

Jurisdiction(s): N	NCD/LCD Document (s): L33684
https://www.cms.gov/medicare-coverage-database/search/lcd-date-search.aspx?DocID=L33684&bc=gAAAAAAAAAAAAA==	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Cahaba Government Benefit Administrators, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC