

Vyxeos® (daunorubicin and cytarabine – liposome) (Intravenous)

Document Number: IC-0314

Last Review Date: 09/01/2020

Date of Origin: 08/29/2017

Dates Reviewed: 08/2017, 07/2018, 07/2019, 07/2020, 09/2020

I. Length of Authorization

Coverage will be provided for a maximum of 2 cycles of induction (5 doses total) and 2 cycles of consolidation (4 doses total) within 6 months. Coverage may not be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Vyxeos single-dose vial: 23 vials total

B. Max Units (per dose and over time) [HCPCS Unit]:

- Induction: 132 billable units per dose (3 vials per dose; 5 doses total)
- Consolidation: 88 billable units per dose (2 vials per dose; 4 doses total)

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

Universal Criteria ¹

- Baseline left ventricular ejection fraction (LVEF) is within normal limits and will be reassessed prior to consolidation and as clinically required; **AND**
- Cumulative lifetime anthracycline (e.g., daunorubicin, etc.) dose does not exceed 550 mg/m² (or 400 mg/m² in patients who received radiation to the mediastinum); **AND**
- Will not be used in combination with other chemotherapy; **AND**

Acute Myeloid Leukemia (AML) † Φ ¹⁻³

- Patient has one of the following sub-types of disease:
 - Therapy-related acute myeloid leukemia (t-AML)
 - AML with myelodysplasia-related changes (AML-MRC)

- Antecedent myelodysplastic syndrome/chronic myelomonocytic leukemia (antecedent MDS/CMML)
- Used for one of the following:
 - Patient has newly diagnosed disease (*Note: For antecedent MDS/CMML, use only allowed in patients age ≥ 60 years in candidates for intensive remission induction therapy*); **OR**
 - Used for re-induction after standard-dose cytarabine induction ‡; **AND**
 - Patients ≥ 60 years with residual disease; **OR**
 - Patients < 60 years with significant residual disease in the absence of a hypocellular marrow and core binding factor (CBF) abnormalities; **OR**
 - Used post-remission ‡; **AND**
 - Patients ≥ 60 years with complete response to previous intensive therapy; **OR**
 - Patients < 60 years with treatment-related disease other than core binding factor (CBF) and/or unfavorable cytogenetics and/or molecular abnormalities

† FDA Approved Indication(s); ‡ Compendia recommended indication(s); ◊ Orphan Drug

IV. Renewal Criteria

Authorizations may not be renewed.

V. Dosage/Administration

Indication	Dose
t-AML, antecedent MDS/CMML & AML-MRC	<p><u>First Induction</u></p> <ul style="list-style-type: none"> • daunorubicin 44 mg/m² and cytarabine 100 mg/m² liposome intravenously days 1, 3 and 5 <p><u>Second induction</u></p> <ul style="list-style-type: none"> • daunorubicin 44 mg/m² and cytarabine 100 mg/m² liposome intravenously days 1 and 3 <ul style="list-style-type: none"> ○ Only for patients who fail to respond to the first induction cycle ○ May be administered 2 to 5 weeks after the first induction cycle if there was no unacceptable toxicity <p><u>Consolidation</u></p> <ul style="list-style-type: none"> • daunorubicin 29 mg/m² and cytarabine 65 mg/m² liposome intravenously days 1 and 3 <ul style="list-style-type: none"> ○ Administer the first consolidation cycle 5 to 8 weeks after the start of the last induction cycle ○ Administer the second consolidation cycle 5 to 8 weeks after the start of the first consolidation cycle if there was not unacceptable toxicity of disease progression

VI. Billing Code/Availability Information

HCPCS code:

- J9153 – Injection, liposomal, 1 mg daunorubicin and 2.27 mg cytarabine: 1 billable unit = 1 mg daunorubicin and 2.27 mg cytarabine

NDC:

- Vyxeos (44 mg daunorubicin and 100 mg cytarabine) liposome, single-dose vial: 68727-0745-xx

VII. References

1. Vyxeos [package insert]. Palo Alto, CA; Jazz Pharmaceuticals, Inc., July 2019. Accessed June 2020.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for cytarabine/daunorubicin liposome. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2020.
3. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Acute Myeloid Leukemia. Version 3.2020. National Comprehensive Cancer Network, 2020. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed August 2020.
4. Lin TL, Ryan RJ, Fadert S, et al. Outcomes in older patients with high-risk/secondary AML who achieved remission with CPX-351 versus 7+3 but did not undergo transplant: Phase 3 exploratory analysis. J Clin Onco; DOI: 10.1200/JCO.2020.38.15_suppl.7537 Journal of Clinical Oncology 38, no. 15_suppl(May 20, 2020)7537-7537.
5. Palmetto GBA. Local Coverage Article (LCA): Billing and Coding: Chemotherapy (A56141). Centers for Medicare & Medicaid Services, Inc. Updated on 05/26/2020 with effective date 04/30/2020. Accessed August 2020.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C92.00	Acute myeloblastic leukemia not having achieved remission
C92.01	Acute myeloblastic leukemia in remission
C92.50	Acute myelomonocytic leukemia not having achieved remission
C92.51	Acute myelomonocytic leukemia in remission

VYXEOS® (daunorubicin and cytarabine - liposome) **Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2020, Magellan Rx Management

ICD-10	ICD-10 Description
C92.60	Acute myeloid leukemia with 11q23-abnormality not having achieved remission
C92.61	Acute myeloid leukemia with 11q23-abnormality in remission
C92.A0	Acute myeloid leukemia with multilineage dysplasia not having achieved remission
C92.A1	Acute myeloid leukemia with multilineage dysplasia in remission
C93.00	Acute monoblastic/monocytic leukemia not having achieved remission
C93.01	Acute monoblastic/monocytic leukemia in remission
C94.00	Acute erythroid leukemia not having achieved remission
C94.01	Acute erythroid leukemia in remission
C94.20	Acute megakaryoblastic leukemia not having achieved remission
C94.21	Acute megakaryoblastic leukemia in remission

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

Jurisdiction(s): J&M	NCD/LCD/Article Document (s): A56141
https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A56141&bc=gAAAAAAAAAAAA	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)

VYXEOS® (daunorubicin and cytarabine - liposome)
Prior Auth Criteria

Medicare Part B Administrative Contractor (MAC) Jurisdictions

Jurisdiction	Applicable State/US Territory	Contractor
15	KY, OH	CGS Administrators, LLC