



## Onivyde® (irinotecan liposome injection) (Intravenous)

-E-

Document Number: IC-0428

Last Review Date: 05/04/2023

Date of Origin: 02/2019

Dates Reviewed: 02/2019, 05/2019, 05/2020, 05/2021, 05/2022, 05/2023

### I. Length of Authorization

Coverage will be provided for 6 months and may be renewed.

### II. Dosing Limits

#### A. Quantity Limit (max daily dose) [NDC Unit]:

- Onivyde 43 mg/10 mL single-dose vial: 4 vials per 14 days

#### B. Max Units (per dose and over time) [HCPCS Unit]:

- All indications: 172 billable units per 14 days

### III. Initial Approval Criteria <sup>1</sup>

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

#### Universal Criteria <sup>1</sup>

- Patient does not have bowel obstruction; **AND**
- Therapy will not be substituted for other drugs containing irinotecan HCl; **AND**

#### Pancreatic Adenocarcinoma † ‡ Φ <sup>1,2</sup>

- Used in combination with fluorouracil and leucovorin; **AND**
- Patient has locally advanced or metastatic disease; **AND**
- Used after disease progression with gemcitabine-based therapy

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

† FDA Approved Indication(s); ‡ Compendia recommended indication; Ⓞ Orphan Drug

#### IV. Renewal Criteria <sup>1</sup>

Coverage can be renewed based upon the following criteria:

- Patient continues to meet universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe diarrhea, severe neutropenia, interstitial lung disease, severe hypersensitivity reactions (including anaphylactic reactions), etc.

#### V. Dosage/Administration <sup>1,3</sup>

Indication	Dose
All Indications	Administer 70 mg/m <sup>2</sup> intravenously every 14 days <u>Note:</u> Patients homozygous for the UGT1A1*28 allele: Administer 50 mg/m <sup>2</sup> every 14 days and may titrate up to 70 mg/m <sup>2</sup> as tolerated in subsequent cycles.

#### VI. Billing Code/Availability Information

HCPCS Code:

- J9205 – Injection, irinotecan liposome, 1 mg: 1 billable unit = 1 mg

NDC:

- Onivyde 43 mg/10 mL single dose vial: 15054-0043-xx

#### VII. References (STANDARD)

1. Onivyde [package insert]. Cambridge, MA: Ipsen Biopharmaceuticals, Inc.; February 2023. Accessed April 2023.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium<sup>®</sup>) irinotecan liposomal. National Comprehensive Cancer Network, 2023. The NCCN Compendium<sup>®</sup> is a derivative work of the NCCN Guidelines<sup>®</sup>. NATIONAL COMPREHENSIVE CANCER NETWORK<sup>®</sup>, NCCN<sup>®</sup>, and NCCN GUIDELINES<sup>®</sup> are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2023.
3. Wang-Gillam A, Li CP, Bodky G, NAPOLI-1 study group. Nanoliposomal irinotecan with fluorouracil and folinic acid in metastatic pancreatic cancer after previous gemcitabine-

based therapy (NAPOLI-1): a global, randomised, open-label, phase 3 trial. *Lancet*. 2016 Feb 6;387(10018):545-557. doi: 10.1016/S0140-6736(15)00986-1. Epub 2015 Nov 29.

## VIII. References (ENHANCED)

- 1e. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Pancreatic Adenocarcinoma, Version 2.2022. National Comprehensive Cancer Network, 2023. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed April 2023.
- 2e. Pelzer U, Schwaner I, Stieler J, et al. Best supportive care (BSC) versus oxaliplatin, folinic acid and 5-fluorouracil (OFF) plus BSC in patients for second-line advanced pancreatic cancer: a phase III-study from the German CONKO-study group. *Eur J Cancer*. 2011 Jul;47(11):1676-81.
- 3e. Oettle H, Riess H, Stieler JM, et al. Second-Line Oxaliplatin, Folinic Acid, and Fluorouracil Versus Folinic Acid and Fluorouracil Alone for Gemcitabine-Refractory Pancreatic Cancer: Outcomes From the CONKO-003 Trial. *Journal of Clinical Oncology* 2014 32:23, 2423-2429.
- 4e. Gill S, Ko YJ, Cripps C, et al. PANCREOX: A Randomized Phase III Study of Fluorouracil/Leucovorin With or Without Oxaliplatin for Second-Line Advanced Pancreatic Cancer in Patients Who Have Received Gemcitabine-Based Chemotherapy. *Journal of Clinical Oncology* 2016 34:32, 3914-3920.
- 5e. Heinemann V, Vehling-Kaiser U, Waldschmidt D, et al. Gemcitabine plus erlotinib followed by capecitabine versus capecitabine plus erlotinib followed by gemcitabine in advanced pancreatic cancer: final results of a randomised phase 3 trial of the 'Arbeitsgemeinschaft Internistische Onkologie' (AIO-PK0104). *Gut*. 2012;62(5):751-9.
- 6e. Yoo C, Hwang JY, Kim JE, et al. A randomised phase II study of modified FOLFIRI.3 vs modified FOLFOX as second-line therapy in patients with gemcitabine-refractory advanced pancreatic cancer. *Br J Cancer*. 2009;101(10):1658-63.
- 7e. Neuzillet C, Hentic O, Rousseau B, et al. FOLFIRI regimen in metastatic pancreatic adenocarcinoma resistant to gemcitabine and platinum-salts. *World J Gastroenterol*. 2012;18(33):4533-41.
- 8e. Assaf E, Verlinde-Carvalho M, Delbaldo C, et al. 5-fluorouracil/leucovorin combined with irinotecan and oxaliplatin (FOLFIRINOX) as second-line chemotherapy in patients with metastatic pancreatic adenocarcinoma. *Oncology*. 2011;80(5-6):301-6.
- 9e. Xiong HQ, Varadhachary GR, Blais JC, et al. Phase 2 trial of oxaliplatin plus capecitabine (XELOX) as second-line therapy for patients with advanced pancreatic cancer. *Cancer*. 2008 Oct 15;113(8):2046-52.
- 10e. Le DT, Durham JN, Smith KN, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science*. 2017;357(6349):409-413.

11e. Portal A, Pernot S, Tougeron D, et al. Nab-paclitaxel plus gemcitabine for metastatic pancreatic adenocarcinoma after Folfirinox failure: an AGEO prospective multicentre cohort. *Br J Cancer*. 2015;113(7):989-95.

12e. Magellan Health, Magellan Rx Management. Onivyde Clinical Literature Review Analysis. Last updated April 2023. Accessed April 2023.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C25.0	Malignant neoplasm of head of pancreas
C25.1	Malignant neoplasm of body of the pancreas
C25.2	Malignant neoplasm of tail of pancreas
C25.3	Malignant neoplasm of pancreatic duct
C25.7	Malignant neoplasm of other parts of pancreas
C25.8	Malignant neoplasm of overlapping sites of pancreas
C25.9	Malignant neoplasm of pancreas, unspecified
Z85.07	Personal history of malignant neoplasm of pancreas

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC

M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC