

Revlimid[®] (lenalidomide) (Oral)

Document Number: IC-0106

Last Review Date: 01/04/2022

Date of Origin: 01/01/2012

Dates Reviewed: 12/2012, 06/2013, 05/2014, 03/2015, 01/2016, 01/2017, 01/2018, 01/2019, 06/2019, 01/2020, 01/2021, 01/2022

I. Length of Authorization

Coverage will be provided for six months and may be renewed.

- Previously untreated Follicular Lymphoma may be renewed for up to 18 cycles.
- Previously treated Follicular Lymphoma and Marginal Zone Lymphoma may be renewed for up to 12 cycles.
- Systemic Light Chain Amyloidosis: Primary treatment or treatment of relapsed/refractory disease as repeat of initial therapy may be renewed for up to 8 cycles.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Units]:

- Revlimid 2.5mg capsules: 1 capsule per day
- Revlimid 5mg capsules: 1 capsule per day
- Revlimid 10mg capsules: 1 capsule per day
- Revlimid 15mg capsules: 1 capsule per day
- Revlimid 20mg capsules: 1 capsule per day
- Revlimid 25mg capsules: 1 capsule per day

B. Max Units (per dose and over time) [HCPCS Units]:

- MDS, MF, CLL/SLL (in combination with rituximab): 10mg daily
- Multiple Myeloma (maintenance therapy), Primary CNS Lymphoma, Systemic Light Chain Amyloidosis (in combination with bortezomib, cyclophosphamide, or ixazomib): 15mg daily
- Follicular Lymphoma/Marginal Zone Lymphoma: 20mg daily
- All other indications: 25mg daily

III. Initial Approval Criteria ^{1,2}

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

Universal Criteria ¹

- Both patient AND prescriber are enrolled in and meet the conditions of the Revlimid REMS[®] program; **AND**

Myelodysplastic Syndrome (MDS) † Φ ^{1,2}

- Patient has myelodysplastic/myeloproliferative neoplasms with ring sideroblasts and thrombocytosis (MDS/MPN-RS-T) overlap disease; **AND**
 - Used as a single agent or in combination with a hypomethylating agent (e.g., azacitidine, decitabine, etc.); **OR**
- Patient has symptomatic anemia; **AND**
 - Patient has lower risk disease (defined as IPSS [Low/Intermediate-1]); **AND**
 - Used as a single agent for del(5q) mutation †; **OR**
 - Patient has lower risk disease (defined as IPSS-R [Very Low, Low, Intermediate]); **AND**
 - Patient does not have del(5q) mutation; **AND**
 - Patient has ring sideroblasts < 15% (or <5% with an SF3B1 mutation); **AND**
 - Patient has a serum erythropoietin (EPO) \leq 500 mU/mL; **AND**
 - Used as a single agent if no response to an erythropoiesis-stimulating agent (ESA) AND no response to an ESA with granulocyte colony-stimulation factor (G-CSF); **OR**
 - Used with ESA following failure of ESA alone; **OR**
 - Patient has a serum EPO > 500 mU/mL; **AND**
 - Patient had no response, intolerance, or a poor probability of response to immunosuppressive therapy (i.e., antithymocyte globulin [ATG] \pm cyclosporine); **OR**
 - Patient has ring sideroblasts \geq 15% (or ring sideroblasts \geq 5% with an SF3B1 mutation); **AND**
 - Patient has a serum EPO \leq 500 mU/mL with no response to ESA with G-CSF AND no response to luspatercept-aamt; **OR**
 - Patient has a serum EPO > 500 mU/mL and no response to luspatercept-aamt

Multiple Myeloma † Φ ^{1,2}

- Used in combination with dexamethasone; **OR**
- Used as primary therapy for active (symptomatic) disease; **AND**
 - Used in combination with dexamethasone AND bortezomib or carfilzomib or daratumumab or cyclophosphamide; **OR**

- Used in combination with dexamethasone AND daratumumab and bortezomib or daratumumab and carfilzomib in patients who are transplant candidates; **OR**
- Used as part of the VRd-lite regimen (bortezomib, lenalidomide, and dexamethasone) for frail patients who are non-transplant candidates; **OR**
- Used as maintenance therapy after response to primary myeloma therapy or following autologous hematopoietic stem cell transplant (auto-HSCT); **AND**
 - Used as a single agent **OR**;
 - Used in combination with bortezomib; **OR**
 - Used in combination with bortezomib and dexamethasone; **OR**
- Used for previously treated multiple myeloma for relapsed or progressive disease; **AND**
 - Used in combination with dexamethasone OR a dexamethasone-containing regimen; **OR**
 - Used as a single agent if patient is steroid-intolerant; **OR**
- Used in combination with dexamethasone for the management of POEMS (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, skin changes) syndrome

Systemic Light Chain Amyloidosis †²

- Used for newly diagnosed disease in combination with dexamethasone and bortezomib; **OR**
- Used for relapsed or refractory disease; **AND**
 - Used in combination with dexamethasone with or without cyclophosphamide or ixazomib; **OR**
 - Used in combination with dexamethasone and bortezomib as repeat of initial therapy if the patient has been relapse-free for several years

Classic Hodgkin's Lymphoma (cHL) †^{2,14}

- Used as a single agent; **AND**
- Used as third-line or greater therapy for relapsed or refractory disease

B-Cell Lymphomas^{1,2}

- Mantle Cell Lymphoma † Φ
 - Used in combination with rituximab for initial/induction therapy or in non-transplant candidates; **OR**
 - Used as a single agent after two prior therapies, one of which included bortezomib †; **OR**
 - Used as subsequent therapy in combination with rituximab; **AND**
 - Used for stable disease; **OR**
 - Used for partial response with substantial disease after induction therapy; **OR**
 - Used for relapsed or progressive disease following prior treatment with chemoimmunotherapy

- Follicular Lymphoma † ⊕
 - Used as first-line therapy in combination with rituximab; **OR**
 - Used as subsequent therapy in patients with refractory or progressive disease in combination with rituximab or obinutuzumab or as a single agent if not a candidate for anti-CD20 monoclonal antibody therapy
- AIDS-Related B-Cell Lymphoma ‡
 - Used as subsequent therapy, with or without rituximab, for relapsed non-germinal center diffuse large B-cell lymphoma in patients who are not transplant candidates
- Diffuse Large B-Cell Lymphoma (DLBCL) ‡ (including Histologic Transformation)
 - Used in combination with tafasitamab-cxix for non-transplant candidates; **AND**
 - Patient had histologic transformation from Nodal Marginal Zone Lymphoma OR from Follicular Lymphoma (without translocations of MYC and BCL2 and/or BCL6); **AND**
 - Patient received at least two prior lines of chemoimmunotherapy for indolent or transformed disease; **OR**
 - Patient received minimal or no chemoimmunotherapy prior to histologic transformation and had no response or progressive disease after chemoimmunotherapy (e.g., anthracycline- or anthracenedione-based regimen unless contraindicated); **OR**
 - Patient had partial response, no response, relapsed, progressive, or refractory disease; **OR**
 - Used with or without rituximab; **AND**
 - Patient had histologic transformation from Nodal Marginal Zone Lymphoma OR from Follicular Lymphoma; **AND**
 - Patient had transformation to non-germinal center disease; **AND**
 - Patient received multiple lines of chemoimmunotherapy for indolent or transformed disease; **OR**
 - Patient is not a transplant candidate and had partial response, no response, relapsed, progressive, or refractory non-germinal center disease
- High-Grade B-Cell Lymphoma ‡
 - Used as subsequent therapy, with or without rituximab, in non-transplant candidate patients with a partial response, no response, relapsed, progressive, or refractory disease
- Marginal Zone Lymphoma † ⊕ (includes Nodal or Splenic MZL & Gastric or Non-Gastric MALT)
 - Used as subsequent therapy for recurrent, relapsed, refractory, or progressive disease in combination with rituximab
- Multicentric Castleman's Disease ‡

- Used as subsequent therapy, with or without rituximab, for disease that has progressed following treatment of relapsed/refractory or progressive disease
- Post-Transplant Lymphoproliferative Disorders (PTLD) ‡
 - Used as subsequent therapy, following first-line chemoimmunotherapy, in patients with partial response, persistent or progressive disease to first-line therapy; **AND**
 - Patient has monomorphic PTLD (non-germinal center B-cell type)

Primary Cutaneous Lymphomas ‡²

- Mycosis Fungoides (MF)/Sezary Syndrome (SS) ‡
 - Used as a single agent; **AND**
 - Used as systemic therapy for primary treatment; **AND**
 - Patient has stage IVA2 non-Sezary or stage IVB visceral disease; **OR**
 - Patient has large cell transformation (LCT) with generalized cutaneous or extracutaneous lesions; **OR**
 - Used as systemic therapy for subsequent treatment
- Primary Cutaneous CD30+ T-Cell Lymphoproliferative Disorders ‡
 - Used as a single agent for relapsed or refractory disease; **AND**
 - Patient has primary cutaneous anaplastic large cell lymphoma (ALCL) with multifocal lesions; **OR**
 - Patient has cutaneous ALCL with regional nodes (excludes systemic ALCL)

T-Cell Lymphomas ‡²

- Peripheral T-Cell Lymphoma (*includes all of the following: peripheral T-cell lymphoma not otherwise specified, angioimmunoblastic T-cell lymphoma, enteropathy-associated T-cell lymphoma, monomorphic epitheliotropic intestinal T-cell lymphoma, nodal peripheral T-cell lymphoma with TFH phenotype, or follicular T-cell lymphoma*) ‡
 - Used as a single agent as second-line or initial palliative therapy
- Adult T-cell Leukemia/Lymphoma ‡
 - Used as a single agent as subsequent therapy for non-responders to first-line therapy for acute or lymphoma subtypes
- Hepatosplenic T-Cell Lymphoma ‡
 - Used as a single agent therapy for refractory disease after two first-line therapy regimens

Primary CNS Lymphoma ‡^{2,13}

- Used as a single-agent or in combination with rituximab; **AND**
 - Used as induction therapy for intolerance to, or patient is not a candidate for, high-dose methotrexate; **OR**
 - Used for relapsed or refractory disease; **AND**

- Patient previously received whole brain radiation therapy; **OR**
- Patient previously received high-dose methotrexate based regimen; **OR**
- Patient received prior high-dose chemotherapy with stem cell rescue (HDT/ASCR)

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma (CLL/SLL) ‡²

- Used as subsequent therapy for patients with or without del(17p)/TP53 mutation; **AND**
- Used as single agent or in combination with rituximab

Myelofibrosis (MF) ^{2,9,10}

- Used in combination with prednisone taper for management of myelofibrosis-associated anemia; **AND**
 - Patient has a serum EPO ≥ 500 mU/mL; **OR**
 - Patient has a serum EPO < 500 mU/mL and no response or loss of response to erythropoietic stimulating agents

Histiocytic Neoplasms – Langerhans Cell Histiocytosis ‡^{2,15}

- Used as a single agent; **AND**
 - Patient has single system multifocal skin disease (including mucosa); **OR**
 - Patient has relapsed/refractory disease

Kaposi Sarcoma ‡^{2,8,19}

- Used as subsequent therapy; **AND**
 - Used as a single agent for patients that do not have HIV; **OR**
 - Used in combination with antiretroviral therapy (ART) for patients with HIV; **AND**
- Patient has relapsed/refractory advanced cutaneous, oral, visceral, or nodal disease; **AND**
- Disease has progressed on or not responded to first-line therapy; **AND**
- Disease has progressed on alternate first-line therapy

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); **Ⓞ** Orphan Drug

IV. Renewal Criteria

Coverage can be renewed based upon the following criteria:

- Patient continues to meet universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: secondary primary malignancies, venous and arterial thromboembolism, hematologic toxicity

(neutropenia, thrombocytopenia), tumor lysis syndrome, hepatotoxicity, severe cutaneous reactions, tumor flare reaction, thyroid disorders (hypothyroidism, hyperthyroidism), hypersensitivity (including angioedema, anaphylaxis, and anaphylactic reactions), etc.

V. Dosage/Administration ^{1,5,9,10,13-31}

Indication	Dose
MDS	10mg daily until disease progression or unacceptable toxicity
Multiple Myeloma	<p><u>Combination Therapy:</u> 25mg daily on Days 1-21 of repeated 28-day cycles in combination with dexamethasone</p> <ul style="list-style-type: none"> For patients who are not eligible for auto-HSCT, treatment should continue until disease progression or unacceptable toxicity. For patients who are auto-HSCT-eligible, hematopoietic stem cell mobilization should occur within 4 cycles of a REVLIMID-containing therapy. <p><u>Maintenance Therapy Following Auto-HSCT:</u> 10mg daily continuously until disease progression or unacceptable toxicity</p> <ul style="list-style-type: none"> After 3 cycles of maintenance therapy, the dose can be increased to 15mg once daily if tolerated.
Previously untreated Follicular Lymphoma	20mg daily for 21 days of a 28-day cycle for up to 18 cycles.
Previously treated Follicular Lymphoma and Marginal Zone Lymphoma (Gastric MALT, Nongastric MALT, Nodal, Splenic)	20mg daily for 21 days of a 28-day cycle for up to 12 cycles in combination with a rituximab-product.
CLL/SLL	<p><u>In Combination with Rituximab:</u> 10 mg daily until disease progression or unacceptable toxicity</p> <p><u>Single Agent:</u> Up to 25 mg daily until disease progression or unacceptable toxicity</p> <p>- OR -</p> <p>25mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity</p>
Primary CNS Lymphoma	15mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity
Myelofibrosis	10mg daily until disease progression or unacceptable toxicity
	- OR -

	10 mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity
Systemic Light Chain Amyloidosis	<p><u>In Combination with Dexamethasone and Bortezomib:</u> 5-15mg daily for 21 days of a 28-day cycle for up to 8 cycles</p> <p><u>In Combination with Dexamethasone and Ixazomib:</u> 15mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity</p> <p><u>In Combination with Dexamethasone and Cyclophosphamide:</u> 15mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity</p> <p><u>In Combination with Dexamethasone:</u> 15mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity</p> <p>- OR -</p> <p>25mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity</p>
T-Cell Lymphomas	<p>25mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity</p> <p>- OR -</p> <p>25mg daily until disease progression or unacceptable toxicity</p>
All other indications	25mg daily for 21 days of a 28-day cycle until disease progression or unacceptable toxicity
<p><i>*Please refer to the prescribing information for dose adjustment for hematologic toxicities (e.g., thrombocytopenia, neutropenia) or renal impairment.</i></p>	

VI. Billing Code/Availability Information

HCPCS Code:

- J8999 – Prescription drug, oral, chemotherapeutic, Not Otherwise Specified

NDC:

- Revlimid 2.5mg capsules: 59572-0402-xx
- Revlimid 5mg capsules: 59572-0405-xx
- Revlimid 10mg capsules: 59572-0410-xx
- Revlimid 15mg capsules: 59572-0415-xx
- Revlimid 20mg capsules: 59572-0420-xx
- Revlimid 25mg capsules: 59572-0425-xx

VII. References

1. Revlimid [package insert]. Summit, NJ; Celgene Corporation; August 2021. Accessed November 2021.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) lenalidomide. National Comprehensive Cancer Network, 2021. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2021.
3. Dimopoulos M, Spencer A, Attal M, et al. Lenalidomide plus dexamethasone for relapsed or refractory multiple myeloma. *N Engl J Med* 2007; 357:2123-32.
4. Weber D, Chen C, Niesvizky R, et al. Lenalidomide plus dexamethasone for relapsed multiple myeloma in North America. *N Engl J Med* 2007; 357:2133-42.
5. Morschhauser F, Fowler NH, Feugier P, et al. Rituximab plus Lenalidomide in Advanced Untreated Follicular Lymphoma. *N Engl J Med*. 2018 Sep 6;379(10):934-947.
6. Facon T, Kumar S, Plesner T, et al; MAIA Trial Investigators. Daratumumab plus lenalidomide and dexamethasone for untreated myeloma. *N Engl J Med*. 2019;380(22):2104-2115. Doi: 10.1056/NEJMoa1817249.
7. Fenaux P, Giagounidis A, Selleslag D, et al. A Randomized phase 3 study of lenalidomide versus placebo in RBC transfusion-dependent patients with low-/intermediate-1-risk myelodysplastic syndromes with del5q. *Blood*. 2011;118(14):3765-3776.
8. Pourcher V, Desnoyer A, Assoumou L, et al. Phase II Trial of Lenalidomide in HIV-Infected Patients with Previously Treated Kaposi's Sarcoma: Results of the ANRS 154 Lenakap Trial. *AIDS Res Hum Retroviruses*. 2017 Jan;33(1):1-10. Doi: 10.1089/AID.2016.0069.
9. Mesa RA, Yao X, Cripe LD, et al. Lenalidomide and prednisone for myelofibrosis: Eastern Cooperative Oncology Group (ECOG) phase 2 trial E4903. *Blood*. 2010 Nov 25;116(22):4436-8. Doi: 10.1182/blood-2010-05-287417.
10. Chihara D, Masarova L, Newberry KJ, et al. Long-term results of a phase II trial of lenalidomide plus prednisone therapy for patients with myelofibrosis. *Leuk Res*. 2016 Sep;48:1-5. Doi: 10.1016/j.leukres.2016.06.007. Epub 2016 Jun 23.
11. Fink AM, Bahlo J, Robrecht S, et al. Lenalidomide maintenance after first-line therapy for high-risk chronic lymphocytic leukaemia (CLLM1): final results from a randomised, double-blind, phase 3 study. *Lancet Haematol*. 2017 Oct;4(10):e475-e486. Doi: 10.1016/S2352-3026(17)30171-0.
12. Chanan-Khan AA, Zaritskey A, Egyed M, et al. Lenalidomide maintenance therapy in previously treated chronic lymphocytic leukaemia (CONTINUUM): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Haematol*. 2017 Nov;4(11):e534-e543. Doi: 10.1016/S2352-3026(17)30168-0.

13. Rubenstein JL, Geng H, Fraser E, et al. Phase 1 investigation of lenalidomide/rituximab plus outcomes of lenalidomide maintenance in relapsed CNS lymphoma. *Blood Adv.* 2018 Jul 10;2(13):1595-1607. Doi: 10.1182/bloodadvances.2017014845.
14. Fehniger TA, Larson S, Trinkaus K, et al. A phase 2 multicenter study of lenalidomide in relapsed or refractory classical Hodgkin lymphoma. *Blood.* 2011 Nov 10;118(19):5119-25. Doi: 10.1182/blood-2011-07-362475.
15. Szturz P, Adam Z, Reháč Z, et al. Lenalidomide proved effective in multisystem Langerhans cell histiocytosis. *Acta Oncol.* 2012 Mar;51(3):412-5. doi: 10.3109/0284186X.2011.631581.
16. Badoux XC, Keating MJ, Wen S, et al. Phase II study of lenalidomide and rituximab as salvage therapy for patients with relapsed or refractory chronic lymphocytic leukemia. *J Clin Oncol.* 2013 Feb 10;31(5):584-91. doi: 10.1200/JCO.2012.42.8623.
17. Chanan-Khan A, Miller KC, Musial L, et al. Clinical efficacy of lenalidomide in patients with relapsed or refractory chronic lymphocytic leukemia: results of a phase II study. *J Clin Oncol.* 2006 Dec 1;24(34):5343-9. doi: 10.1200/JCO.2005.05.0401.
18. Ferrajoli A, Lee BN, Schlette EJ, et al. Lenalidomide induces complete and partial remissions in patients with relapsed and refractory chronic lymphocytic leukemia. *Blood.* 2008 Jun 1;111(11):5291-7. doi: 10.1182/blood-2007-12-130120.
19. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Kaposi Sarcoma Version 2.2021. National Comprehensive Cancer Network, 2021. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc.” To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed November 2021.
20. Kastiris E, Dialoupi I, Gavriatopoulou M, et al. Primary treatment of light-chain amyloidosis with bortezomib, lenalidomide, and dexamethasone. *Blood Adv.* 2019 Oct 22;3(20):3002-3009. doi: 10.1182/bloodadvances.2019000147.
21. Cohen OC, Sharpley F, Gillmore JD, et al. Use of ixazomib, lenalidomide and dexamethasone in patients with relapsed amyloid light-chain amyloidosis. *Br J Haematol.* 2020 May;189(4):643-649. doi: 10.1111/bjh.16401.
22. Kumar SK, Hayman SR, Buadi FK, et al. Lenalidomide, cyclophosphamide, and dexamethasone (CRd) for light-chain amyloidosis: long-term results from a phase 2 trial. *Blood.* 2012 May 24;119(21):4860-7. doi: 10.1182/blood-2012-01-407791.
23. Palladini G, Russo P, Milani P, et al. A phase II trial of cyclophosphamide, lenalidomide and dexamethasone in previously treated patients with AL amyloidosis. *Haematologica.* 2013 Mar;98(3):433-6. doi: 10.3324/haematol.2012.073593.
24. Santhorawala V, Wright DG, Rosenzweig M, et al. Lenalidomide and dexamethasone in the treatment of AL amyloidosis: results of a phase 2 trial. *Blood.* 2007 Jan 15;109(2):492-6. doi: 10.1182/blood-2006-07-030544.

25. Dispenzieri A, Lacy MQ, Zeldenrust SR, et al. The activity of lenalidomide with or without dexamethasone in patients with primary systemic amyloidosis. *Blood*. 2007 Jan 15;109(2):465-70. doi: 10.1182/blood-2006-07-032987.
26. Dispenzieri A, Lacy M, Zeldenrust SR, et al. Long Term Follow-up of Patients with Immunoglobulin Light Chain Amyloidosis Treated with Lenalidomide and Dexamethasone. *Blood*. 2008 Nov 16; 112(11): 1737. <https://doi.org/10.1182/blood.V112.11.1737.1737>.
27. Toumishey E, Prasad A, Dueck G, et al. Final report of a phase 2 clinical trial of lenalidomide monotherapy for patients with T-cell lymphoma. *Cancer*. 2015 Mar 1;121(5):716-23. doi: 10.1002/cncr.29103.
28. Morschhauser F, Fitoussi O, Haioun C, et al. A phase 2, multicentre, single-arm, open-label study to evaluate the safety and efficacy of single-agent lenalidomide (Revlimid) in subjects with relapsed or refractory peripheral T-cell non-Hodgkin lymphoma: the EXPECT trial. *Eur J Cancer*. 2013 Sep;49(13):2869-76. doi: 10.1016/j.ejca.2013.04.029.
29. Ishida T, Fujiwara H, Nosaka K, et al. Multicenter Phase II Study of Lenalidomide in Relapsed or Recurrent Adult T-Cell Leukemia/Lymphoma: ATLL-002. *J Clin Oncol*. 2016 Dec;34(34):4086-4093. doi: 10.1200/JCO.2016.67.7732.
30. Martin P, Jung SH, Pitcher B, et al. A phase II trial of lenalidomide plus rituximab in previously untreated follicular non-Hodgkin's lymphoma (NHL): CALGB 50803 (Alliance). *Ann Oncol*. 2017 Nov 1;28(11):2806-2812. doi: 10.1093/annonc/mdx496.
31. Fowler NH, Davis RE, Rawal S, et al. Safety and activity of lenalidomide and rituximab in untreated indolent lymphoma: an open-label, phase 2 trial. *Lancet Oncol*. 2014 Nov;15(12):1311-8. doi: 10.1016/S1470-2045(14)70455-3.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C46.0	Kaposi's sarcoma of skin
C46.1	Kaposi's sarcoma of soft tissue
C46.2	Kaposi's sarcoma of palate
C46.3	Kaposi's sarcoma of lymph nodes
C46.4	Kaposi's sarcoma of gastrointestinal sites
C46.50	Kaposi's sarcoma of unspecified lung
C46.51	Kaposi's sarcoma of right lung
C46.52	Kaposi's sarcoma of left lung
C46.7	Kaposi's sarcoma of other sites
C46.9	Kaposi's sarcoma, unspecified
C81.10	Nodular sclerosis classical Hodgkin lymphoma, unspecified site
C81.11	Nodular sclerosis classical Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.12	Nodular sclerosis classical Hodgkin lymphoma, intrathoracic lymph nodes

ICD-10	ICD-10 Description
C81.13	Nodular sclerosis classical Hodgkin lymphoma, intra-abdominal lymph nodes
C81.14	Nodular sclerosis classical Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.15	Nodular sclerosis classical Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.16	Nodular sclerosis classical Hodgkin lymphoma, intrapelvic lymph nodes
C81.17	Nodular sclerosis classical Hodgkin lymphoma, spleen
C81.18	Nodular sclerosis classical Hodgkin lymphoma, lymph nodes of multiple sites
C81.19	Nodular sclerosis classical Hodgkin lymphoma, extranodal and solid organ sites
C81.20	Mixed cellularity classical Hodgkin lymphoma, unspecified site
C81.21	Mixed cellularity classical Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.22	Mixed cellularity classical Hodgkin lymphoma, intrathoracic lymph nodes
C81.23	Mixed cellularity classical Hodgkin lymphoma, intra-abdominal lymph nodes
C81.24	Mixed cellularity classical Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.25	Mixed cellularity classical Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.26	Mixed cellularity classical Hodgkin lymphoma, intrapelvic lymph nodes
C81.27	Mixed cellularity classical Hodgkin lymphoma, spleen
C81.28	Mixed cellularity classical Hodgkin lymphoma, lymph nodes of multiple sites
C81.29	Mixed cellularity classical Hodgkin lymphoma, extranodal and solid organ sites
C81.30	Lymphocyte depleted classical Hodgkin lymphoma, unspecified site
C81.31	Lymphocyte depleted classical Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.32	Lymphocyte depleted classical Hodgkin lymphoma, intrathoracic lymph nodes
C81.33	Lymphocyte depleted classical Hodgkin lymphoma, intra-abdominal lymph nodes
C81.34	Lymphocyte depleted classical Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.35	Lymphocyte depleted classical Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.36	Lymphocyte depleted classical Hodgkin lymphoma, intrapelvic lymph nodes
C81.37	Lymphocyte depleted classical Hodgkin lymphoma, spleen
C81.38	Lymphocyte depleted classical Hodgkin lymphoma, lymph nodes of multiple sites
C81.39	Lymphocyte depleted classical Hodgkin lymphoma, extranodal and solid organ sites
C81.40	Lymphocyte-rich classical Hodgkin lymphoma, unspecified site
C81.41	Lymphocyte-rich classical Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.42	Lymphocyte-rich classical Hodgkin lymphoma, intrathoracic lymph nodes
C81.43	Lymphocyte-rich classical Hodgkin lymphoma, intra-abdominal lymph nodes
C81.44	Lymphocyte-rich classical Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.45	Lymphocyte-rich classical Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.46	Lymphocyte-rich classical Hodgkin lymphoma, intrapelvic lymph nodes

ICD-10	ICD-10 Description
C81.47	Lymphocyte-rich classical Hodgkin lymphoma, spleen
C81.48	Lymphocyte-rich classical Hodgkin lymphoma, lymph nodes of multiple sites
C81.49	Lymphocyte-rich classical Hodgkin lymphoma, extranodal and solid organ sites
C81.70	Other Hodgkin lymphoma, unspecified site
C81.71	Other Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.72	Other Hodgkin lymphoma, intrathoracic lymph nodes
C81.73	Other Hodgkin lymphoma, intra-abdominal lymph nodes
C81.74	Other Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.75	Other Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.76	Other Hodgkin lymphoma, intrapelvic lymph nodes
C81.77	Other Hodgkin lymphoma, spleen
C81.78	Other Hodgkin lymphoma, extranodal and solid organ sites
C81.79	Other Hodgkin lymphoma, extranodal and solid organ sites
C81.90	Hodgkin lymphoma, unspecified, unspecified site
C81.91	Hodgkin lymphoma, unspecified, lymph nodes of head, face, and neck
C81.92	Hodgkin lymphoma, unspecified, intrathoracic lymph nodes
C81.93	Hodgkin lymphoma, unspecified, intra-abdominal lymph nodes
C81.94	Hodgkin lymphoma, unspecified, lymph nodes of axilla and upper limb
C81.95	Hodgkin lymphoma, unspecified, lymph nodes of inguinal region and lower limb
C81.96	Hodgkin lymphoma, unspecified, intrapelvic lymph nodes
C81.97	Hodgkin lymphoma, unspecified, spleen
C81.98	Hodgkin lymphoma, unspecified, lymph nodes of multiple sites
C81.99	Hodgkin lymphoma, unspecified, extranodal and solid organ sites
C82.00	Follicular lymphoma grade I, unspecified site
C82.01	Follicular lymphoma grade I, lymph nodes of head, face and neck
C82.02	Follicular lymphoma, grade I, intrathoracic lymph nodes
C82.03	Follicular lymphoma grade I, intra-abdominal lymph nodes
C82.04	Follicular lymphoma grade I, lymph nodes of axilla and upper limb
C82.05	Follicular lymphoma grade I, lymph nodes of inguinal regional and lower limb
C82.06	Follicular lymphoma grade I, intrapelvic lymph nodes
C82.07	Follicular lymphoma grade I, spleen
C82.08	Follicular lymphoma grade I, lymph nodes of multiple sites
C82.09	Follicular lymphoma grade I, extranodal and solid organ sites
C82.10	Follicular lymphoma grade II, unspecified site

ICD-10	ICD-10 Description
C82.11	Follicular lymphoma grade II, lymph nodes of head, face and neck
C82.12	Follicular lymphoma, grade II, intrathoracic lymph nodes
C82.13	Follicular lymphoma grade II, intra-abdominal lymph nodes
C82.14	Follicular lymphoma grade II, lymph nodes of axilla and upper limb
C82.15	Follicular lymphoma grade II, lymph nodes of inguinal region and lower limb
C82.16	Follicular lymphoma grade II, intrapelvic lymph nodes
C82.17	Follicular lymphoma grade II, spleen
C82.18	Follicular lymphoma grade II, lymph nodes of multiple sites
C82.19	Follicular lymphoma grade II, extranodal and solid organ sites
C82.20	Follicular lymphoma grade III, unspecified, unspecified site
C82.21	Follicular lymphoma grade III, unspecified, lymph nodes of head, face and neck
C82.22	Follicular lymphoma, grade III, unspecified, intrathoracic lymph nodes
C82.23	Follicular lymphoma grade III, unspecified, intra-abdominal lymph nodes
C82.24	Follicular lymphoma grade III, unspecified, lymph nodes of axilla and upper limb
C82.25	Follicular lymphoma grade III, unspecified, lymph nodes of inguinal region and lower limb
C82.26	Follicular lymphoma grade III, unspecified, intrapelvic lymph nodes
C82.27	Follicular lymphoma grade III, unspecified, spleen
C82.28	Follicular lymphoma grade III, unspecified, lymph nodes of multiple sites
C82.29	Follicular lymphoma grade III, unspecified, extranodal and solid organ sites
C82.30	Follicular lymphoma grade IIIa, unspecified site
C82.31	Follicular lymphoma grade IIIa, lymph nodes of head, face and neck
C82.32	Follicular lymphoma, grade IIIa, intrathoracic lymph nodes
C82.33	Follicular lymphoma grade IIIa, intra-abdominal lymph nodes
C82.34	Follicular lymphoma grade IIIa, lymph nodes of axilla and upper limb
C82.35	Follicular lymphoma grade IIIa, lymph nodes of inguinal region and lower limb
C82.36	Follicular lymphoma grade IIIa, intrapelvic lymph nodes
C82.37	Follicular lymphoma grade IIIa, spleen
C82.38	Follicular lymphoma grade IIIa, lymph nodes of multiple sites
C82.39	Follicular lymphoma grade IIIa, extranodal and solid organ sites
C82.40	Follicular lymphoma grade IIIb, unspecified site
C82.41	Follicular lymphoma grade IIIb, lymph nodes of head, face and neck
C82.42	Follicular lymphoma, grade IIIb, intrathoracic lymph nodes
C82.43	Follicular lymphoma grade IIIb, intra-abdominal lymph nodes
C82.44	Follicular lymphoma grade IIIb, lymph nodes of axilla and upper limb

ICD-10	ICD-10 Description
C82.45	Follicular lymphoma grade IIIb, lymph nodes of inguinal region and lower limb
C82.46	Follicular lymphoma grade IIIb, intrapelvic lymph nodes
C82.47	Follicular lymphoma grade IIIb, spleen
C82.48	Follicular lymphoma grade IIIb, lymph nodes of multiple sites
C82.49	Follicular lymphoma grade IIIb, extranodal and solid organ sites
C82.50	Diffuse follicle center lymphoma, unspecified site
C82.51	Diffuse follicle center lymphoma, lymph nodes of head, face and neck
C82.52	Diffuse follicle center lymphoma, intrathoracic lymph nodes
C82.53	Diffuse follicle center lymphoma, intra-abdominal lymph nodes
C82.54	Diffuse follicle center lymphoma, lymph nodes of axilla and upper limb
C82.55	Diffuse follicle center lymphoma, lymph nodes of inguinal region and lower limb
C82.56	Diffuse follicle center lymphoma, intrapelvic lymph nodes
C82.57	Diffuse follicle center lymphoma, spleen
C82.58	Diffuse follicle center lymphoma, lymph nodes of multiple sites
C82.59	Diffuse follicle center lymphoma, extranodal and solid organ sites
C82.60	Cutaneous follicle center lymphoma, unspecified site
C82.61	Cutaneous follicle center lymphoma, lymph nodes of head, face and neck
C82.62	Cutaneous follicle center lymphoma, intrathoracic lymph nodes
C82.63	Cutaneous follicle center lymphoma, intra-abdominal lymph nodes
C82.64	Cutaneous follicle center lymphoma, lymph nodes of axilla and upper limb
C82.65	Cutaneous follicle center lymphoma, lymph nodes of inguinal region and lower limb
C82.66	Cutaneous follicle center lymphoma, intrapelvic lymph nodes
C82.67	Cutaneous follicle center lymphoma, spleen
C82.68	Cutaneous follicle center lymphoma, lymph nodes of multiple sites
C82.69	Cutaneous follicle center lymphoma, extranodal and solid organ sites
C82.80	Other types of follicular lymphoma, unspecified site
C82.81	Other types of follicular lymphoma, lymph nodes of head, face, and neck
C82.82	Other types of follicular lymphoma, intrathoracic lymph nodes
C82.83	Other types of follicular lymphoma, intra-abdominal lymph nodes
C82.84	Other types of follicular lymphoma, lymph nodes of axilla and upper limb
C82.85	Other types of follicular lymphoma, lymph nodes of inguinal region and lower limb
C82.86	Other types of follicular lymphoma, intrapelvic lymph nodes
C82.87	Other types of follicular lymphoma, spleen
C82.88	Other types of follicular lymphoma, lymph nodes of multiple sites

ICD-10	ICD-10 Description
C82.89	Other types of follicular lymphoma, extranodal and solid organ sites
C82.90	Follicular lymphoma, unspecified, unspecified site
C82.91	Follicular lymphoma, unspecified, lymph nodes of head, face and neck
C82.92	Follicular lymphoma, unspecified, intrathoracic lymph nodes
C82.93	Follicular lymphoma, unspecified, intra-abdominal lymph nodes
C82.94	Follicular lymphoma, unspecified, lymph nodes of axilla and upper limb
C82.95	Follicular lymphoma, unspecified lymph nodes of inguinal region and lower limb
C82.96	Follicular lymphoma, unspecified, intrapelvic lymph nodes
C82.97	Follicular lymphoma, unspecified, spleen
C82.98	Follicular lymphoma, unspecified, lymph nodes of multiple sites
C82.99	Follicular lymphoma, unspecified, extranodal and solid organ sites
C83.00	Small cell B-cell lymphoma, unspecified site
C83.01	Small cell B-cell lymphoma, lymph nodes of head, face and neck
C83.02	Small cell B-cell lymphoma, intrathoracic lymph nodes
C83.03	small cell B-cell lymphoma, intra-abdominal lymph nodes
C83.04	Small cell B-cell lymphoma, lymph nodes of axilla and upper limb
C83.05	Small cell B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.06	Small cell B-cell lymphoma, intrapelvic lymph nodes
C83.07	Small cell B-cell lymphoma, spleen
C83.08	Small cell B-cell lymphoma, lymph nodes of multiple sites
C83.09	Small cell B-cell lymphoma, extranodal and solid organ sites
C83.10	Mantle cell lymphoma, unspecified site
C83.11	Mantle cell lymphoma, lymph nodes of head, face and neck
C83.12	Mantle cell lymphoma, intrathoracic lymph nodes
C83.13	Mantle cell lymphoma, intra-abdominal lymph nodes
C83.14	Mantle cell lymphoma, lymph nodes of axilla and upper limb
C83.15	Mantle cell lymphoma, lymph nodes of inguinal region and lower limb
C83.16	Mantle cell lymphoma, intrapelvic lymph nodes
C83.17	Mantle cell lymphoma, spleen
C83.18	Mantle cell lymphoma, lymph nodes of multiple sites
C83.19	Mantle cell lymphoma, extranodal and solid organ sites
C83.30	Diffuse large B-cell lymphoma, unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face and neck
C83.32	Diffuse large B-cell lymphoma intrathoracic lymph nodes

ICD-10	ICD-10 Description
C83.33	Diffuse large B-cell lymphoma, intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma, lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma, intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma, lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma, extranodal and solid organ sites
C83.80	Other non-follicular lymphoma, unspecified site
C83.81	Other non-follicular lymphoma, lymph nodes of head, face and neck
C83.82	Other non-follicular lymphoma, intrathoracic lymph nodes
C83.83	Other non-follicular lymphoma, intra-abdominal lymph nodes
C83.84	Other non-follicular lymphoma, lymph nodes of axilla and upper limb
C83.85	Other non-follicular lymphoma, lymph nodes of inguinal region and lower limb
C83.86	Other non-follicular lymphoma, intrapelvic lymph nodes
C83.87	Other non-follicular lymphoma, spleen
C83.88	Other non-follicular lymphoma, lymph nodes of multiple sites
C83.89	Other non-follicular lymphoma, extranodal and solid organ sites
C83.90	Non-follicular (diffuse) lymphoma, unspecified site
C83.91	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of head, face, and neck
C83.92	Non-follicular (diffuse) lymphoma, unspecified intrathoracic lymph nodes
C83.93	Non-follicular (diffuse) lymphoma, unspecified intra-abdominal lymph nodes
C83.94	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of axilla and upper limb
C83.95	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of inguinal region and lower limb
C83.96	Non-follicular (diffuse) lymphoma, unspecified intrapelvic lymph nodes
C83.97	Non-follicular (diffuse) lymphoma, unspecified spleen
C83.98	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of multiple sites
C83.99	Non-follicular (diffuse) lymphoma, unspecified extranodal and solid organ sites
C84.00	Mycosis fungoides, unspecified site
C84.01	Mycosis fungoides, lymph nodes of head, face and neck
C84.02	Mycosis fungoides, intrathoracic lymph nodes
C84.03	Mycosis fungoides, intra-abdominal lymph nodes
C84.04	Mycosis fungoides, lymph nodes of axilla and upper limb
C84.05	Mycosis fungoides, lymph nodes of inguinal region and lower limb
C84.06	Mycosis fungoides, intrapelvic lymph nodes

ICD-10	ICD-10 Description
C84.07	Mycosis fungoides, spleen
C84.08	Mycosis fungoides, lymph nodes of multiple sites
C84.09	Mycosis fungoides, extranodal and solid organ sites
C84.10	Sézary disease unspecified site
C84.11	Sézary disease lymph nodes of head, face, and neck
C84.12	Sézary disease intrathoracic lymph nodes
C84.13	Sézary disease intra-abdominal lymph nodes
C84.14	Sézary disease lymph nodes of axilla and upper limb
C84.15	Sézary disease lymph nodes of inguinal region and lower limb
C84.16	Sézary disease intrapelvic lymph nodes
C84.17	Sézary disease spleen
C84.18	Sézary disease lymph nodes of multiple sites
C84.19	Sézary disease extranodal and solid organ sites
C84.40	Peripheral T-cell lymphoma, unspecified site
C84.41	Peripheral T-cell lymphoma, lymph nodes of head, face and neck
C84.42	Peripheral T-cell lymphoma, intrathoracic lymph nodes
C84.43	Peripheral T-cell lymphoma, intra-abdominal lymph nodes
C84.44	Peripheral T-cell lymphoma, lymph nodes of axilla and upper limb
C84.45	Peripheral T-cell lymphoma, lymph nodes of inguinal region and lower limb
C84.46	Peripheral T-cell lymphoma, intrapelvic lymph nodes
C84.47	Peripheral T-cell lymphoma, spleen
C84.48	Peripheral T-cell lymphoma, lymph nodes of multiple sites
C84.49	Peripheral T-cell lymphoma, extranodal and solid organ sites
C84.90	Mature T/NK-cell lymphomas, unspecified, unspecified site
C84.91	Mature T/NK-cell lymphomas, unspecified, lymph nodes of head, face and neck
C84.92	Mature T/NK-cell lymphomas, unspecified, intrathoracic lymph nodes
C84.93	Mature T/NK-cell lymphomas, unspecified, intra-abdominal lymph nodes
C84.94	Mature T/NK-cell lymphomas, unspecified, lymph nodes of axilla and upper limb
C84.95	Mature T/NK-cell lymphomas, unspecified, lymph nodes of inguinal region and lower limb
C84.96	Mature, T/NK-cell lymphomas, unspecified, intrapelvic lymph nodes
C84.97	Mature T/NK-cell lymphomas, unspecified, spleen
C84.98	Mature T/NK-cell lymphomas, unspecified, lymph nodes of multiple sites
C84.99	Mature T/NK-cell lymphomas, unspecified, extranodal and solid organ sites
C84.Z0	Other mature T/NK-cell lymphomas, unspecified site

ICD-10	ICD-10 Description
C84.Z1	Other mature T/NK-cell lymphomas, lymph nodes of head, face and neck
C84.Z2	Other mature T/NK-cell lymphomas, intrathoracic lymph nodes
C84.Z3	Other mature T/NK-cell lymphomas, intra-abdominal lymph nodes
C84.Z4	Other mature T/NK-cell lymphomas, lymph nodes of axilla and upper limb
C84.Z5	Other mature T/NK-cell lymphomas, lymph nodes of inguinal region and lower limb
C84.Z6	Other mature T/NK-cell lymphomas, intrapelvic lymph nodes
C84.Z7	Other mature T/NK-cell lymphomas, spleen
C84.Z8	Other mature T/NK-cell lymphomas, lymph nodes of multiple sites
C84.Z9	Other mature T/NK-cell lymphomas, extranodal and solid organ sites
C85.10	Unspecified B-cell lymphoma, unspecified site
C85.11	Unspecified B-cell lymphoma, lymph nodes of head, face, and neck
C85.12	Unspecified B-cell lymphoma, intrathoracic lymph nodes
C85.13	Unspecified B-cell lymphoma, intra-abdominal lymph nodes
C85.14	Unspecified B-cell lymphoma, lymph nodes of axilla and upper limb
C85.15	Unspecified B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.16	Unspecified B-cell lymphoma, intrapelvic lymph nodes
C85.17	Unspecified B-cell lymphoma, spleen
C85.18	Unspecified B-cell lymphoma, lymph nodes of multiple sites
C85.19	Unspecified B-cell lymphoma, extranodal and solid organ sites
C85.20	Mediastinal (thymic) large B-cell lymphoma, unspecified sites
C85.21	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck
C85.22	Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes
C85.23	Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes
C85.24	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb
C85.25	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.26	Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes
C85.27	Mediastinal (thymic) large B-cell lymphoma, spleen
C85.28	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites
C85.29	Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites
C85.80	Other specified types of non-Hodgkin lymphoma, unspecified site
C85.81	Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face and neck
C85.82	Other specified types of non-Hodgkin lymphoma, intrathoracic lymph nodes
C85.83	Other specified types of non-Hodgkin lymphoma, intra-abdominal lymph nodes
C85.84	Other specified types of non-Hodgkin lymphoma, lymph nodes of axilla and upper limb

ICD-10	ICD-10 Description
C85.85	Other specified types of non-Hodgkin lymphoma, lymph nodes of inguinal region of lower limb
C85.86	Other specified types of non-Hodgkin lymphoma, intrapelvic lymph nodes
C85.87	Other specified types of non-Hodgkin lymphoma, spleen
C85.88	Other specified types of non-Hodgkin lymphoma, lymph nodes of multiple sites
C85.89	Other specified types of non-Hodgkin lymphoma, extranodal and solid organ sites
C85.99	Non-Hodgkin lymphoma, unspecified, extranodal and solid organ sites
C86.1	Hepatosplenic T-cell lymphoma
C86.2	Enteropathy-type (intestinal) T-cell lymphoma
C86.5	Angioimmunoblastic T-cell lymphoma
C86.6	Primary cutaneous CD30-positive T-cell proliferations
C88.4	Extranodal marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue (MALT-lymphoma)
C90.00	Multiple myeloma not having achieved remission
C90.02	Multiple myeloma in relapse
C90.10	Plasma cell leukemia not having achieved remission
C90.12	Plasma cell leukemia in relapse
C90.20	Extramedullary plasmacytoma not having achieved remission
C90.22	Extramedullary plasmacytoma in relapse
C90.30	Solitary plasmacytoma not having achieved remission
C90.32	Solitary plasmacytoma in relapse
C91.10	Chronic lymphocytic leukemia of B-cell type not having achieved remission
C91.12	Chronic lymphocytic leukemia of B-cell type in relapse
C91.50	Adult T-cell lymphoma/leukemia (HTLV-1-associated) not having achieved remission
C91.52	Adult T-cell lymphoma/leukemia (HTLV-1-associated) in relapse
C93.10	Chronic myelomonocytic leukemia not having achieved remission
C94.40	Acute panmyelosis with myelofibrosis not having achieved remission
C94.41	Acute panmyelosis with myelofibrosis, in remission
C94.42	Acute panmyelosis with myelofibrosis, in relapse
C94.6	Myelodysplastic disease, not classified
C96.0	Multifocal and multisystemic (disseminated) Langerhans-cell histiocytosis
C96.2	Malignant mast cell neoplasm
C96.5	Multifocal and unisystemic Langerhans-cell histiocytosis
C96.6	Unifocal Langerhans-cell histiocytosis
C96.9	Malignant neoplasm of lymphoid, hematopoietic and related tissue, unspecified
C96.Z	Other specified malignant neoplasms of lymphoid, hematopoietic and related tissue

ICD-10	ICD-10 Description
D46.0	Refractory anemia without ring sideroblasts, so stated
D46.1	Refractory anemia with ring sideroblasts
D46.20	Refractory anemia with excess of blasts, unspecified
D46.21	Refractory anemia with excess of blasts 1
D46.4	Refractory anemia, unspecified
D46.9	Myelodysplastic syndrome, unspecified
D46.A	Refractory cytopenia with multilineage dysplasia
D46.B	Refractory cytopenia with multilineage dysplasia and ring sideroblasts
D46.C	Myelodysplastic syndrome with isolated del(5q) chromosomal abnormality
D46.Z	Other myelodysplastic syndromes
D47.1	Chronic myeloproliferative disease
D47.4	Osteomyelofibrosis
D47.7	Other specified neoplasms of uncertain or unknown behavior of lymphoid, hematopoietic, and related tissue
D47.9	Neoplasm of uncertain behavior of lymphoid, hematopoietic and related tissue, unspecified
D47.Z1	Post-transplant lymphoproliferative disorder (PTLD)
D47.Z2	Castleman disease
D75.81	Myelofibrosis
E31.9	Polyglandular dysfunction, unspecified
E85.81	Light chain (AL) amyloidosis
E85.89	Other amyloidosis
E85.9	Amyloidosis, unspecified
G62.9	Polyneuropathy, unspecified
G90.9	Disorder of the skin and subcutaneous tissue, unspecified
L98.9	Disorder of the skin and subcutaneous tissue, unspecified
Z85.79	Personal history of other malignant neoplasms of lymphoid, hematopoietic and related tissues

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC