

## Stelara® (ustekinumab) (Intravenous/Subcutaneous)

Document Number: IH-0117

Last Review Date: 08/04/2020

Date of Origin: 02/15/2011

Dates Reviewed: 03/2011, 06/2011, 09/2011, 12/2011, 03/2012, 06/2012, 09/2012, 03/2013, 06/2013, 09/2013, 11/2013, 12/2013, 03/2014, 06/2014, 09/2014, 12/2014, 03/2015, 06/2015, 09/2015, 12/2015, 03/2016, 06/2016, 9/2016, 10/2016, 11/2016, 03/2017, 06/2017, 09/2017, 10/2017, 03/2018, 06/2018, 10/2018, 10/2019, 12/2019, 07/2020, 08/2020

### I. Length of Authorization

#### Crohn's Disease and Ulcerative Colitis:

Coverage will be provided for 8 weeks initially and may be renewed in 6 month intervals thereafter.

#### All other indications:

Coverage will be provided for 6 months and may be renewed.

### II. Dosing Limits

#### A. Quantity Limit (max daily dose) [NDC Unit]:

<p><b><u>Subcutaneous</u></b></p> <ul style="list-style-type: none"> <li>• Stelara 45 mg vial/prefilled syringe:             <ul style="list-style-type: none"> <li>○ Loading: 1 syringe at weeks 0 &amp; 4</li> <li>○ Maintenance: 1 syringe every 12 weeks</li> </ul> </li> <li>• Stelara 90 mg prefilled syringe:             <ul style="list-style-type: none"> <li>○ Loading: 1 syringe at weeks 0 &amp; 4</li> <li>○ Maintenance: 1 syringe every 8 weeks</li> </ul> </li> </ul>
<p><b><u>Intravenous</u></b></p> <ul style="list-style-type: none"> <li>• Stelara 130 mg (5 mg/mL) single-dose vial: 4 vials</li> </ul>

#### B. Max Units (per dose and over time) [HCPCS Unit]:

Indication	Max Units
Plaque Psoriasis & Psoriatic Arthritis with co-existent moderate-severe Plaque Psoriasis	<p><u>Subcutaneous Loading (J3357):</u></p> <ul style="list-style-type: none"> <li>• 90 billable units at weeks 0 &amp; 4; maintenance dosing 12 weeks later</li> </ul> <p><u>Subcutaneous Maintenance (J3357):</u></p> <ul style="list-style-type: none"> <li>• 90 billable units every 12 weeks</li> </ul>
Psoriatic Arthritis	<p><u>Subcutaneous Loading (J3357):</u></p>

Indication	Max Units
	<ul style="list-style-type: none"> <li>45 billable units at weeks 0 &amp; 4; maintenance dosing 12 weeks later</li> </ul> <u>Subcutaneous Maintenance (J3357):</u> <ul style="list-style-type: none"> <li>45 billable units every 12 weeks</li> </ul>
Crohn's Disease & Ulcerative Colitis	<u>Intravenous Induction (J3358):</u> <ul style="list-style-type: none"> <li>520 billable units</li> </ul> <u>Subcutaneous Maintenance (J3357):</u> <ul style="list-style-type: none"> <li>90 billable units 8 weeks after induction &amp; every 8 weeks thereafter</li> </ul>

### III. Initial Approval Criteria <sup>1</sup>

- The use of samples and free goods do not qualify as an established clinical response.

Coverage is provided in the following conditions:

- Patient is at least 18 years of age (unless otherwise specified); **AND**
- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**
- Patient is up to date with all vaccinations, in accordance with current vaccination guidelines, prior to initiating therapy; **AND**

#### Universal Criteria <sup>1</sup>

- Patient has been evaluated and screened for the presence of latent (tuberculosis) TB infection prior to initiating treatment and will receive ongoing monitoring for presence of TB during treatment; **AND**
- Patient does not have an active infection, including clinically important localized infections; **AND**
- Therapy will not be administered concurrently with live vaccines; **AND**
- Patient is not on concurrent treatment with a TNF-inhibitor, biologic response modifier or other non-biologic agent (i.e., apremilast, tofacitinib, baricitinib); **AND**

#### Plaque Psoriasis † <sup>1-5,25,26,30</sup>

- Patient is at least 6 years of age; **AND**
- Patient has moderate to severe plaque psoriasis for at least 6 months with at least one of the following:
  - Involvement of at least 3% of body surface area (BSA); **OR**
  - Psoriasis Area and Severity Index (PASI) score of 10 or greater; **OR**
  - Incapacitation or serious emotional consequences due to plaque location (i.e. hands, feet, head and neck, or genitalia) or with intractable pruritis; **AND**
- Patient did not respond adequately (or is not a candidate) to a 3 month minimum trial of topical agents (i.e., anthralin, coal tar preparations, corticosteroids, emollients, immunosuppressives, keratolytics, retinoic acid derivatives, and/or vitamin D analogues); **AND**

- Patient did not respond adequately (or is not a candidate) to a 3 month minimum trial of at least one systemic agent (i.e., immunosuppressives, retinoic acid derivatives, and/or methotrexate); **OR**
- Patient did not respond adequately (or is not a candidate\*) to a 3 month minimum trial of phototherapy (i.e., psoralens with UVA light [PUVA] or UVB with coal tar or dithranol)

#### **Psoriatic Arthritis (PsA) †<sup>1,6-9,15,16,17,27</sup>**

- Patient has documented moderate to severe active disease ; **AND**
  - For patients with predominantly axial disease OR active enthesitis and/or dactylitis, an adequate trial and failure of at least TWO (2) non-steroidal anti-inflammatory agents (NSAIDs), unless use is contraindicated; **OR**
  - For patients with peripheral arthritis, a trial and failure of at least a 3 month trial of ONE oral disease-modifying anti-rheumatic agent (DMARD) such as methotrexate, azathioprine, sulfasalazine, or hydroxychloroquine

#### **Crohn's Disease †<sup>1,10-12,14,18,24</sup>**

- Documented moderate to severely active disease; **AND**
- Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum (3) month trial of corticosteroids or immunomodulators (e.g. azathioprine, 6-mercaptopurine, or methotrexate)

#### **Ulcerative Colitis †<sup>1,13,19-23,29</sup>**

- Documented moderate to severely active disease; **AND**
- Documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum (3) month trial of corticosteroids or immunomodulators (e.g. azathioprine, 6-mercaptopurine, or methotrexate)

**\*Examples of contraindications to phototherapy (PUVA or UVB) include the following:<sup>31,32</sup>**

- Xeroderma pigmentosum
- Pregnancy or lactation (PUVA only)
- Lupus Erythematosus
- History of one of the following: photosensitivity diseases (e.g., chronic actinic dermatitis, solar urticaria), melanoma, non-melanoma skin cancer, treatment with arsenic or ionizing radiation
- Immunosuppression in an organ transplant patient

† FDA Approved Indication(s); ‡ Compendia recommended indication(s); Ⓢ Orphan Drug

## **IV. Renewal Criteria<sup>1</sup>**

Coverage can be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**

- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: serious infections, malignancy, severe hypersensitivity reactions, reversible posterior leukoencephalopathy syndrome (RPLS), non-infectious pneumonia, etc; **AND**

#### **Plaque Psoriasis** <sup>15,16</sup>

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as redness, thickness, scaliness, and/or the amount of surface area involvement (a total BSA involvement  $\leq 1\%$ ), and/or an improvement on a disease activity scoring tool [e.g. a 75% reduction in the PASI score from when treatment started (PASI 75) or a 50% reduction in the PASI score (PASI 50) and a four-point reduction in the DLQI from when treatment started].

#### **Psoriatic Arthritis (PsA)** <sup>14</sup>

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts and/or an improvement on a disease activity scoring tool [e.g. defined as an improvement in at least 2 of the 4 Psoriatic Arthritis Response Criteria (PsARC), 1 of which must be joint tenderness or swelling score, with no worsening in any of the 4 criteria].

#### **Crohn's Disease** <sup>13</sup>

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as endoscopic activity, number of liquid stools, presence and severity of abdominal pain, presence of abdominal mass, body weight compared to IBW, hematocrit, presence of extra intestinal complications, use of anti-diarrheal drugs, tapering or discontinuation of corticosteroid therapy, and/or an improvement on a disease activity scoring tool [e.g. an improvement on the Crohn's Disease Activity Index (CDAI) score or the Harvey-Bradshaw Index score].

#### **Ulcerative Colitis** <sup>19-23</sup>

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as stool frequency, rectal bleeding, and/or endoscopic activity, tapering or discontinuation of corticosteroid therapy, normalization of C-reactive protein (CRP) or fecal calprotectin (FC), and/or an improvement on a disease activity scoring tool [e.g. an improvement on the Ulcerative Colitis Endoscopic Index of Severity (UCEIS) score or the Mayo Score].

## V. Dosage/Administration

Indication	Dose
Plaque Psoriasis & Psoriatic Arthritis with co-existent moderate-severe Plaque Psoriasis	<u>Adult Subcutaneous Loading Dose:</u> <ul style="list-style-type: none"> <li>&lt;100 kg: 45 mg at weeks 0 &amp; 4, then begin maintenance dosing 12 weeks later</li> <li>&gt;100 kg: 90 mg at weeks 0 &amp; 4, then begin maintenance dosing 12 weeks later</li> </ul> <u>Adult Subcutaneous Maintenance Dose:</u> <ul style="list-style-type: none"> <li>&lt;100 kg: 45 mg every 12 weeks</li> <li>&gt;100 kg: 90 mg every 12 weeks</li> </ul>
	<u>Pediatric Subcutaneous Loading Dose:</u> <ul style="list-style-type: none"> <li>&lt;60 kg: 0.75 mg/kg at weeks 0 &amp; 4, then begin maintenance dosing 12 weeks later</li> <li>60 – 100 kg: 45 mg at weeks 0 &amp; 4, then begin maintenance dosing 12 weeks later</li> <li>&gt;100 kg: 90 mg at weeks 0 &amp; 4, then begin maintenance dosing 12 weeks later</li> </ul> <u>Pediatric Subcutaneous Maintenance Dose:</u> <ul style="list-style-type: none"> <li>&lt;60 kg: 0.75 mg/kg every 12 weeks</li> <li>60 – 100 kg: 45 mg every 12 weeks</li> <li>&gt;100 kg: 90 mg every 12 weeks</li> </ul>
Psoriatic Arthritis	<u>Subcutaneous Loading Dose:</u> <ul style="list-style-type: none"> <li>45 mg at weeks 0 &amp; 4, then begin maintenance dosing 12 weeks later</li> </ul> <u>Subcutaneous Maintenance Dose:</u> <ul style="list-style-type: none"> <li>45 mg every 12 weeks</li> </ul>
Crohn's Disease & Ulcerative Colitis	<u>Intravenous Induction Dose (one-time only):</u> <ul style="list-style-type: none"> <li>≤ 55 kg: 260 mg</li> <li>&gt; 55 kg to 85 kg: 390 mg</li> <li>&gt; 85 kg: 520 mg</li> </ul> <u>Subcutaneous Maintenance Dose:</u> <ul style="list-style-type: none"> <li>90 mg given 8 weeks after the initial IV dose, then every 8 weeks thereafter</li> </ul>

## VI. Billing Code/Availability Information

### HCPCS Code:

- J3357 – Ustekinumab, for subcutaneous injection, 1 mg; 1 billable unit = 1 mg
- J3358 – Ustekinumab, for intravenous injection, 1 mg; 1 billable unit = 1 mg

### NDC:

- Subcutaneous
  - Stelara 45 mg single-dose vial (SDV) and prefilled (PF) syringe: 57894-0060-xx
  - Stelara 90 mg prefilled (PF) syringe: 57894-0061-xx
- Intravenous
  - Stelara 130 mg (5 mg/mL) single-dose vial (SDV): 57894-0054-xx

## VII. References

- Stelara [package insert]. Horsham, PA; Janssen Biotech, Inc; March 2020. Accessed June 2020.

2. Leonardi CL, Kimball AB, Papp KA, et al, “Efficacy and Safety of Ustekinumab, a Human Interleukin-12/23 Monoclonal Antibody, in Patients With Psoriasis: 76-Week Results from a Randomised, Double-Blind, Placebo-Controlled Trial (PHOENIX 1),” *Lancet*, 2008, 371(9625): 1665-74.
3. Papp KA, Langley RG, Lebwohl M, et al, “Efficacy and Safety of Ustekinumab, a Human Interleukin-12/23 Monoclonal Antibody, in Patients With Psoriasis: 52-Week Results from a Randomised, Double-Blind, Placebo-Controlled Trial (PHOENIX 2),” *Lancet*, 2008, 371(9625): 1675-84.
4. Hsu S, Papp KA, Lebwohl MG, et al. Consensus guidelines for the management of plaque psoriasis. *Arch Dermatol*. 2012 Jan;148(1):95-102.
5. Papp KA, Griffiths CE, Gordon K, et al. Long-term safety of ustekinumab in patients with moderate-to-severe psoriasis: final results from 5 years of follow-up. *Br J Dermatol*. 2013 Apr;168(4):844-54.
6. Menter A, Gottlieb A, Feldman SR, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 1. Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2008 May;58(5):826-50. doi: 10.1016/j.jaad.2008.02.039.
7. Gottlieb A, Korman NJ, Gordon KB, Feldman SR, Lebwohl M, Koo JY, Van Voorhees AS, Elmets CA, Leonardi CL, Beutner KR, Bhushan R, Menter A. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 2. Psoriatic arthritis: overview and guidelines of care for treatment with an emphasis on the biologics. *J Am Acad Dermatol* 2008 May;58(5):851-64.
8. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis*. 2015 Dec 7. pii: annrheumdis-2015-208337. doi: 10.1136/annrheumdis-2015-208337.
9. Sing JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Care Res*, 71: 2-29. doi:10.1002/acr.23789.
10. Lichtenstein GR, Hanauer SB, Sandborn WJ, Practice Parameters Committee of American College of Gastroenterology. Management of Crohn’s disease in adults. *Am J Gastroenterol*. 2009;104(2):465.
11. Terdiman JP, Gruss CB, Heidelbaugh JJ, et al. American Gastroenterological Association Institute guideline on the use of thiopurines, methotrexate, and anti-TNF- $\alpha$  biologic drugs for the induction and maintenance of remission in inflammatory Crohn's disease. *Gastroenterology*. 2013 Dec;145(6):1459-63. doi: 10.1053/j.gastro.2013.10.047.
12. Gomollón F, Dignass A, Annese V, et al. EUROPEAN Evidence-based consensus on the diagnosis and management of Crohn's disease 2016: Part 1: Diagnosis and medical management. *J Crohns Colitis*. 2016 Sep 22. pii: jjw168.
13. Harbord M, Eliakim R, Bettenworth D, et al. Third European Evidence-based Consensus on Diagnosis and Management of Ulcerative Colitis. Part 2: Current Management. *J Crohns Colitis*. 2017 Jan 28. doi: 10.1093/ecco-jcc/jjx009.

14. National Institute for Health and Care Excellence. NICE 2012. Crohn's Disease: Management. Published 10 October 2012. Clinical Guideline [CG152]. <https://www.nice.org.uk/guidance/cg152/resources/crohns-disease-management-pdf-35109627942085>.
15. National Institute for Health and Care Excellence. NICE 2017. Certolizumab pegol and secukinumab for treating active psoriatic arthritis after inadequate response to DMARDs. Published 24 May 2017. Technology Appraisal Guidance [TA445]. <https://www.nice.org.uk/guidance/TA445/chapter/1-Recommendations>. Accessed August 2017.
16. National Institute for Health and Care Excellence. NICE 2008. Infliximab for the treatment of adults with psoriasis. Published 23 January 2008. Technology Appraisal Guidance [TA134]. <https://www.nice.org.uk/guidance/ta134/resources/infliximab-for-the-treatment-of-adults-with-psoriasis-pdf-82598193811141>.
17. Smith CH, Jabbar-Lopez ZK, Yiu ZK, et al. British Association of Dermatologists guidelines for biologic therapy for psoriasis 2017. *Br J Dermatol*. 2017 Sep;177(3):628-636. doi: 10.1111/bjd.15665.
18. Lichtenstein GR, Loftus EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol* 2018; 113:481–517; doi: 10.1038/ajg.2018.27
19. Sands BE, Sandborn WJ, Panaccione R, et al. UNIFI Study Group. Ustekinumab as Induction and Maintenance Therapy for Ulcerative Colitis. *N Engl J Med*. 2019 Sep 26;381(13):1201-1214. doi: 10.1056/NEJMoa1900750.
20. Lewis JD, Chuai S, Nessel L, et al. Use of the Non-invasive Components of the Mayo Score to Assess Clinical Response in Ulcerative Colitis. *Inflamm Bowel Dis*. 2008 Dec; 14(12): 1660–1666. doi: 10.1002/ibd.20520
21. Paine ER. Colonoscopic evaluation in ulcerative colitis. *Gastroenterol Rep (Oxf)*. 2014 Aug; 2(3): 161–168.
22. Walsh AJ, Bryant RV, Travis SPL. Current best practice for disease activity assessment in IBD. *Nature Reviews Gastroenterology & Hepatology* 13, 567–579 (2016) doi:10.1038/nrgastro.2016.128
23. Kornbluth, A, Sachar, DB; Practice Parameters Committee of the American College of Gastroenterology. Ulcerative colitis practice guidelines in adults: American College Of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol*. 2010 Mar;105(3):501-23.
24. Feagan BG, Sandborn WJ, Gasink C, UNITI–IM–UNITI Study Group et al. Ustekinumab as Induction and Maintenance Therapy for Crohn's Disease. *N Engl J Med*. 2016 Nov 17;375(20):1946-1960. doi: 10.1056/NEJMoa1602773.
25. Leonardi CL, Kimball AB, Papp KA, PHOENIX 1 study investigators. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 76-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 1). *Lancet*. 2008;371(9625):1665.
26. Papp KA, Langley RG, Lebwohl M, PHOENIX 2 study investigators. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 52-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 2). *Lancet*. 2008;371(9625):1675.

27. Landells I, Marano C, Hsu MC, et al. Ustekinumab in adolescent patients age 12 to 17 years with moderate-to-severe plaque psoriasis: results of the randomized phase 3 CADMUS study. *J Am Acad Dermatol*. 2015;73(4):594.
28. McInnes IB, Kavanaugh A, Gottlieb AB, PSUMMIT 1 Study Group. Efficacy and safety of ustekinumab in patients with active psoriatic arthritis: 1 year results of the phase 3, multicentre, double-blind, placebo-controlled PSUMMIT 1 trial. *Lancet*. 2013;382(9894):780. Epub 2013 Jun 13.
29. Ritchlin C, Rahman P, Kavanaugh A, PSUMMIT 2 Study Group. Efficacy and safety of the anti-IL-12/23 p40 monoclonal antibody, ustekinumab, in patients with active psoriatic arthritis despite conventional non-biological and biological anti-tumour necrosis factor therapy: 6-month and 1-year results of the phase 3, multicentre, double-blind, placebo-controlled, randomised PSUMMIT 2 trial. *Ann Rheum Dis*. 2014;73(6):990. Epub 2014 Jan 30.
30. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019 Feb 13. pii: S0190-9622(18)33001-9. <https://doi.org/10.1016/j.jaad.2018.11.057>.
31. Richard EG. (2019). Psoralen plus ultraviolet A (PUVA) photochemotherapy. In Elmetts CA, Corona R (Eds.), *UptoDate*. Available from [https://www.uptodate.com/contents/psoralen-plus-ultraviolet-a-puva-photochemotherapy?sectionName=Skin%20cancer&search=psoriasis%20phototherapy&topicRef=5666&anchor=H31513976&source=see\\_link#H2099103](https://www.uptodate.com/contents/psoralen-plus-ultraviolet-a-puva-photochemotherapy?sectionName=Skin%20cancer&search=psoriasis%20phototherapy&topicRef=5666&anchor=H31513976&source=see_link#H2099103).
32. Honigsman H. (2018). UVB therapy (broadband and narrowband). In Elmetts CA, Corona R (Eds.), *UptoDate*. Available from [https://www.uptodate.com/contents/uvb-therapy-broadband-and-narrowband?sectionName=SHORT-%20AND%20LONG-TERM%20ADVERSE%20EFFECTS&search=psoriasis%20phototherapy&topicRef=5666&anchor=H10844620&source=see\\_link#H10844627](https://www.uptodate.com/contents/uvb-therapy-broadband-and-narrowband?sectionName=SHORT-%20AND%20LONG-TERM%20ADVERSE%20EFFECTS&search=psoriasis%20phototherapy&topicRef=5666&anchor=H10844620&source=see_link#H10844627).
33. Noridian Healthcare Solutions, LLC. Local Coverage Articles (LCA): Chemotherapy Administration (A52991). Centers for Medicare & Medicaid Services, Inc. Updated on 12/16/2019 with effective date 1/01/2020. Accessed June 2020.
34. Noridian Healthcare Solutions, LLC. Local Coverage Articles (LCA): Chemotherapy Administration (A52953). Centers for Medicare & Medicaid Services, Inc. Updated on 12/16/2019 with effective date 1/01/2020. Accessed June 2020.

## Appendix 1 – Covered Diagnosis Codes

### Subcutaneous (J3357)

ICD-10	ICD-10 Description
K50.00	Crohn's disease of small intestine without complications
K50.011	Crohn's disease of small intestine with rectal bleeding
K50.012	Crohn's disease of small intestine with intestinal obstruction
K50.013	Crohn's disease of small intestine with fistula
K50.014	Crohn's disease of small intestine with abscess



ICD-10	ICD-10 Description
K50.018	Crohn's disease of small intestine with other complication
K50.019	Crohn's disease of small intestine with unspecified complications
K50.10	Crohn's disease of large intestine without complications
K50.111	Crohn's disease of large intestine with rectal bleeding
K50.112	Crohn's disease of large intestine with intestinal obstruction
K50.113	Crohn's disease of large intestine with fistula
K50.114	Crohn's disease of large intestine with abscess
K50.118	Crohn's disease of large intestine with other complication
K50.119	Crohn's disease of large intestine with unspecified complications
K50.80	Crohn's disease of both small and large intestine without complications
K50.811	Crohn's disease of both small and large intestine with rectal bleeding
K50.812	Crohn's disease of both small and large intestine with intestinal obstruction
K50.813	Crohn's disease of both small and large intestine with fistula
K50.814	Crohn's disease of both small and large intestine with abscess
K50.818	Crohn's disease of both small and large intestine with other complication
K50.819	Crohn's disease of both small and large intestine with unspecified complications
K50.90	Crohn's disease, unspecified, without complications
K50.911	Crohn's disease, unspecified, with rectal bleeding
K50.912	Crohn's disease, unspecified, with intestinal obstruction
K50.913	Crohn's disease, unspecified, with fistula
K50.914	Crohn's disease, unspecified, with abscess
K50.918	Crohn's disease, unspecified, with other complication
K50.919	Crohn's disease, unspecified, with unspecified complications
K51.00	Ulcerative (chronic) pancolitis without complications
K51.011	Ulcerative (chronic) pancolitis with rectal bleeding
K51.012	Ulcerative (chronic) pancolitis with intestinal obstruction
K51.013	Ulcerative (chronic) pancolitis with fistula
K51.014	Ulcerative (chronic) pancolitis with abscess
K51.018	Ulcerative (chronic) pancolitis with other complication
K51.019	Ulcerative (chronic) pancolitis with unspecified complications
K51.20	Ulcerative (chronic) proctitis without complications
K51.211	Ulcerative (chronic) proctitis with rectal bleeding
K51.212	Ulcerative (chronic) proctitis with intestinal obstruction
K51.213	Ulcerative (chronic) proctitis with fistula

ICD-10	ICD-10 Description
K51.214	Ulcerative (chronic) proctitis with abscess
K51.218	Ulcerative (chronic) proctitis with other complication
K51.219	Ulcerative (chronic) proctitis with unspecified complications
K51.30	Ulcerative (chronic) rectosigmoiditis without complications
K51.311	Ulcerative (chronic) rectosigmoiditis with rectal bleeding
K51.312	Ulcerative (chronic) rectosigmoiditis with intestinal obstruction
K51.313	Ulcerative (chronic) rectosigmoiditis with fistula
K51.314	Ulcerative (chronic) rectosigmoiditis with abscess
K51.318	Ulcerative (chronic) rectosigmoiditis with other complication
K51.319	Ulcerative (chronic) rectosigmoiditis with unspecified complications
K51.50	Left sided colitis without complications
K51.511	Left sided colitis with rectal bleeding
K51.512	Left sided colitis with intestinal obstruction
K51.513	Left sided colitis with fistula
K51.514	Left sided colitis with abscess
K51.518	Left sided colitis with other complication
K51.519	Left sided colitis with unspecified complications
K51.80	Other ulcerative colitis without complications
K51.811	Other ulcerative colitis with rectal bleeding
K51.812	Other ulcerative colitis with intestinal obstruction
K51.813	Other ulcerative colitis with fistula
K51.814	Other ulcerative colitis with abscess
K51.818	Other ulcerative colitis with other complication
K51.819	Other ulcerative colitis with unspecified complications
K51.90	Ulcerative colitis, unspecified, without complications
K51.911	Ulcerative colitis, unspecified with rectal bleeding
K51.912	Ulcerative colitis, unspecified with intestinal obstruction
K51.913	Ulcerative colitis, unspecified with fistula
K51.914	Ulcerative colitis, unspecified with abscess
K51.918	Ulcerative colitis, unspecified with other complication
K51.919	Ulcerative colitis, unspecified with unspecified complications
K52.1	Toxic gastroenteritis and colitis
L40.0	Psoriasis vulgaris
L40.50	Arthropathic psoriasis, unspecified

ICD-10	ICD-10 Description
L40.51	Distal interphalangeal psoriatic arthropathy
L40.52	Psoriatic arthritis mutilans
L40.53	Psoriatic spondylitis
L40.59	Other psoriatic arthropathy
R19.7	Diarrhea, unspecified

### Intravenous (J3358)

ICD-10	ICD-10 Description
K50.00	Crohn's disease of small intestine without complications
K50.011	Crohn's disease of small intestine with rectal bleeding
K50.012	Crohn's disease of small intestine with intestinal obstruction
K50.013	Crohn's disease of small intestine with fistula
K50.014	Crohn's disease of small intestine with abscess
K50.018	Crohn's disease of small intestine with other complication
K50.019	Crohn's disease of small intestine with unspecified complications
K50.10	Crohn's disease of large intestine without complications
K50.111	Crohn's disease of large intestine with rectal bleeding
K50.112	Crohn's disease of large intestine with intestinal obstruction
K50.113	Crohn's disease of large intestine with fistula
K50.114	Crohn's disease of large intestine with abscess
K50.118	Crohn's disease of large intestine with other complication
K50.119	Crohn's disease of large intestine with unspecified complications
K50.80	Crohn's disease of both small and large intestine without complications
K50.811	Crohn's disease of both small and large intestine with rectal bleeding
K50.812	Crohn's disease of both small and large intestine with intestinal obstruction
K50.813	Crohn's disease of both small and large intestine with fistula
K50.814	Crohn's disease of both small and large intestine with abscess
K50.818	Crohn's disease of both small and large intestine with other complication
K50.819	Crohn's disease of both small and large intestine with unspecified complications
K50.90	Crohn's disease, unspecified, without complications
K50.911	Crohn's disease, unspecified, with rectal bleeding
K50.912	Crohn's disease, unspecified, with intestinal obstruction
K50.913	Crohn's disease, unspecified, with fistula

ICD-10	ICD-10 Description
K50.914	Crohn's disease, unspecified, with abscess
K50.918	Crohn's disease, unspecified, with other complication
K50.919	Crohn's disease, unspecified, with unspecified complications
K51.00	Ulcerative (chronic) pancolitis without complications
K51.011	Ulcerative (chronic) pancolitis with rectal bleeding
K51.012	Ulcerative (chronic) pancolitis with intestinal obstruction
K51.013	Ulcerative (chronic) pancolitis with fistula
K51.014	Ulcerative (chronic) pancolitis with abscess
K51.018	Ulcerative (chronic) pancolitis with other complication
K51.019	Ulcerative (chronic) pancolitis with unspecified complications
K51.20	Ulcerative (chronic) proctitis without complications
K51.211	Ulcerative (chronic) proctitis with rectal bleeding
K51.212	Ulcerative (chronic) proctitis with intestinal obstruction
K51.213	Ulcerative (chronic) proctitis with fistula
K51.214	Ulcerative (chronic) proctitis with abscess
K51.218	Ulcerative (chronic) proctitis with other complication
K51.219	Ulcerative (chronic) proctitis with unspecified complications
K51.30	Ulcerative (chronic) rectosigmoiditis without complications
K51.311	Ulcerative (chronic) rectosigmoiditis with rectal bleeding
K51.312	Ulcerative (chronic) rectosigmoiditis with intestinal obstruction
K51.313	Ulcerative (chronic) rectosigmoiditis with fistula
K51.314	Ulcerative (chronic) rectosigmoiditis with abscess
K51.318	Ulcerative (chronic) rectosigmoiditis with other complication
K51.319	Ulcerative (chronic) rectosigmoiditis with unspecified complications
K51.50	Left sided colitis without complications
K51.511	Left sided colitis with rectal bleeding
K51.512	Left sided colitis with intestinal obstruction
K51.513	Left sided colitis with fistula
K51.514	Left sided colitis with abscess
K51.518	Left sided colitis with other complication
K51.519	Left sided colitis with unspecified complications
K51.80	Other ulcerative colitis without complications
K51.811	Other ulcerative colitis with rectal bleeding
K51.812	Other ulcerative colitis with intestinal obstruction

ICD-10	ICD-10 Description
K51.813	Other ulcerative colitis with fistula
K51.814	Other ulcerative colitis with abscess
K51.818	Other ulcerative colitis with other complication
K51.819	Other ulcerative colitis with unspecified complications
K51.90	Ulcerative colitis, unspecified, without complications
K51.911	Ulcerative colitis, unspecified with rectal bleeding
K51.912	Ulcerative colitis, unspecified with intestinal obstruction
K51.913	Ulcerative colitis, unspecified with fistula
K51.914	Ulcerative colitis, unspecified with abscess
K51.918	Ulcerative colitis, unspecified with other complication
K51.919	Ulcerative colitis, unspecified with unspecified complications
K52.1	Toxic gastroenteritis and colitis
R19.7	Diarrhea, unspecified

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <http://www.cms.gov/medicare-coverage-database/search/advanced-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

### J3358

<b>Jurisdiction(s):</b> E	<b>NCD/LCD/LCA Document (s):</b> A52953
<a href="https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A52953&amp;bc=gAAAAAAAAAAAA">https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A52953&amp;bc=gAAAAAAAAAAAA</a>	
<b>Jurisdiction(s):</b> F	<b>NCD/LCD/LCA Document (s):</b> A52991
<a href="https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A52991&amp;bc=gAAAAAAAAAAAA">https://www.cms.gov/medicare-coverage-database/search/article-date-search.aspx?DocID=A52991&amp;bc=gAAAAAAAAAAAA</a>	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)

### Medicare Part B Administrative Contractor (MAC) Jurisdictions

Jurisdiction	Applicable State/US Territory	Contractor
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC