



Bavencio® (avelumab) (Intravenous)

Document Number: IC-0295

Last Review Date: 12/01/2022

Date of Origin: 04/25/2017

Dates Reviewed: 04/2017, 05/2017, 08/2017, 11/2017, 02/2018, 05/2018, 09/2018, 11/2018, 03/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020, 08/2020, 12/2020, 03/2021, 06/2021, 09/2021, 12/2021, 03/2022, 06/2022, 09/2022, 12/2022

I. Length of Authorization

Coverage will be provided for 6 months and may be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Bavencio 200 mg/10 mL single-dose vial: 4 vials per 14 days

B. Max Units (per dose and over time) [HCPCS Unit]:

- 80 billable units (800 mg) every 14 days (all indications)

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age, unless otherwise indicated; **AND**

Universal Criteria

- Patient has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy (e.g., nivolumab, pembrolizumab, dostarlimab, atezolizumab, durvalumab, cemiplimab, nivolumab/relatlimab-rmbw, etc.), unless otherwise specified; **AND**

Merkel Cell Carcinoma (MCC) † ‡ ◊ ^{1,2,4,5}

- Patient is at least 12 years of age; **AND**
- Used as single-agent therapy; **AND**
- Patient has metastatic or recurrent disseminated disease

Urothelial Carcinoma (Bladder Cancer) † ‡ ^{1,4,6,8,16}

- Used as single-agent therapy; **AND**
 - Patient has one of the following diagnoses:

- Locally advanced or metastatic urothelial carcinoma †
 - Muscle invasive bladder cancer with local recurrence or persistent disease in a preserved bladder
 - Metastatic or local bladder cancer recurrence post cystectomy
 - Metastatic upper genitourinary (GU) tract tumors
 - Metastatic urothelial carcinoma of the prostate
 - Recurrent or metastatic primary carcinoma of the urethra (*excluding recurrence of stage T3-4 disease or palpable inguinal lymph nodes*)
- AND**

- Used for disease that progressed during or following platinum-containing chemotherapy*; **OR**
- Used as second-line treatment after chemotherapy other than a platinum; **OR**
- Used as first-line maintenance treatment; **AND**
 - Patient has locally advanced or metastatic urothelial carcinoma (inclusive of bladder, upper GU tract, urethra, and/or prostate cancer); **AND**
 - Patient has not progressed with first-line platinum-containing chemotherapy

* **Note:** 6,7,17,20

- *If patient was progression free for > 12 months after platinum therapy, consider re-treatment with platinum-based therapy if the patient is still platinum eligible (see below for cisplatin- or platinum-ineligible comorbidities).*
 - *Cisplatin-ineligible comorbidities may include the following: CrCl < 60 mL/min, PS ≥ 2, hearing loss of ≥ 25 decibels (dB) at two contiguous frequencies, grade ≥ 2 peripheral neuropathy, or NYHA class ≥ 3. Carboplatin may be substituted for cisplatin particularly in those patients with a CrCl < 60 mL/min or a PS of 2.*
 - *Platinum-ineligible comorbidities may include the following: CrCl < 30 mL/min, PS ≥ 3, grade ≥ 2 peripheral neuropathy, or NYHA class > 3, etc.*

Renal Cell Carcinoma (RCC) † ‡ 1,4,9,14

- Used in combination with axitinib; **AND**
- Used as first-line therapy; **AND**
- Used for the treatment of advanced, relapsed, or stage IV disease and clear cell histology

Gestational Trophoblastic Neoplasia ‡ 4,13,15

- Used as single-agent therapy for multiagent chemotherapy-resistant disease; **AND**
 - Patient has intermediate placental site trophoblastic tumor (PSTT) or epithelioid trophoblastic tumor (ETT); **AND**
 - Patient has recurrent or progressive disease; **AND**
 - Patient was previously treated with a platinum-based regimen; **OR**
 - Patient has high-risk disease (i.e., prognostic score ≥ 7 or FIGO stage IV disease)

Endometrial Carcinoma (Uterine Neoplasms) ‡ 4,18

- Used as single-agent therapy; **AND**
- Patient has recurrent or metastatic disease; **AND**
- Used as second-line treatment for microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) tumors

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓞ Orphan Drug

IV. Renewal Criteria ^{1,2}

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe or life-threatening infusion-related reactions, hepatotoxicity, severe and fatal immune-mediated adverse reactions (e.g., pneumonitis, hepatitis, colitis, endocrinopathies, nephritis with renal dysfunction, dermatitis/dermatologic adverse reactions, etc.), major adverse cardiovascular events (MACE) when used in combination with axitinib, complications of allogeneic hematopoietic stem cell transplantation (HSCT), etc.

V. Dosage/Administration ^{1,13,18}

Indication	Dose
All Indications	Administer 800 mg intravenously every 14 days, until disease progression or unacceptable toxicity
<p><u>Dosing should be calculated using actual body weight and not flat dosing (as applicable) based on the following:</u></p> <p><u>Weight > 60 kg:</u></p> <ul style="list-style-type: none"> • Standard dose 800 mg IV every 2 weeks <p><u>Weight is ≤ 60kg:</u></p> <ul style="list-style-type: none"> • Use 600 mg IV every 2 weeks <p><i>Note: This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide. Patient-specific variables should be taken into account.</i></p>	

VI. Billing Code/Availability Information

HCPCS Code:

- J9023 – Injection, avelumab, 10 mg; 1 billable unit = 10 mg

NDC:

- Bavencio 200 mg/10 mL single-dose vial: 44087-3535-xx

VII. References

1. Bavencio [package insert]. Rockland, MA; EMD Serono, Inc; July 2022. Accessed November 2022.
2. Kaufman HL, Russell J, Hamid O, et al. Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial. *Lancet Oncol.* 2016 Oct;17(10):1374-1385.
3. Novakovic AM, Wilkins JJ, Dai H, et al. Changing body weight-based dosing to a flat dose for avelumab in metastatic Merkel cell and advanced urothelial carcinoma. *Clin Pharmacol Ther.* 2019 Sep 25.
4. Referenced with permission from the NCCN Drugs and Biologics Compendium (NCCN Compendium®) avelumab. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2022.
5. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Merkel Cell Carcinoma. Version 2.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2022.
6. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Bladder Cancer. Version 2.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2022.
7. Gupta S, Sonpavde G, Grivas P, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol.* 2019 Mar 1;37(7_suppl):451.
8. Patel MR, Ellerton J, Infante JR, et al. Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-label, phase 1 trial. *Lancet Oncol.* 2018 Jan;19(1):51-64. Doi: 10.1016/S1470-2045(17)30900-2. Epub 2017 Dec 5.
9. Motzer RJ, Penkov K, Haanen J, et al. Avelumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma. *N Engl J Med.* 2019 Mar 21;380(12):1103-1115. Doi: 10.1056/NEJMoa1816047. Epub 2019 Feb 16.

10. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. *J Oncol Pract.* 2018 Mar;14(3):e130-e136.
11. Hematology/Oncology Pharmacy Association (2019). Intravenous Cancer Drug Waste Issue Brief. Retrieved from http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug_Waste_2019.pdf
12. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. *BMJ.* 2016 Feb 29;352:i788.
13. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Gestational Trophoblastic Neoplasia. Version 1.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2022.
14. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Kidney Cancer. Version 3.2023. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2022.
15. You B, Bolze PA, Lotz JP, et al. Avelumab in Patients With Gestational Trophoblastic Tumors With Resistance to Single-Agent Chemotherapy: Cohort A of the TROPHIMMUN Phase II Trial. *J Clin Oncol.* 2020 Sep 20;38(27):3129-3137. Doi: 10.1200/JCO.20.00803. Epub 2020 Jul 27.
16. Powles T, Park SH, Voog E, et al. Avelumab Maintenance Therapy for Advanced or Metastatic Urothelial Carcinoma. *N Engl J Med.* 2020 Sep 24;383(13):1218-1230. Doi: 10.1056/NEJMoa2002788. Epub 2020 Sep 18. PMID: 32945632.
17. Bellmunt, J. (2022). Treatment of metastatic urothelial cancer of the bladder and urinary tract. In Lerner SP, Shah S (Eds.), *UptoDate*. Accessed November 1, 2022. Available from https://www.uptodate.com/contents/treatment-of-metastatic-urothelial-cancer-of-the-bladder-and-urinary-tract?search=cisplatin%20ineligible&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1.
18. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Uterine Neoplasms. Version 1.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network,

Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed November 2022.

19. Galsky MD, Balar AV, Black PC, et al. Society for Immunotherapy of Cancer (SITC) clinical practice guideline on immunotherapy for the treatment of urothelial cancer. *Journal for Immunotherapy of Cancer* 2021;9:e002552. doi: 10.1136/jitc-2021-002552.
20. Gupta S, Bellmunt J, Plimack ER, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). *J Clin Oncol.* 2022 June 1;40(16_suppl):4577.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C4A.0	Merkel cell carcinoma of lip
C4A.10	Merkel cell carcinoma of eyelid, including canthus
C4A.111	Merkel cell carcinoma of right upper eyelid, including canthus
C4A.112	Merkel cell carcinoma of right lower eyelid, including canthus
C4A.121	Merkel cell carcinoma of left upper eyelid, including canthus
C4A.122	Merkel cell carcinoma of left lower eyelid, including canthus
C4A.20	Merkel cell carcinoma of unspecified ear and external auricular canal
C4A.21	Merkel cell carcinoma of right ear and external auricular canal
C4A.22	Merkel cell carcinoma of left ear and external auricular canal
C4A.30	Merkel cell carcinoma of unspecified part of face
C4A.31	Merkel cell carcinoma of nose
C4A.39	Merkel cell carcinoma of other parts of face
C4A.4	Merkel cell carcinoma of scalp and neck
C4A.51	Merkel cell carcinoma of anal skin
C4A.52	Merkel cell carcinoma of skin of breast
C4A.59	Merkel cell carcinoma of other part of trunk
C4A.60	Merkel cell carcinoma of unspecified upper limb, including shoulder
C4A.61	Merkel cell carcinoma of right upper limb, including shoulder
C4A.62	Merkel cell carcinoma of left upper limb, including shoulder
C4A.70	Merkel cell carcinoma of unspecified lower limb, including hip
C4A.71	Merkel cell carcinoma of right lower limb, including hip
C4A.72	Merkel cell carcinoma of left lower limb, including hip
C4A.8	Merkel cell carcinoma of overlapping sites
C4A.9	Merkel cell carcinoma, unspecified
C54.0	Malignant neoplasm of isthmus uteri
C54.1	Malignant neoplasm of endometrium
C54.2	Malignant neoplasm of myometrium
C54.3	Malignant neoplasm of fundus uteri
C54.8	Malignant neoplasm of overlapping sites of corpus uteri

ICD-10	ICD-10 Description
C54.9	Malignant neoplasm of corpus uteri, unspecified
C55	Malignant neoplasm of uterus, part unspecified
C58	Malignant neoplasm of placenta
C61	Malignant neoplasm of prostate
C64.1	Malignant neoplasm of right kidney, except renal pelvis
C64.2	Malignant neoplasm of left kidney, except renal pelvis
C64.9	Malignant neoplasm of unspecified kidney, except renal pelvis
C65.1	Malignant neoplasm of right renal pelvis
C65.2	Malignant neoplasm of left renal pelvis
C65.9	Malignant neoplasm of unspecified renal pelvis
C66.1	Malignant neoplasm of right ureter
C66.2	Malignant neoplasm of left ureter
C66.9	Malignant neoplasm of unspecified ureter
C67.0	Malignant neoplasm of trigone of bladder
C67.1	Malignant neoplasm of dome of bladder
C67.2	Malignant neoplasm of lateral wall of bladder
C67.3	Malignant neoplasm of anterior wall of bladder
C67.4	Malignant neoplasm of posterior wall of bladder
C67.5	Malignant neoplasm of bladder neck
C67.6	Malignant neoplasm of ureteric orifice
C67.7	Malignant neoplasm of urachus
C67.8	Malignant neoplasm of overlapping sites of bladder
C67.9	Malignant neoplasm of bladder, unspecified
C68.0	Malignant neoplasm of urethra
C7B.1	Secondary Merkel cell carcinoma
D09.0	Carcinoma in situ of bladder
D39.2	Neoplasm of uncertain behavior of placenta
O01.9	Hydatidiform mole, unspecified
Z85.51	Personal history of malignant neoplasm of bladder
Z85.59	Personal history of malignant neoplasm of other urinary tract organ
Z85.821	Personal history of Merkel cell carcinoma

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at:

<https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC