



Yescarta® (axicabtagene ciloleucel) (Intravenous)

Document Number: IC-0333

Last Review Date: 05/02/2022

Date of Origin: 10/31/2017

Dates Reviewed: 10/2017, 11/2018, 10/2019, 12/2019, 11/2020, 04/2021, 11/2021, 05/2022

I. Length of Authorization

Coverage will be provided for one treatment course (1 dose of Yescarta) and may not be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- 1 infusion bag of up to 200 million autologous anti-cd19 CAR-positive viable T-cells

B. Max Units (per dose and over time) [HCPCS Unit]:

- 1 billable unit (1 infusion of up to 200 million autologous anti-cd19 CAR-positive viable T-cells)

III. Initial Approval Criteria ¹

- Submission of medical records related to the medical necessity criteria is **REQUIRED** on all requests for authorizations. Records will be reviewed at the time of submission. Please provide documentation via direct upload through the PA web portal or by fax.

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**
- Patient does not have a clinically significant active systemic infection or inflammatory disorder; **AND**
- Patient has not received live vaccines within 6 weeks prior to the start of lymphodepleting chemotherapy, and will not receive live vaccines during axicabtagene ciloleucel treatment and until immune recovery following treatment; **AND**
- Patient has been screened for hepatitis B virus (HBV), hepatitis C virus (HCV), and human immunodeficiency virus (HIV) in accordance with clinical guidelines prior to collection of cells (leukapheresis); **AND**
- Prophylaxis for infection will be followed according to local guidelines; **AND**

- Healthcare facility has enrolled in the YESCARTA & TECARTUS REMS Program and training has been given to providers on the management of cytokine release syndrome (CRS) and neurological toxicities; **AND**
- Patient has not received prior CAR-T therapy; **AND**
- Patient has not received prior anti-CD19 therapy, (e.g., blinatumomab, etc.) OR patient previously received anti-CD19 therapy and re-biopsy indicates CD-19 positive disease; **AND**
- Used as single agent therapy (not applicable to lymphodepleting or additional chemotherapy while awaiting manufacture); **AND**
- Patient has not received prior allogeneic hematopoietic stem cell transplantation (HSCT); **AND**
- Patient has an ECOG performance status of 0-1; **AND**
- Patient does not have primary central nervous system lymphoma; **AND**

B-Cell Lymphomas † ‡ ◻ 1,7,12,13

- Patient has histologic transformation of follicular lymphoma to DLBCL; **AND**
 - Disease is refractory to first-line chemoimmunotherapy; **OR**
 - Used for treatment of disease that has relapsed within 12 months of first-line chemoimmunotherapy; **OR**
 - Patient has received two (2) or more prior lines of chemoimmunotherapy which must have included an anthracycline or anthracenedione-based regimen, unless contraindicated; **OR**
- Patient has histologic transformation of nodal marginal zone lymphoma to DLBCL OR Richter's transformation of CLL to DLBCL; **AND**
 - Patient has received two (2) or more prior lines of chemoimmunotherapy which must have included an anthracycline or anthracenedione-based regimen, unless contraindicated; **OR**
- Patient has AIDS-related large B-cell lymphoma (e.g., diffuse large B-cell lymphoma, primary effusion lymphoma, and HHV8-positive diffuse large B-cell lymphoma, not otherwise specified), DLBCL, primary mediastinal large B-cell lymphoma (PMBCL), high grade B-cell lymphoma, or monomorphic post-transplant lymphoproliferative disorder (B-cell type); **AND**
 - Used as additional therapy for disease relapse >12 months after completion of first-line therapy in patients with intention to proceed to transplant who have partial response following second-line therapy (*Note: Intention to proceed to transplant does NOT apply to monomorphic post-transplant lymphoproliferative disorder*); **OR**
 - Used as additional therapy for patients with primary refractory disease (partial response, no response, or progression) or disease relapse <12 months after completion of first-line therapy; **OR**

- Used for treatment of disease that is in second or greater relapse in patients with partial response, no response, or progressive disease following therapy for relapsed or refractory disease; **OR**
- Patient has nodal marginal zone lymphoma (MZL), nongastric MALT lymphoma (noncutaneous), gastric MALT lymphoma, grade 1-2 follicular lymphoma; **AND**
 - Disease is relapsed, refractory, or progressive after two (2) or more prior chemoimmunotherapy regimens; **OR**
- Patient has splenic marginal zone lymphoma (MZL); **AND**
 - Used for recurrent disease after two (2) or more prior chemoimmunotherapy regimens

† FDA Approved Indication(s); ‡ Compendium Recommended Indication(s); Ⓞ Orphan Drug

IV. Renewal Criteria ¹

Coverage cannot be renewed.

V. Dosage/Administration ¹

| Indication | Dose |
|---|---|
| B-Cell Lymphomas | <p><u>Lymphodepleting chemotherapy:</u></p> <ul style="list-style-type: none"> • Administer cyclophosphamide 500 mg/m² and fludarabine 30 mg/m² intravenously on the fifth, fourth, and third day before infusion of Yescarta. <p><u>Yescarta infusion:</u></p> <ul style="list-style-type: none"> • Each single infusion bag of Yescarta contains a suspension of chimeric antigen receptor (CAR)-positive T cells in approximately 68 mL. The target dose is 2 × 10⁶ CAR-positive viable T cells per kg body weight, with a maximum of 2 × 10⁸ CAR-positive viable T cells. |
| <p>For autologous use only. For intravenous use only.</p> <ul style="list-style-type: none"> • Yescarta is prepared from the patient's peripheral blood mononuclear cells, which are obtained via a standard leukapheresis procedure. • One treatment course consists of lymphodepleting chemotherapy followed by a single infusion of Yescarta. • Confirm Yescarta availability prior to starting the lymphodepleting regimen. • Confirm the patient's identity matches the patient identifiers on the YESCARTA product bag. | |
| <p><u>Premedication:</u></p> <ul style="list-style-type: none"> • Premedicate with 650 mg acetaminophen and 12.5 mg diphenhydramine 1 hour prior to infusion. Only consider the use of prophylactic systemic corticosteroids in patients after weighing the potential benefits and risks. <p><u>Monitoring after infusion:</u></p> <ul style="list-style-type: none"> • Monitor patients at least daily for 7 days at the certified healthcare facility following infusion for signs and symptoms of CRS and neurologic toxicities. • Instruct patients to remain within proximity of the certified healthcare facility for at least 4 weeks following infusion. • Instruct patients to refrain from driving or hazardous activities for at least 8 weeks following infusion. | |

- Store infusion bag in the vapor phase of liquid nitrogen (less than or equal to minus 150°C). Thaw prior to infusion.
- In case of manufacturing failure, a second manufacturing may be attempted.
- Additional chemotherapy (not the lymphodepletion) may be necessary while the patient awaits the product.
- Ensure that 2 doses of tocilizumab and emergency equipment are available prior to infusion and during the recovery period.
- Yescarta contains human blood cells that are genetically modified with replication incompetent retroviral vector. Follow universal precautions and local biosafety guidelines for handling and disposal.

VI. Billing Code/Availability Information

HCPCS Code:

- Q2041 - Axicabtagene ciloleucel, up to 200 million autologous anti-cd19 car-positive viable t cells, including leukapheresis and dose preparation procedures, per therapeutic dose

NDC:

- Yescarta suspension for intravenous infusion; 1 infusion bag (68 mL): 71287-0119-xx

VII. References

1. Yescarta [package insert]. Santa Monica, CA; Kite Pharma, Inc., April 2022. Accessed April 2022.
2. Locke FL, Neelapu SS, Bartlett NL, et al. Phase 1 Results of ZUMA-1: A Multicenter Study of KTE-C19 Anti-CD19 CAR T Cell Therapy in Refractory Aggressive Lymphoma. *Mol Ther*. 2017 Jan 4;25(1):285-295.
3. Mejstrikova E, Hrusak O, Borowitz MJ, et al. CD19-negative relapse of pediatric B-cell precursor acute lymphoblastic leukemia following blinatumomab treatment. *Blood Cancer J*. 2017; 659. DOI 10.1038/s41408-017-0023-x
4. Ruella M, Maus MV. Catch me if you can: Leukemia Escape after CD19-Directed T Cell Immunotherapies. *Computational and Structural Biotechnology Journal* 14 (2016) 357–362.
5. Braig F, Brandt A, Goebeler M, et al. Resistance to anti-CD19/CD3 BiTE in acute lymphoblastic leukemia may be mediated by disrupted CD19 membrane trafficking. *Blood*; 129:1, 2017 Jan.
6. Majzner RG, Mackall CL. Tumor Antigen Escape from CAR T-cell Therapy. *Cancer Discov* 2018;8:1219-1226.
7. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) axicabtagene ciloleucel. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed April 2022.
8. Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol*. 2019;20(1):31-42. doi:10.1016/S1470-2045(18)30864-7.

9. Neelapu SS, Jacobson CA, Oluwole OO, et al. Outcomes of older patients in ZUMA-1, a pivotal study of axicabtagene ciloleucel in refractory large B-cell lymphoma. *Blood*. 2020;135(23):2106-2109. doi:10.1182/blood.2019004162.
10. Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med*. 2017;377(26):2531-2544. doi: 10.1056/NEJMoa1707447.
11. Jacobson CA, Chavez JC, Sehgal AR, et al. Interim analysis of ZUMA-5: A phase II study of axicabtagene ciloleucel (axi-cel) in patients (pts) with relapsed/refractory indolent non-Hodgkin lymphoma (R/R iNHL). *Journal of Clinical Oncology* 2020 38:15_suppl, 8008-8008.
12. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2022.
13. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for B-Cell Lymphomas Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed April 2022.
14. Locke FL, Miklos DB, Jacobson CA, et al; All ZUMA-7 Investigators and Contributing Kite Members. Axicabtagene Ciloleucel as Second-Line Therapy for Large B-Cell Lymphoma. *N Engl J Med*. 2022 Feb 17;386(7):640-654. doi: 10.1056/NEJMoa2116133.

Appendix 1 – Covered Diagnosis Codes

| ICD-10 | ICD-10 Description |
|--------|--|
| C82.00 | Follicular lymphoma grade I, unspecified site |
| C82.01 | Follicular lymphoma grade I, lymph nodes of head, face and neck |
| C82.02 | Follicular lymphoma, grade I, intrathoracic lymph nodes |
| C82.03 | Follicular lymphoma grade I, intra-abdominal lymph nodes |
| C82.04 | Follicular lymphoma grade I, lymph nodes of axilla and upper limb |
| C82.05 | Follicular lymphoma grade I, lymph nodes of inguinal regional and lower limb |
| C82.06 | Follicular lymphoma grade I, intrapelvic lymph nodes |
| C82.07 | Follicular lymphoma grade I, spleen |
| C82.08 | Follicular lymphoma grade I, lymph nodes of multiple sites |
| C82.09 | Follicular lymphoma grade I, extranodal and solid organ sites |
| C82.10 | Follicular lymphoma grade II, unspecified site |
| C82.11 | Follicular lymphoma grade II, lymph nodes of head, face and neck |

| | |
|--------|---|
| C82.12 | Follicular lymphoma, grade II, intrathoracic lymph nodes |
| C82.13 | Follicular lymphoma grade II, intra-abdominal lymph nodes |
| C82.14 | Follicular lymphoma grade II, lymph nodes of axilla and upper limb |
| C82.15 | Follicular lymphoma grade II, lymph nodes of inguinal region and lower limb |
| C82.16 | Follicular lymphoma grade II, intrapelvic lymph nodes |
| C82.17 | Follicular lymphoma grade II, spleen |
| C82.18 | Follicular lymphoma grade II, lymph nodes of multiple sites |
| C82.19 | Follicular lymphoma grade II, extranodal and solid organ sites |
| C82.20 | Follicular lymphoma grade III, unspecified, unspecified site |
| C82.21 | Follicular lymphoma grade III, unspecified, lymph nodes of head, face and neck |
| C82.22 | Follicular lymphoma, grade III, unspecified, intrathoracic lymph nodes |
| C82.23 | Follicular lymphoma grade III, unspecified, intra-abdominal lymph nodes |
| C82.24 | Follicular lymphoma grade III, unspecified, lymph nodes of axilla and upper limb |
| C82.25 | Follicular lymphoma grade III, unspecified, lymph nodes of inguinal region and lower limb |
| C82.26 | Follicular lymphoma grade III, unspecified, intrapelvic lymph nodes |
| C82.27 | Follicular lymphoma grade III, unspecified, spleen |
| C82.28 | Follicular lymphoma grade III, unspecified, lymph nodes of multiple sites |
| C82.29 | Follicular lymphoma grade III, unspecified, extranodal and solid organ sites |
| C82.30 | Follicular lymphoma grade IIIa, unspecified site |
| C82.31 | Follicular lymphoma grade IIIa, lymph nodes of head, face and neck |
| C82.32 | Follicular lymphoma, grade IIIa, intrathoracic lymph nodes |
| C82.33 | Follicular lymphoma grade IIIa, intra-abdominal lymph nodes |
| C82.34 | Follicular lymphoma grade IIIa, lymph nodes of axilla and upper limb |
| C82.35 | Follicular lymphoma grade IIIa, lymph nodes of inguinal region and lower limb |
| C82.36 | Follicular lymphoma grade IIIa, intrapelvic lymph nodes |
| C82.37 | Follicular lymphoma grade IIIa, spleen |
| C82.38 | Follicular lymphoma grade IIIa, lymph nodes of multiple sites |
| C82.39 | Follicular lymphoma grade IIIa, extranodal and solid organ sites |
| C82.40 | Follicular lymphoma grade IIIb, unspecified site |
| C82.41 | Follicular lymphoma grade IIIb, lymph nodes of head, face, and neck |
| C82.42 | Follicular lymphoma grade IIIb, intrathoracic lymph nodes |
| C82.43 | Follicular lymphoma grade IIIb, intra-abdominal lymph nodes |
| C82.44 | Follicular lymphoma grade IIIb, lymph nodes of axilla and upper limb |
| C82.45 | Follicular lymphoma grade IIIb, lymph nodes of inguinal region and lower limb |
| C82.46 | Follicular lymphoma grade IIIb, intrapelvic lymph nodes |
| C82.47 | Follicular lymphoma grade IIIb, spleen |

| | |
|--------|---|
| C82.48 | Follicular lymphoma grade IIIb, lymph nodes of multiple sites |
| C82.49 | Follicular lymphoma grade IIIb, extranodal and solid organ sites |
| C82.50 | Diffuse follicle center lymphoma, unspecified site |
| C82.51 | Diffuse follicle center lymphoma, lymph nodes of head, face and neck |
| C82.52 | Diffuse follicle center lymphoma, intrathoracic lymph nodes |
| C82.53 | Diffuse follicle center lymphoma, intra-abdominal lymph nodes |
| C82.54 | Diffuse follicle center lymphoma, lymph nodes of axilla and upper limb |
| C82.55 | Diffuse follicle center lymphoma, lymph nodes of inguinal region and lower limb |
| C82.56 | Diffuse follicle center lymphoma, intrapelvic lymph nodes |
| C82.57 | Diffuse follicle center lymphoma, spleen |
| C82.58 | Diffuse follicle center lymphoma, lymph nodes of multiple sites |
| C82.59 | Diffuse follicle center lymphoma, extranodal and solid organ sites |
| C82.60 | Cutaneous follicle center lymphoma, unspecified site |
| C82.61 | Cutaneous follicle center lymphoma, lymph nodes of head, face and neck |
| C82.62 | Cutaneous follicle center lymphoma, intrathoracic lymph nodes |
| C82.63 | Cutaneous follicle center lymphoma, intra-abdominal lymph nodes |
| C82.64 | Cutaneous follicle center lymphoma, lymph nodes of axilla and upper limb |
| C82.65 | Cutaneous follicle center lymphoma, lymph nodes of inguinal region and lower limb |
| C82.66 | Cutaneous follicle center lymphoma, intrapelvic lymph nodes |
| C82.67 | Cutaneous follicle center lymphoma, spleen |
| C82.68 | Cutaneous follicle center lymphoma, lymph nodes of multiple sites |
| C82.69 | Cutaneous follicle center lymphoma, extranodal and solid organ sites |
| C82.80 | Other types of follicular lymphoma, unspecified site |
| C82.81 | Other types of follicular lymphoma, lymph nodes of head, face and neck |
| C82.82 | Other types of follicular lymphoma, intrathoracic lymph nodes |
| C82.83 | Other types of follicular lymphoma, intra-abdominal lymph nodes |
| C82.84 | Other types of follicular lymphoma, lymph nodes of axilla and upper limb |
| C82.85 | Other types of follicular lymphoma, lymph nodes of inguinal region and lower limb |
| C82.86 | Other types of follicular lymphoma, intrapelvic lymph nodes |
| C82.87 | Other types of follicular lymphoma, spleen |
| C82.88 | Other types of follicular lymphoma, lymph nodes of multiple sites |
| C82.89 | Other types of follicular lymphoma, extranodal and solid organ sites |
| C82.90 | Follicular lymphoma, unspecified, unspecified site |
| C82.91 | Follicular lymphoma, unspecified, lymph nodes of head, face and neck |
| C82.92 | Follicular lymphoma, unspecified, intrathoracic lymph nodes |
| C82.93 | Follicular lymphoma, unspecified, intra-abdominal lymph nodes |

| | |
|--------|--|
| C82.94 | Follicular lymphoma, unspecified, lymph nodes of axilla and upper limb |
| C82.95 | Follicular lymphoma, unspecified lymph nodes of inguinal region and lower limb |
| C82.96 | Follicular lymphoma, unspecified, intrapelvic lymph nodes |
| C82.97 | Follicular lymphoma, unspecified, spleen |
| C82.98 | Follicular lymphoma, unspecified, lymph nodes of multiple sites |
| C82.99 | Follicular lymphoma, unspecified, extranodal and solid organ sites |
| C83.00 | Small cell B-cell lymphoma, unspecified site |
| C83.01 | Small cell B-cell lymphoma, lymph nodes of head, face and neck |
| C83.02 | Small cell B-cell lymphoma, intrathoracic lymph nodes |
| C83.03 | small cell B-cell lymphoma, intra-abdominal lymph nodes |
| C83.04 | Small cell B-cell lymphoma, lymph nodes of axilla and upper limb |
| C83.05 | Small cell B-cell lymphoma, lymph nodes of inguinal region and lower limb |
| C83.06 | Small cell B-cell lymphoma, intrapelvic lymph nodes |
| C83.07 | Small cell B-cell lymphoma, spleen |
| C83.08 | Small cell B-cell lymphoma, lymph nodes of multiple sites |
| C83.09 | Small cell B-cell lymphoma, extranodal and solid organ sites |
| C83.30 | Diffuse large B-cell lymphoma unspecified site |
| C83.31 | Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck |
| C83.32 | Diffuse large B-cell lymphoma intrathoracic lymph nodes |
| C83.33 | Diffuse large B-cell lymphoma intra-abdominal lymph nodes |
| C83.34 | Diffuse large B-cell lymphoma lymph nodes of axilla and upper limb |
| C83.35 | Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb |
| C83.36 | Diffuse large B-cell lymphoma intrapelvic lymph nodes |
| C83.37 | Diffuse large B-cell lymphoma, spleen |
| C83.38 | Diffuse large B-cell lymphoma lymph nodes of multiple sites |
| C83.39 | Diffuse large B-cell lymphoma extranodal and solid organ sites |
| C83.80 | Other non-follicular lymphoma, unspecified site |
| C83.81 | Other non-follicular lymphoma, lymph nodes of head, face and neck |
| C83.82 | Other non-follicular lymphoma, intrathoracic lymph nodes |
| C83.83 | Other non-follicular lymphoma, intra-abdominal lymph nodes |
| C83.84 | Other non-follicular lymphoma, lymph nodes of axilla and upper limb |
| C83.85 | Other non-follicular lymphoma, lymph nodes of inguinal region and lower limb |
| C83.86 | Other non-follicular lymphoma, intrapelvic lymph nodes |
| C83.87 | Other non-follicular lymphoma, spleen |
| C83.88 | Other non-follicular lymphoma, lymph nodes of multiple sites |
| C83.89 | Other non-follicular lymphoma, extranodal and solid organ sites |

| | |
|--------|--|
| C83.90 | Non-follicular (diffuse) lymphoma, unspecified site |
| C83.91 | Non-follicular (diffuse) lymphoma, unspecified lymph nodes of head, face, and neck |
| C83.92 | Non-follicular (diffuse) lymphoma, unspecified intrathoracic lymph nodes |
| C83.93 | Non-follicular (diffuse) lymphoma, unspecified intra-abdominal lymph nodes |
| C83.94 | Non-follicular (diffuse) lymphoma, unspecified lymph nodes of axilla and upper limb |
| C83.95 | Non-follicular (diffuse) lymphoma, unspecified lymph nodes of inguinal region and lower limb |
| C83.96 | Non-follicular (diffuse) lymphoma, unspecified intrapelvic lymph nodes |
| C83.97 | Non-follicular (diffuse) lymphoma, unspecified spleen |
| C83.98 | Non-follicular (diffuse) lymphoma, unspecified lymph nodes of multiple sites |
| C83.99 | Non-follicular (diffuse) lymphoma, unspecified extranodal and solid organ sites |
| C85.10 | Unspecified B-cell lymphoma, unspecified site |
| C85.11 | Unspecified B-cell lymphoma, lymph nodes of head, face, and neck |
| C85.12 | Unspecified B-cell lymphoma, intrathoracic lymph nodes |
| C85.13 | Unspecified B-cell lymphoma, intra-abdominal lymph nodes |
| C85.14 | Unspecified B-cell lymphoma, lymph nodes of axilla and upper limb |
| C85.15 | Unspecified B-cell lymphoma, lymph nodes of inguinal region and lower limb |
| C85.16 | Unspecified B-cell lymphoma, intrapelvic lymph nodes |
| C85.17 | Unspecified B-cell lymphoma, spleen |
| C85.18 | Unspecified B-cell lymphoma, lymph nodes of multiple sites |
| C85.19 | Unspecified B-cell lymphoma, extranodal and solid organ sites |
| C85.20 | Mediastinal (thymic) large B-cell lymphoma, unspecified site |
| C85.21 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck |
| C85.22 | Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes |
| C85.23 | Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes |
| C85.24 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb |
| C85.25 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb |
| C85.26 | Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes |
| C85.27 | Mediastinal (thymic) large B-cell lymphoma, spleen |
| C85.28 | Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites |
| C85.29 | Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites |
| C85.80 | Other specified types of non-Hodgkin lymphoma, unspecified site |
| C85.81 | Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face and neck |
| C85.82 | Other specified types of non-Hodgkin lymphoma, intrathoracic lymph nodes |
| C85.83 | Other specified types of non-Hodgkin lymphoma, intra-abdominal lymph nodes |
| C85.84 | Other specified types of non-Hodgkin lymphoma, lymph nodes of axilla and upper limb |
| C85.85 | Other specified types of non-Hodgkin lymphoma, lymph nodes of inguinal region of lower limb |

| | |
|--------|---|
| C85.86 | Other specified types of non-Hodgkin lymphoma, intrapelvic lymph nodes |
| C85.87 | Other specified types of non-Hodgkin lymphoma, spleen |
| C85.88 | Other specified types of non-Hodgkin lymphoma, lymph nodes of multiple sites |
| C85.89 | Other specified types of non-Hodgkin lymphoma, extranodal and solid organ sites |
| C88.4 | Extranodal marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue [MALT-lymphoma] |
| C91.10 | Chronic lymphocytic leukemia of B-cell type not having achieved remission |
| C91.12 | Chronic lymphocytic leukemia of B-cell type in relapse |
| D47.Z1 | Post-transplant lymphoproliferative disorder (PTLD) |

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Article and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

| Medicare Part B Administrative Contractor (MAC) Jurisdictions | | |
|---|---|---|
| Jurisdiction | Applicable State/US Territory | Contractor |
| E (1) | CA, HI, NV, AS, GU, CNMI | Noridian Healthcare Solutions, LLC |
| F (2 & 3) | AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ | Noridian Healthcare Solutions, LLC |
| 5 | KS, NE, IA, MO | Wisconsin Physicians Service Insurance Corp (WPS) |
| 6 | MN, WI, IL | National Government Services, Inc. (NGS) |
| H (4 & 7) | LA, AR, MS, TX, OK, CO, NM | Novitas Solutions, Inc. |
| 8 | MI, IN | Wisconsin Physicians Service Insurance Corp (WPS) |
| N (9) | FL, PR, VI | First Coast Service Options, Inc. |
| J (10) | TN, GA, AL | Palmetto GBA, LLC |
| M (11) | NC, SC, WV, VA (excluding below) | Palmetto GBA, LLC |
| L (12) | DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA) | Novitas Solutions, Inc. |
| K (13 & 14) | NY, CT, MA, RI, VT, ME, NH | National Government Services, Inc. (NGS) |
| 15 | KY, OH | CGS Administrators, LLC |