

Enhertu® (fam-trastuzumab deruxtecan-nxki)

(Intravenous)

Document Number: IC-0522

Last Review Date: 06/06/2022 Date of Origin: 02/04/2020 Dates Reviewed: 02/2020, 06/2020, 09/2020, 02/2021, 03/2022, 06/2022

I. Length of Authorization

Coverage will be provided for 6 months and may be renewed.

II. Dosing Limits

- A. Quantity Limit (max daily dose) [NDC Unit]:
 - Enhertu 100 mg vial: 7 vials every 21 days

B. Max Units (per dose and over time) [HCPCS Unit]:

- Breast Cancer: 600 billable units every 21 days
- All other indications: 700 billable units every 21 days

III. Initial Approval Criteria 1

Coverage is provided in the following conditions:

• Patient is at least 18 years of age; AND

Universal Criteria¹

- Left ventricular ejection fraction (LVEF) is within normal limits prior to initiating therapy and will be assessed at regular intervals (e.g., every 3 months) during treatment; **AND**
- Used as a single agent; AND
- Therapy will not be substituted with or for any trastuzumab-based formulation (i.e., trastuzumab [or trastuzumab biosimilar product], ado-trastuzumab emtansine, trastuzumab-hyaluronidase, pertuzumab/trastuzumab and hyaluronidase-zzxf, etc.); **AND**

Breast Cancer **† ‡** ^{1,2,4,8,15,16}

- Patient has human epidermal growth factor receptor 2 (HER2)-positive* disease as determined by an FDA-approved or CLIA-compliant test*; **AND**
- Patient has recurrent unresectable (local or regional) or metastatic disease OR inflammatory breast cancer with no response to preoperative systemic therapy; **AND**
 - \circ $\;$ Used as subsequent therapy; OR

Proprietary & Confidential © 2022 Magellan Health, Inc.



 Used as first-line therapy in patients with who experience disease progression during or within 6 months of neoadjuvant or adjuvant therapy (12 months for pertuzumabcontaining regimens)

Gastric, Esophageal, and Esophagogastric Junction Cancers $\ddagger \ddagger \Phi^{1,2,9,17,18}$

- Patient has human epidermal growth factor receptor 2 (HER2)-positive* disease as determined by an FDA-approved or CLIA-compliant test*; **AND**
- Patient has adenocarcinoma histology; AND
- Patient is not a surgical candidate OR has unresectable locally advanced, recurrent, or metastatic disease; **AND**
- Used as subsequent therapy

Colorectal Cancer (CRC) ‡ ^{2,10-12}

- Patient has human epidermal growth factor receptor 2 (HER2)-positive* disease as determined by an FDA-approved or CLIA-compliant test*; **AND**
- Patient has RAS and BRAF wild-type (WT) disease; AND
- Used as subsequent therapy for progression of advanced or metastatic disease after at least one prior line of treatment in the advanced or metastatic disease setting

Appendiceal Adenocarcinoma – Colon Cancer ‡ 2,11

- Patient has human epidermal growth factor receptor 2 (HER2)-positive* disease as determined by an FDA-approved or CLIA-compliant test*; **AND**
- Patient has RAS and BRAF wild-type (WT) disease; **AND**
- Used as subsequent therapy for progression of advanced or metastatic disease after at least one prior line of treatment in the advanced or metastatic disease setting

Non-Small Cell Lung Cancer (NSCLC) ‡ 2,13,14

- Patient has ERBB2 (HER2) mutation positive disease as determined by an FDA-approved or CLIA-complaint test*; **AND**
- Patient has metastatic disease

*HER2-positive overexpression criteria

Breast Cancer: 4,5

- Immunohistochemistry (IHC) assay 3+; **OR**
- Dual-probe in situ hybridization (ISH) assay HER2/CEP17 ratio \geq 2.0 AND average HER2 copy number \geq 4.0 signals/cell; **OR**
- Dual-probe in situ hybridization (ISH) assay AND concurrent IHC indicating one of the following:
 - $\circ~$ HER2/CEP17 ratio ≥ 2.0 AND average HER2 copy number < 4.0 signals/cell AND concurrent IHC 3+; \mathbf{OR}



- $\circ~$ HER2/CEP17 ratio < 2.0 AND average HER2 copy number \geq 6.0 signals/cell AND concurrent IHC 2+ or 3+; \mathbf{OR}
- $\circ~$ HER2/CEP17 ratio < 2.0 AND average HER2 copy number \geq 4.0 and < 6.0 signals/cell AND concurrent IHC 3+

Gastric, Esophageal, and Esophagogastric Junction Cancer: 17-19

- Immunohistochemistry (IHC) assay 3+; OR
- Fluorescence in situ hybridization (FISH) or in situ hybridization (ISH) assay AND concurrent IHC indicating one of the following:
 - HER2/CEP17 ratio \geq 2.0 AND concurrent IHC 2+; **OR**
 - \circ Average HER2 copy number ≥ 6.0 signals/cell AND concurrent IHC 2+

Colorectal Cancer and Appendiceal Adenocarcinoma: ^{11,12}

- Immunohistochemistry (IHC) assay 3+; **OR**
- Fluorescence in situ hybridization (FISH) HER2/CEP17 ratio ≥ 2 AND concurrent IHC 2+; OR
- Next-generation sequencing (NGS) panel HER2 amplification

♦ If confirmed using an FDA approved assay - http://www.fda.gov/companiondiagnostics

† FDA approved indication(s); **‡** Compendia Recommended Indication(s); **Φ** Orphan Drug *(only applies to Gastric and Esophagogastric Junction Cancers)*

IV. Renewal Criteria 1,10,13

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: pulmonary toxicity (e.g., interstitial lung disease, pneumonitis), neutropenia/febrile neutropenia, left ventricular dysfunction/symptomatic congestive heart failure, etc.; **AND**
- Left ventricular ejection fraction (LVEF) within the previous 3 months as follows:
 - \circ LVEF is > 45% and <u>absolute</u> decrease is \leq 20% from baseline; **OR**
 - \circ LVEF is 40% to 45% and <u>absolute</u> decrease is < 10% from baseline

V. Dosage/Administration ^{1,11-13}

Indication	Dose
Breast Cancer	Administer 5.4 mg/kg given as an intravenous infusion every 3 weeks (21-day cycle) until disease progression or unacceptable toxicity
	Administer 6.4 mg/kg given as an intravenous infusion every 3 weeks (21-day cycle) until disease progression or unacceptable toxicity



Annondicoal	Administer 6.4 mg/kg given as an intravenous infusion every 3 weeks (21-day cycle) until disease progression or unacceptable toxicity
0	Administer 6.4 mg/kg given as an intravenous infusion every 3 weeks (21-day cycle) until disease progression or unacceptable toxicity

VI. Billing Code/Availability Information

HCPCS Code:

• J9358 – Injection, fam-trastuzumab deruxtecan-nxki, 1 mg: 1 billable unit = 1 mg

NDC:

• Enhertu 100 mg single-dose vial: 65597-0406-xx

VII. References

- 1. Enhertu [package insert]. Basking Ridge, NJ; Daiichi Sankyo, Inc; May 2022. Accessed May 2022.
- 2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) fam-trastuzumab deruxtecan-nxki. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2022.
- Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. J Oncol Pract. 2018 Mar;14(3):e130-e136.
- 4. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for Breast Cancer, Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK[®], NCCN[®], and NCCN GUIDELINES[®] are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
- 5. Wolff AC, Hammond EH, Allison KH, et al. Human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Focused Update. J Clin Oncol 2018;36:2105-2122.
- Hematology/Oncology Pharmacy Association (2019). Intravenous Cancer Drug Waste Issue Brief. Retrieved from <u>http://www.hoparx.org/images/hopa/advocacy/Issue-</u> Briefs/Drug_Waste_2019.pdf
- 7. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. BMJ. 2016 Feb 29;352:i788.



- Modi S, Saura C, Yamashita T, et al; DESTINY-Breast01 Investigators. Trastuzumab Deruxtecan in Previously Treated HER2-Positive Breast Cancer. N Engl J Med. 2019 Dec 11. Doi: 10.1056/NEJMoa1914510.
- Shitara K, Bang YJ, Iwasa S, et al; DESTINY-Gastric01 Investigators. Trastuzumab Deruxtecan in Previously Treated HER2-Positive Gastric Cancer. N Engl J Med. 2020 Jun 18;382(25):2419-2430.
- Siena S, Di Bartolomeo M, Raghav KPS, et al. A phase II, multicenter, open-label study of trastuzumab deruxtecan in patients with HER2-expressing metastatic colorectal cancer (mCRC): DESTINY-CRC01. J Clin Oncol 2020;38(suppl; abstr 4000).
- 11. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for Colon Cancer, Version 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
- 12. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for Rectal Cancer 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
- 13. Smit EF, Nakagawa K, Nagasaka M, et al. Trastuzumab deruxtecan (T-DXd; DS-8201) in patients with HER2-mutated metastatic non-small cell lung cancer (NSCLC): interim results of DESTINY-Lung01[abstract]. J Clin Oncol 2020;38:Abstract 9504.
- 14. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for Non-Small Cell Lung Cancer, Version 3.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
- 15. Gennari A, André F, Barrios CH, et al.; ESMO Guidelines Committee. Electronic address: clinicalguidelines@esmo.org. ESMO Clinical Practice Guideline for the diagnosis, staging and treatment of patients with metastatic breast cancer. Ann Oncol. 2021 Dec;32(12):1475-1495. doi: 10.1016/j.annonc.2021.09.019. Epub 2021 Oct 19. PMID: 34678411.
- 16. Cortés J, Kim SB, Chung WP, et al; DESTINY-Breast03 Trial Investigators. Trastuzumab Deruxtecan versus Trastuzumab Emtansine for Breast Cancer. N Engl J Med. 2022 Mar 24;386(12):1143-1154. doi: 10.1056/NEJMoa2115022.
- Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for Esophageal and Esophagogastric Junction Cancers, Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK[®], NCCN[®], and NCCN GUIDELINES[®] are trademarks owned by the

ENHERTU® (fam-trastuzumab deruxtecan-nxki)



National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.

- 18. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) for Gastric Cancer, Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
- Bartley AN, Washington MK, Colasacco C, et al. HER2 Testing and Clinical Decision Making in Gastroesophageal Adenocarcinoma: Guideline From the College of American Pathologists, American Society for Clinical Pathology, and the American Society of Clinical Oncology. J Clin Oncol. 2017 Feb;35(4):446-464. doi: 10.1200/JCO.2016.69.4836.
- 20. First Coast Service Options, Inc. Local Coverage Article: Billing and Coding: Trastuzumab -Trastuzumab Biologics (A56660). Centers for Medicare & Medicaid Services, Inc. Updated on 10/08/2021 with effective date of 10/01/2021. Accessed May 2022.

ICD-10	ICD-10 Description		
C15.3	Malignant neoplasm of upper third of esophagus		
C15.4	Malignant neoplasm of middle third of esophagus		
C15.5	Malignant neoplasm of the lower third of esophagus		
C15.8	Malignant neoplasm of overlapping sites of esophagus		
C15.9	Malignant neoplasm of esophagus, unspecified		
C16.0	Malignant neoplasm of cardia		
C16.1	Malignant neoplasm of fundus of stomach		
C16.2	Malignant neoplasm of body of stomach		
C16.3	Malignant neoplasm of pyloric antrum		
C16.4	Malignant neoplasm of pylorus		
C16.5	Malignant neoplasm of lesser curvature of stomach, unspecified		
C16.6	Malignant neoplasm of greater curvature of stomach, unspecified		
C16.8	Malignant neoplasm of overlapping sites of stomach		
C16.9	Malignant neoplasm of stomach, unspecified		
C18.0	Malignant neoplasm of cecum		
C18.1	Malignant neoplasm of appendix		
C18.2	Malignant neoplasm of ascending colon		
C18.3	Malignant neoplasm of hepatic flexure		
C18.4	Malignant neoplasm of transverse colon		
C18.5	Malignant neoplasm of splenic flexure		
Page 6	ENHERTU® (fam-trastuzumab deruxtecan-nxki) Prior Auth Criteria Proprietary Information. Restricted Access – Do not disseminate or copy without approval. ©2022 Munutum De Manuel		

Appendix 1 – Covered Diagnosis Codes

©2022, Magellan Rx Management

C18.6	Malignant neoplasm of descending colon	
C18.7	Malignant neoplasm of sigmoid colon	
C18.8	Malignant neoplasm of overlapping sites of colon	
C18.9	Malignant neoplasm of colon, unspecified	
C19	Malignant neoplasm of rectosigmoid junction	
C20	Malignant neoplasm of rectum	
C21.8	Malignant neoplasm of overlapping sites of rectum, anus and anal canal	
C33	Malignant neoplasm of trachea	
C34.00	Malignant neoplasm of unspecified main bronchus	
C34.01	Malignant neoplasm of right main bronchus	
C34.02	Malignant neoplasm of left main bronchus	
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung	
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung	
C34.12	Malignant neoplasm of upper lobe, left bronchus or lung	
C34.2	Malignant neoplasm of middle lobe, bronchus or lung	
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung	
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung	
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung	
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus and lung	
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung	
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung	
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung	
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung	
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung	
C50.011	Malignant neoplasm of nipple and areola, right female breast	
C50.012	Malignant neoplasm of nipple and areola, left female breast	
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast	
C50.021	Malignant neoplasm of nipple and areola, right male breast	
C50.022	Malignant neoplasm of nipple and areola, left male breast	
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast	
C50.111	Malignant neoplasm of central portion of right female breast	
C50.112	Malignant neoplasm of central portion of left female breast	
C50.119	Malignant neoplasm of central portion of unspecified female breast	
C50.121	Malignant neoplasm of central portion of right male breast	
C50.122	Malignant neoplasm of central portion of left male breast	



C50.129	Malignant neoplasm of central portion of unspecified male breast	
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast	
C50.212	Malignant neoplasm of upper-inner quadrant of left female breast	
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast	
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast	
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast	
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast	
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast	
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast	
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast	
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast	
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast	
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast	
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast	
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast	
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast	
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast	
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast	
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast	
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast	
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast	
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast	
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast	
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast	
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast	
C50.611	Malignant neoplasm of axillary tail of right female breast	
C50.612	Malignant neoplasm of axillary tail of left female breast	
C50.619	Malignant neoplasm of axillary tail of unspecified female breast	
C50.621	Malignant neoplasm of axillary tail of right male breast	
C50.622	Malignant neoplasm of axillary tail of left male breast	
C50.629	Malignant neoplasm of axillary tail of unspecified male breast	
C50.811	Malignant neoplasm of overlapping sites of right female breast	
C50.812	Malignant neoplasm of overlapping sites of left female breast	
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast	
C50.821	Malignant neoplasm of overlapping sites of right male breast	



C50.822	Malignant neoplasm of overlapping sites of left male breast		
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast		
C50.911	Malignant neoplasm of unspecified site of right female breast		
C50.912	Malignant neoplasm of unspecified site of left female breast		
C50.919	Malignant neoplasm of unspecified site of unspecified female breast		
C50.921	Malignant neoplasm of unspecified site of right male breast		
C50.922	Malignant neoplasm of unspecified site of left male breast		
C50.929	Malignant neoplasm of unspecified site of unspecified male breast		
C78.00	Secondary malignant neoplasm of unspecified lung		
C78.01	Secondary malignant neoplasm of right lung		
C78.02	Secondary malignant neoplasm of left lung		
C78.6	Secondary malignant neoplasm of retroperitoneum and peritoneum		
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct		
D37.1	Neoplasm of uncertain behavior of stomach		
D37.8	Neoplasm of uncertain behavior of other specified digestive organs		
D37.9	Neoplasm of uncertain behavior of digestive organ, unspecified		
Z85.00	Personal history of malignant neoplasm of unspecified digestive organ		
Z85.01	Personal history of malignant neoplasm of esophagus		
Z85.028	Personal history of other malignant neoplasm of stomach		
Z85.038	Personal history of other malignant neoplasm of large intestine		
Z85.118	Personal history of other malignant neoplasm of bronchus and lung		

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

Jurisdiction(s): N(9)	NCD/LCD/LCA Document (s): A56660	
https://www.cms.gov/medicare-coverage-database/new-search/search-		
results.aspx?keyword=a56660&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMC		
<u>D%2C6%2C3%2C5%2C1%2CF%2CP</u>		



Medicare Part B Administrative Contractor (MAC) Jurisdictions			
Jurisdiction	Applicable State/US Territory	Contractor	
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC	
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC	
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)	
6	MN, WI, IL	National Government Services, Inc. (NGS)	
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.	
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)	
N (9)	FL, PR, VI	First Coast Service Options, Inc.	
J (10)	TN, GA, AL	Palmetto GBA, LLC	
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC	
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.	
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)	
15	KY, OH	CGS Administrators, LLC	

