

Rituximab:

Rituxan[®], Truxima[®], Ruxience[®], Riabni[™] (Intravenous)

-E-

Document Number: BH-0477

Last Review Date: 07/05/2022

Date of Origin: 06/03/2019

Dates Reviewed: 06/2019, 10/2019, 11/2019, 01/2020, 04/2020, 07/2020, 10/2020, 01/2021, 04/2021, 07/2021, 10/2021, 01/2022, 04/2022, 07/2022

I. Length of Authorization ^{1-5,23-25,44,62,80,94-98,102,108,115-118,128-130}

Coverage will be provided for 6 months (12 months initially for pemphigus vulgaris) and may be renewed unless otherwise specified.

- Maintenance therapy for oncology indications (excluding ALL, Hairy Cell Leukemia, Mantle Cell Lymphoma, and induction/consolidation of Pediatric B-Cell Acute Leukemia/Aggressive Mature B-Cell Lymphomas) may be renewed for up to a maximum of 2 years.
 - ALL* and Mantle Cell Lymphoma may be renewed until disease progression or intolerable toxicity. (**ALL may be renewed for up to a total of 18 doses*)
 - Hairy Cell Leukemia may not be renewed.
 - Induction/Consolidation of Pediatric B-Cell Acute Leukemia and Aggressive Mature B-Cell Lymphomas may not be renewed.
- Management of Immunotherapy-Related Toxicities:
 - Myalgias/Myositis/Myasthenia Gravis/Encephalitis may not be renewed.
 - Bullous dermatitis may be renewed for a maximum of 18 months (4 total doses).
- Relapse therapy for Pemphigus Vulgaris must be at least 16 weeks past a prior infusion
- Chronic Graft-Versus-Host Disease (cGVHD) may not be renewed.
- Hematopoietic Cell Transplantation may not be renewed.
- Lupus Nephritis may be renewed ONLY in patients experiencing a disease relapse.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Rituxan 100 mg/10 mL injection: 12 vials per 28 day supply
- Rituxan 500 mg/50 mL injection: 8 vials per 28 day supply
- Truxima 100 mg/10 mL injection: 12 vials per 28 day supply
- Truxima 500 mg/50 mL injection: 8 vials per 28 day supply

- Ruxience 100 mg/10 mL injection: 12 vials per 28 day supply
- Ruxience 500 mg/50 mL injection: 8 vials per 28 day supply
- Riabni 100 mg/10 mL injection: 12 vials per 28 day supply
- Riabni 500 mg/50 mL injection: 8 vials per 28 day supply

B. Max Units (per dose and over time) [HCPCS Unit]:

Oncology Indications
<p><u>Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Leukemia (SLL):</u></p> <ul style="list-style-type: none"> • Initial therapy: <ul style="list-style-type: none"> ○ Loading dose: 100 billable units x 1 dose ○ Subsequent doses: 130 billable units every 28 days x 5 doses per 6 months • Renewal therapy: 130 billable units every 8 weeks
<p><u>ALL</u></p> <ul style="list-style-type: none"> • 100 billable units twice weekly x 18 doses
<p><u>Hairy Cell Leukemia</u></p> <ul style="list-style-type: none"> • 100 billable units weekly x 8 doses
<p><u>cGVHD</u></p> <ul style="list-style-type: none"> • 100 billable units weekly x 8 doses
<p><u>Hematopoietic Cell Transplantation</u></p> <ul style="list-style-type: none"> • Initial dose: 100 billable units x 1 dose before transplant • Subsequent doses: 250 billable units x 3 doses after transplant
<p><u>All other oncology indications:</u></p> <ul style="list-style-type: none"> • Initial therapy: 100 billable units weekly x 8 doses per 6 months • Renewal therapy: 100 billable units every 8 weeks per 6 months
Non-Oncology Indications
<p><u>Rheumatoid Arthritis (RA):</u></p> <ul style="list-style-type: none"> • 100 billable units every 14 days x 2 doses in a 16 week period
<p><u>Pemphigus Vulgaris:</u></p> <ul style="list-style-type: none"> • Initiation: 100 billable units weekly x 4 doses in a 12 month period • Maintenance: 50 billable units every 16 weeks
<p><u>GPA(WG)/MPA:</u></p> <ul style="list-style-type: none"> • Induction: 100 billable units weekly x 4 doses in a 4 month period • Initial Maintenance: 50 billable units x 2 doses in a 6 month period • Subsequent Maintenance: 50 billable units every 6 months
<p><u>All other non-oncology indications:</u></p> <ul style="list-style-type: none"> • 100 billable units weekly x 4 doses in a 6 month period

III. Initial Approval Criteria ¹⁻⁴

Coverage is provided in the following conditions:

Rituxan, Ruxience, and Rituxan Hycela are Non-Preferred products. **The Preferred products are Truxima and Riabni.**

Rituxan, Ruxience, and Rituxan Hycela may be considered medically necessary if:

- Patient has experienced a therapeutic failure or intolerance with Truxima AND Riabni; **OR**
- Rituxan, Ruxience, and Rituxan Hycela is requested for an indication for which Truxima AND Riabni have not been FDA-approved.

**RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

- Patient age is at least 18 years of age (unless otherwise specified); **AND**

Universal Criteria ¹⁻⁴

- Patient does not have a severe, active infection; **AND**
- Patient has been screened for the presence of hepatitis B (HBV) infection (i.e., HBsAg and anti-HBc) prior to initiating therapy and patients with evidence of current or prior HBV infection will be monitored for HBV reactivation during treatment; **AND**
- Patient has not received a live vaccine within 28 days prior to starting treatment and live vaccines will not be administered concurrently while on treatment; **AND**

Oncology Indications ¹⁻⁵

- Patient CD20 antigen expression is positive (*excluding use for cGVHD, Hematopoietic Cell Transplantation, and Management of Immunotherapy-Related Toxicity*); **AND**

Pediatric Mature B-Cell Acute Leukemia † ¹

- Patient is at least 6 months of age; **AND**
- Used in combination with chemotherapy for previously untreated disease

Adult Acute Lymphoblastic Leukemia (ALL) ‡ ^{5,57-59,93}

- Patient has Philadelphia chromosome-negative (Ph-) disease; **AND**
 - Used for induction/consolidation treatment; **AND**
 - Patient is < 60 years of age* without substantial comorbidities; **AND**
 - Used in combination with GRAALL-2005 regimen (daunorubicin, vincristine, prednisone, pegaspargase, and cyclophosphamide); **OR**
 - Used in combination with HyperCVAD regimen (hyper-fractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone, alternating with high-dose methotrexate and cytarabine); **OR**
 - Used in combination with Linker 4-drug regimen (daunorubicin, vincristine, pegaspargase, and prednisone); **OR**
 - Patient is ≥ 65 years of age or with substantial comorbidities (*used for induction treatment only*); **AND**
 - Used in combination with GMALL regimen (idarubicin, dexamethasone, vincristine, cyclophosphamide, and cytarabine)

**NCCN recommendations for ALL may be applicable to adolescent and young adult (AYA) patients within the age range of 15-39 years.*

Central Nervous System (CNS) Cancer ‡ ^{5,15,44,9e}

- Patient has primary CNS lymphoma; **AND**
 - Used as a component of induction therapy in combination with a methotrexate-containing regimen

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



Adult Hodgkin Lymphoma † 5,82,83

- Patient has nodular lymphocyte-predominant disease

Chronic Lymphocytic Leukemia/Small lymphocytic lymphoma (CLL/SLL) † ‡ ◊ 1-5,23e,24e,28e-30e,36e,38e,42e,43e,45e,61e,161e

- Used in combination with fludarabine and cyclophosphamide (FC); **AND**
 - Used as first-line therapy in patients < 65 years of age; **OR**
 - Used as subsequent therapy; **OR**
- Patient has disease without del(17p)/TP53 mutation; **AND**
 - Used as first-line therapy in combination with bendamustine (*patients ≥ 65 years, or younger patients with or without significant comorbidities; excluding use in frail patients [i.e., creatine clearance (CrCl) <70 mL/min]*); **AND**

- Use of rituximab in combination with bendamustine will be restricted to patients with a contraindication or intolerance to one of the following regimens:
 - Chlorambucil + obinutuzumab (*patients ≥ 65 years of age and younger patients with significant comorbidities*)
 - Fludarabine + rituximab + cyclophosphamide (*patients < 65 years of age and without significant comorbidities*); **OR**

- Used as subsequent therapy in combination with one of the following:
 - Bendamustine
 - Idelalisib
 - Lenalidomide
 - Venetoclax; **AND**

In combination with bendamustine ONLY (*patients <65 years without significant comorbidities*):

- Patient must demonstrate an inadequate response to FCR (fludarabine, cyclophosphamide, and rituximab), unless there is a contraindication or intolerance, prior to approval of rituximab in combination with bendamustine; **OR**

In combination with lenalidomide ONLY:

- Patient must demonstrate an inadequate response to one of the following regimens, unless there is a contraindication or intolerance, prior to approval of rituximab in combination with lenalidomide:
 - Venetoclax + rituximab
 - Idelalisib + rituximab

➤ FCR (fludarabine, cyclophosphamide, rituximab) (*patients <65 years without significant comorbidities*); **OR**

- Patient has disease with del(17p)/TP53 mutation; **AND**
 - Used as subsequent therapy in combination with one of the following:
 - Idelalisib
 - Lenalidomide
 - Venetoclax; **AND**

In combination with lenalidomide ONLY:

- Patient must demonstrate an inadequate response to one of the following, unless there is a contraindication or intolerance, prior to approval of rituximab in combination with lenalidomide:
 - Venetoclax + rituximab
 - Idelalisib + rituximab; **OR**

- Used as first-line therapy for histologic (Richter's) transformation to diffuse large B-cell lymphoma in combination with cyclophosphamide, doxorubicin, and vincristine based regimens or as a component of OFAR (oxaliplatin, fludarabine, cytarabine, and rituximab)

Waldenström's Macroglobulinemia/Lymphoplasmacytic Lymphoma ‡^{5,67e,72e}

Adult B-Cell Lymphomas ‡ ‡ Φ^{1-5,44} including, but not limited to, the following:

- AIDS-Related B-Cell Lymphoma ‡
 - Disease is related to Burkitt Lymphoma, diffuse large B-cell lymphoma (DLBCL), HHV-8 positive DLBCL (not otherwise specified), or primary effusion lymphoma (PEL)
- Burkitt Lymphoma ‡
 - Used in combination with chemotherapy
- Castleman Disease ‡
 - Patient has multicentric disease
- Diffuse Large B-Cell Lymphoma ‡ Φ
- Low-grade or Follicular Lymphoma ‡ Φ
- Gastric & Non-Gastric (Noncutaneous) MALT Lymphoma ‡
- High Grade B-Cell Lymphomas ‡
- Mantle Cell Lymphoma ‡
- Nodal & Splenic Marginal Zone Lymphoma ‡
- Histologic transformation of Indolent Lymphomas to Diffuse Large B-Cell Lymphoma ‡
- Post-transplant lymphoproliferative disorder (PTLD) (B-cell type) ‡

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

- Patient has had solid organ transplant or allogeneic hematopoietic stem cell transplantation

Pediatric Aggressive Mature B-Cell Lymphomas (Diffuse Large B-Cell Lymphoma, Burkitt Lymphoma, & Burkitt-like Lymphoma) †^{1,5,50,121}

- Patient is at least 6 months of age*; **AND**
- Used in combination with chemotherapy

**Pediatric Aggressive Mature B-Cell Lymphoma may be applicable to adolescent and young adult (AYA) patients older than 18 years of age and less than 39 years of age, who are treated in the pediatric oncology setting.*

Hairy Cell Leukemia ‡⁵

- Used as a single agent; **AND**
 - Used for relapsed or refractory disease if unable to receive purine analogs (i.e., cladribine or pentostatin); **OR**
- Used in combination with cladribine; **AND**
 - Used as initial therapy; **OR**
 - Used for relapsed or refractory disease; **OR**
- Used in combination with vemurafenib; **AND**
 - Used for less than complete response or relapse within 2 years of complete response following initial treatment with cladribine or pentostatin; **OR**
 - Used for progression after relapsed or refractory therapy

Chronic Graft-Versus-Host Disease (cGVHD) ‡^{5,22-25,45}

- Patient is post-allogeneic stem cell transplant (generally 3 or more months); **AND**
- Used as additional therapy in combination with corticosteroids; **AND**
- Patient has no response (e.g., steroid-refractory disease) to first-line therapy options; **AND**
- Patient must try and have an inadequate response, contraindication, or intolerance to at least a three (3) month trial of ibrutinib

Hematopoietic Cell Transplantation ‡⁵

- Used as conditioning for allogeneic transplant as part of a non-myeloablative regimen in combination with cyclophosphamide and fludarabine

Management of Immunotherapy-Related Toxicities ‡^{5,62}

- Patient has been receiving therapy with an immune checkpoint inhibitor (e.g., cemiplimab, nivolumab, pembrolizumab, atezolizumab, avelumab, durvalumab, ipilimumab, dostarlimab, etc.); **AND**
 - Patient has non-viral encephalitis related to their immunotherapy; **AND**

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

- Patient is autoimmune-encephalopathy-antibody positive; **OR**
- Patient has had limited to no improvement after 7 to 14 days on pulse-dose methylprednisolone with or without intravenous immunoglobulin (IVIG); **OR**
- Patient has bullous dermatitis related to their immunotherapy; **AND**
 - Used as additional therapy for moderate (G2), severe (G3) or life-threatening (G4) disease; **OR**
- Patient has moderate, severe, or life-threatening myalgias or myositis related to their immunotherapy that are steroid-refractory; **OR**
- Patient has severe (G3-4) myasthenia gravis related to their immunotherapy that is refractory to plasmapheresis or IVIG

Non-Oncology Indications

- Patient is not on concurrent treatment with another TNF-inhibitor, biologic response modifier or other non-biologic agent (i.e., apremilast tofacitinib, baricitinib, upadacitinib); **AND**

Rheumatoid Arthritis (RA) † 1-3,46-49,112,113

- Documented moderate to severe active disease; **AND**
- Used in combination with methotrexate unless the patient has a contraindication or intolerance; **AND**
- Patient tried and failed at least a 3 month trial with ONE oral disease modifying anti-rheumatic drug (DMARD) (e.g., methotrexate, azathioprine, auranofin, hydroxychloroquine, penicillamine, sulfasalazine, leflunomide, etc.); **AND**
- Previous failure with one or more preferred TNF antagonists at least one of which should be a self-injectable; **AND**
- Physician has assessed baseline disease severity utilizing an objective measure/tool; **AND**
- Patient has not had treatment with rituximab in the previous 4 months

Pemphigus Vulgaris † ☐ 1,10,11,35,36,38,61,80,114

- Patient has a diagnosis of pemphigus vulgaris as determined by one or more of the following clinical features:
 - Appearance of lesions, erosions and/or blisters
 - Nikolsky sign (induction of blistering via mechanical pressure at the edge of a blister or on normal skin)
 - Characteristic scarring and lesion distribution; **AND**
- Histopathologic confirmation by skin/mucous membrane biopsy; **AND**
- Positive direct immunofluorescence (DIF) microscopy result OR presence of autoantibodies as detected by indirect immunofluorescence (IIF) or enzyme-linked immunosorbent assay (ELISA); **AND**

- Patient has moderate to severe disease as assessed utilizing an objective measure/tool (i.e., PDAI, PSS, ABSIS, etc.); **AND**
- Used in combination with glucocorticoids (e.g., prednisone, prednisolone, etc.); **AND**
- Other causes of blistering or erosive skin and mucous membrane diseases have been ruled out

Granulomatosis with Polyangiitis (GPA) (Wegener’s Granulomatosis) and Microscopic Polyangiitis (MPA) † Φ^{1-4,125}

- Patient is at least 2 years of age; **AND**
- Used in combination with glucocorticoids (e.g., prednisone, methylprednisolone, etc.)

Thrombocytopenic Purpura † Φ^{6-9,63,127}

- Patient has previously failed or has a contraindication or intolerance to therapy with corticosteroids; **AND**
- Patient is at increased risk for bleeding as indicated by platelet count (within the previous 28 days) less than $30 \times 10^9/L$ (30,000/mm³); **AND**
- Patient diagnosis includes one of the following:
 - Primary thrombocytopenia or Idiopathic (Immune) thrombocytopenia purpura (ITP)
 - Evan’s syndrome

Thrombotic Thrombocytopenic Purpura (TTP) † Φ^{16-18,20,21,126}

- Patient is at increased risk for bleeding as indicated by platelet count (within the previous 28 days) less than $30 \times 10^9/L$ (30,000/mm³); **AND**
- Patient has immune-mediated or acquired disease with ADAMTS13-deficiency; **AND**
 - Used in combination with corticosteroids and therapeutic plasma exchange (TPE); **OR**
 - Used as a single agent as prophylactic therapy for patients in remission

Autoimmune Hemolytic Anemia (AIHA) † Φ²⁶⁻³²

- Patient has warm-reactive disease refractory to or dependent on glucocorticoids; **OR**
- Patient has cold agglutinin disease with symptomatic anemia, transfusion-dependence, and/or disabling circulatory symptoms

Lupus Nephritis † Φ¹¹⁵⁻¹¹⁷

- Patient has disease that is non-responsive or refractory to standard first-line therapy (e.g., mycophenolate mofetil, mycophenolic acid, cyclophosphamide, calcineurin inhibitors [e.g., tacrolimus]); **AND**
- Used as a single agent or add-on therapy in combination with mycophenolate mofetil, mycophenolic acid, cyclophosphamide

Myasthenia Gravis (unrelated to immunotherapy-related toxicity) † Φ¹¹⁸⁻¹²⁰

- Patient has muscle-specific tyrosine kinase (MuSK)-antibody positive disease; **AND**

- Patient is refractory to standard first-line therapy (e.g., glucocorticoids, azathioprine, mycophenolate mofetil, etc.)

Neuromyelitis Optica Spectrum Disorder (NMOSD) ‡⁹⁰⁻⁹²

- Patient has a confirmed diagnosis based on the following:
 - Patient was found to be seropositive for aquaporin-4 (AQP-4) IgG antibodies; **AND**
 - Patient has at least one core clinical characteristic §; **AND**
 - Alternative diagnoses have been excluded (e.g., multiple sclerosis, sarcoidosis, cancer, chronic infection, etc.); **OR**
 - Patient was found to be seronegative for AQP-4 IgG antibodies **OR** has unknown AQP-4-IgG status; **AND**
 - Patient has at least two core clinical characteristics § occurring as a result of one or more clinical attacks; **AND**
 - Patient experienced **ALL** of the following:
 - At least 1 core clinical characteristic must be optic neuritis, acute myelitis with LETM*, or area postrema syndrome; **AND**
 - Dissemination in space (≥2 different core clinical characteristics §); **AND**
 - Fulfillment of additional MRI requirements, as applicable **ψ**; **AND**
 - Alternative diagnoses have been excluded (e.g., multiple sclerosis, sarcoidosis, cancer, chronic infection, etc.); **AND**
- Used as a single agent or in combination with immunosuppressive therapy (e.g., azathioprine, methotrexate, mycophenolate, etc.)

§ Core Clinical Characteristics of NMOSD⁹⁰

- Optic neuritis
- Acute myelitis
- Area postrema syndrome: episode of otherwise unexplained hiccups or nausea and vomiting
- Acute brainstem syndrome
- Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions
- Symptomatic cerebral syndrome with NMOSD-typical brain lesions

ψ Additional MRI requirements for NMOSD without AQP4-IgG and NMOSD with unknown AQP4-IgG status⁹⁰

- Acute optic neuritis: requires brain MRI showing (a) normal findings or only nonspecific white matter lesions, **OR** (b) optic nerve MRI with T2-hyperintense lesion or T1-weighted gadolinium-enhancing lesion extending over >1/2 optic nerve length or involving optic chiasm
- Acute myelitis: requires associated intramedullary MRI lesion extending over ≥3 contiguous segments (LETM) **OR** ≥3 contiguous segments of focal spinal cord atrophy in patients with history compatible with acute myelitis
- Area postrema syndrome: requires associated dorsal medulla/area postrema lesions
- Acute brainstem syndrome: requires associated peri-ependymal brainstem lesions

*LETM = longitudinally extensive transverse myelitis lesions

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

† FDA-labeled indication(s); ‡ Compendia recommended indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria¹⁻⁴

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe infusion-related reactions, tumor lysis syndrome (TLS), severe mucocutaneous reactions, progressive multifocal leukoencephalopathy (PML), hepatitis B virus reactivation, serious bacterial, fungal, or viral infections, cardiovascular adverse reactions (e.g., ventricular fibrillation, myocardial infarction, cardiogenic shock, cardiac arrhythmias), renal toxicity, bowel obstruction or perforation, etc.; **AND**

Oncology Indications^{1-5,23-25,44,50,94-98,129,130,102-104,128}

- Patient has not exceeded dosing or duration limits as defined in Sections I, II, and V; **AND**

Acute Lymphoblastic Leukemia (ALL) only

- Treatment response or stabilization of disease as indicated by CBC, bone marrow cytogenic analysis, QPCR, or FISH

Hairy Cell Leukemia

- Coverage may not be renewed

Pediatric B-cell Acute Leukemia and Aggressive Mature B-Cell Lymphomas (induction or consolidation therapy)

- Coverage may not be renewed

Chronic Graft-Versus-Host Disease (cGVHD)

- Coverage may not be renewed

Hematopoietic Cell Transplantation

- Coverage may not be renewed

Management of Immunotherapy-Related Toxicities

- Coverage for use in the treatment of myalgias/myositis/myasthenia gravis/encephalitis may not be renewed.
- Coverage for use in bullous dermatitis: Patient has not exceeded a maximum of 18 months of therapy (4 total doses).

All Other Oncology Indications

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread

Non-Oncology Indications ^{1-5,7-12,34}

Rheumatoid Arthritis (RA)

- Disease response as indicated by improvement in signs and symptoms compared to baseline such as the number of tender and swollen joint counts, reduction of C-reactive protein, improvement of patient global assessment, and/or an improvement on a disease activity scoring tool [e.g. an improvement on a composite scoring index such as Disease Activity Score-28 (DAS28) of 1.2 points or more or a $\geq 20\%$ improvement on the American College of Rheumatology-20 (ACR20) criteria]; **AND**

Dose escalation (up to the maximum dose and frequency specified below) may occur upon clinical review on a case by case basis provided that the patient has:

- Shown an initial response to therapy; **AND**
- Received a minimum of one maintenance dose at the dose and interval specified below; **AND**
- Responded to therapy with subsequent loss of response

Thrombocytopenic Purpura (ITP or Evan's Syndrome)

- Disease response as indicated by the achievement and maintenance of a platelet count of at least $50 \times 10^9/L$ as necessary to reduce the risk for bleeding

Thrombotic Thrombocytopenic Purpura (TTP)

- Disease response as indicated by an increase in ADAMTS13 activity with a reduction in thrombotic risk

Granulomatosis with Polyangiitis (GPA) (Wegener's granulomatosis) and Microscopic polyangiitis (MPA)

- Disease response as indicated by disease control and improvement in signs and symptoms of condition compared to baseline; **AND**
- A decrease frequency in the occurrence of major relapses (defined by the reappearance of clinical and/or laboratory signs of vasculitis activity that could lead to organ failure or damage, or could be life threatening)

Pemphigus Vulgaris ^{1,10,11,35}

- Patient is currently receiving tapering doses of corticosteroids or has discontinued use of corticosteroids; **AND**
 - Disease response as indicated by complete epithelialization of lesions and improvement in signs and symptoms of condition compared to baseline; **OR**
 - Patient has not experienced continued development of new lesions, continued extension of old lesions, or failure of established lesions to begin to heal despite therapy; **OR**
 - For Relapses ONLY: Patient previously had active disease control; **AND**

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



- Patient has the appearance of 3 or more new lesions a month that do not heal spontaneously within 1 week, or by the extension of established lesions

Autoimmune Hemolytic Anemia (AIHA)

- Disease response as indicated by improvement in anemia signs and symptoms (e.g., dyspnea, fatigue, etc.) as well as: improvement in laboratory values (Hb/Hct), reduced transfusion needs, and/or reduced glucocorticoid use

Lupus Nephritis ¹¹⁵⁻¹¹⁷

- Coverage may only be renewed in patients experiencing a disease relapse (e.g., increased serum creatinine, increase in protein urine excretion, decrease in eGFR, etc.)

Myasthenia Gravis (unrelated to immunotherapy-related toxicity) ¹¹⁸⁻¹²⁰

- Disease response as indicated by a decrease in the daily dose of corticosteroids and/or an improvement in signs and symptoms compared to baseline.

NMOSD ^{90,91}

- Disease response as indicated by stabilization/improvement in any of the following: neurologic symptoms as evidenced by a decrease in acute relapses, stability reduced hospitalizations, reduction/discontinuation in plasma exchange treatments, and/or reduction/discontinuation of corticosteroids without relapse

V. Dosage/Administration ^{1-5,9,23-26,32,34,40,42,44,50,62,80,83-89,91,94-98,102-111,115-118,122-125,129,130}

Indication		Dose
CLL/SLL	Initial Therapy	375 mg/m ² intravenously (IV) weekly for 8 doses; OR 375 mg/m ² IV cycle 1, then 500 mg/m ² every 28 days cycles 2-6 (6 total doses); OR 375 mg/m ² IV cycle 1, followed by 500 mg/m ² every 2 weeks for 4 doses, then 500 mg/m ² every 28 days for 3 doses (8 total doses)
	<i>Renewal Therapy</i>	375 mg/m ² IV every 3 months; OR 500 mg/ m ² IV every 8 weeks
Adult B-Cell Lymphomas, Waldenström's, Castleman's, or Adult HL	Initial Therapy	375 mg/m ² IV once weekly for 4 - 8 doses in a 6 month period
	<i>Renewal Therapy</i>	375 mg/m ² IV once weekly for 4 doses per 6 month period; OR 375 mg/ m ² IV every 8 weeks
Pediatric Aggressive Mature B-Cell Lymphoma		Induction* [courses 1 and 2 (COPDAM1 and COPDAM2)] 375 mg/m ² IV, two doses during each of the induction courses (Day -2 and Day 1). <i>During the 1st induction course, prednisone is given as part of the chemotherapy course, and should be administered prior to rituximab. Rituximab will be given 48 hours after the first infusion of rituximab.</i> Consolidation* [courses 1 and 2 (CYM/CYVE)]

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



Indication	Dose
	<p>375 mg/m² IV, one dose during each of the consolidation courses (Day 1)</p> <p><u>Relapsed/Refractory</u></p> <p>RCYVE – 375mg/m² IV on day 1 of each 21-day cycle</p> <p>RICE – 375 mg/m² IV on days 1 and 3 of courses 1 and 2, and on day 1 only of course 3 if needed.</p> <p><i>*Note: dosing and dosing schedules are highly variable and dependent on regimen used, please refer to NCCN and PI for different protocols.</i></p>
<p>Pediatric Mature B-Cell Acute Leukemia</p>	<p><u>Induction* [courses 1 and 2 (COPDAM1 and COPDAM2)]</u></p> <p>375 mg/m² IV, two doses during each of the induction courses (Day -2 and Day 1).</p> <p><i>During the 1st induction course, prednisone is given as part of the chemotherapy course, and should be administered prior to rituximab. Rituximab will be given 48 hours after the first infusion of rituximab.</i></p> <p><u>Consolidation* [courses 1 and 2 (CYM/CYVE)]</u></p> <p>375 mg/m² IV, one dose during each of the consolidation courses (Day 1)</p> <p><i>*Note: dosing and dosing schedules are highly variable and dependent on regimen used, please refer to NCCN and PI for different protocols.</i></p>
<p>CNS Lymphoma</p>	<p><u>Intravenous administration</u></p> <p><u>Initial Therapy:</u> 375 mg/m² IV once weekly for 4 - 8 doses in a 6 month period</p> <p><u>Renewal Therapy:</u> 375 mg/m² IV once weekly for 4 doses per 6 month period; OR</p> <p>375 mg/ m² IV every 8 weeks</p> <p><u>Intrathecal/Intraventricular administration</u></p> <p>10-40 mg weekly to every 3 weeks</p>
<p>ALL</p>	<p>375 mg/m² IV up to twice weekly for a total of 16 to 18 infusions (e.g., induction [days 1 and 7], salvage reinduction when necessary [days 1 and 7], consolidation [4 infusions: blocks 1, 3, 4, and 6], late intensification [days 1 and 7], late consolidation [2 infusions: blocks 7 and 9], and maintenance [6 infusions])</p>
<p>Hairy Cell Leukemia</p>	<p>375 mg/m² IV once weekly for 4 - 8 doses</p>
<p>RA</p>	<p>1,000 mg IV on days 1 and 15, repeated every 24 weeks. May repeat up to every 16 weeks in patients requiring more frequent dosing based on clinical evaluation.</p>
<p>Pemphigus Vulgaris</p>	<p><u>Initiation</u></p> <p>1,000 IV mg on days 1 and 15; OR</p> <p>375 mg/m² IV weekly for 4 doses</p> <p><u>Maintenance</u></p>

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.
 ©2022, Magellan Rx Management



Indication	Dose
	<p>500 mg IV at month 12 and repeat every 6 months thereafter or based on clinical evaluation.</p> <p><u>Relapse</u></p> <p>1,000 IV mg upon relapse, resumption of glucocorticoids may be considered.</p> <p><i>*Subsequent infusions (maintenance and relapse) should be no sooner than 16 weeks after the previous infusion.</i></p>
AIHA	375 mg/m ² IV weekly for 4 doses in a 6 month period
Thrombocytopenic Purpura or Thrombotic Thrombocytopenic Purpura (TTP)	<p>375 mg/m² IV weekly for 4 doses; OR</p> <p>1,000 mg IV on days 1 and 15</p>
Immunotherapy Toxicity Treatment	<p><u>Bullous dermatitis</u></p> <p>1,000 mg IV every 2 weeks for 2 doses, then 500 mg IV at months 12 and 18 as needed</p> <p><u>Myalgias/Myositis</u></p> <p>375 mg/m² IV weekly for 4 doses</p> <p><u>Myasthenia Gravis</u></p> <p>375 mg/m² IV weekly for 4 doses; OR</p> <p>500 mg/m² IV every 2 weeks for 2 doses</p> <p><u>Encephalitis</u></p> <p>1,000 mg IV every 2 weeks for 2 doses; OR</p> <p>375 mg/m² IV weekly for 4 doses</p>
GPA (WG), MPA	<p><u>Induction (Pediatric and Adult)</u></p> <p>375 mg/m² IV weekly for 4 doses; OR</p> <ul style="list-style-type: none"> – Adults: 1,000 mg IV on days 1 and 15; OR – Pediatric (up to a maximum of 1,000 mg per dose): <ul style="list-style-type: none"> ○ 575 mg/m² IV on days 1 and 15 (BSA ≤1.5m²) ○ 750 mg/m² IV on days 1 and 15 (BSA >1.5m²) <p><u>Maintenance</u></p> <ul style="list-style-type: none"> – Pediatric: <ul style="list-style-type: none"> ○ 250 mg/m² IV on days 1 and 15, then 250 mg/m² IV every 6 months thereafter based on clinical evaluation – Adult: <ul style="list-style-type: none"> ○ 500 mg IV on days 1 and 15, then 500 mg IV every 6 months thereafter based on clinical evaluation. <p><i>*Initial MAINTENANCE infusions should be no sooner than 16 weeks and no later than 24 weeks after the previous infusion if rituximab was used for initial induction therapy.</i></p>

Indication	Dose
	<i>*Initial MAINTENANCE infusions should be initiated within 4 weeks following disease control when initial induction occurred with other standard of care immunosuppressants.</i>
cGVHD	375 mg/m ² IV weekly for 4 doses, then 375 mg/m ² IV monthly for 4 months -OR- 375 mg/m ² IV weekly for 4 doses (Note: A second course of 4 weekly doses may be administered 8 weeks after initial therapy for patients with lack of or incomplete response.) -OR- 375 mg/m ² IV weekly for 4 - 8 doses
Hematopoietic Cell Transplantation	375 mg/m ² IV for 1 day before transplant, then 1000 mg/m ² IV on days 1, 8, and 15 after transplant
NMOSD	1,000 mg IV once on days 1 and 15, repeat every 6 months -OR- 375 mg/m ² once weekly for 4 weeks, repeat every 6 months
Lupus Nephritis	1,000 mg IV on days 1 and 15 -OR- 375 mg/m ² IV once weekly for 4 doses
Myasthenia Gravis (unrelated to immunotherapy-related toxicity)	1,000 mg IV on days 1 and 15, may repeat a full or partial course every 6 months -OR- 375 mg/m ² IV once weekly for 4 doses, may repeat a full or partial course every 6 months
<i>Abbreviations: COP = Cyclophosphamide, Oncovin (vincristine), Prednisone; COPDAM = Cyclophosphamide, Oncovin (vincristine), Prednisolone, Adriamycin (doxorubicin), Methotrexate; CYM = CYtarabine (Aracytine, Ara-C), Methotrexate; CYVE = CYtarabine (Aracytine, Ara-C), VEposide (VP16)</i>	

VI. Billing Code/Availability Information

HCPCS Code:

- J9312 – Injection, rituximab, 10 mg; 1 billable unit = 10 mg (*Rituxan IV only*)
- Q5115 – Injection, rituximab-abbs, biosimilar, (truxima), 10 mg; 1 billable unit = 10 mg
- Q5119 – Injection, rituximab-pvvr, biosimilar, (ruxience), 10 mg; 1 billable unit = 10 mg
- Q5123 – Injection, rituximab-arrx, biosimilar, (riabni), 10 mg; 1 billable unit = 10 mg

NDC(s):

- Rituxan 100 mg/10 mL single-dose vial for injection: 50242-0051-xx
- Rituxan 500 mg/50 mL single-dose vial for injection: 50242-0053-xx
- Truxima 100 mg/10 mL single-dose vial for injection: 63459-0103-xx

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™) Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

- Truxima 500 mg/50 mL single-dose vial for injection: 63459-0104-xx
- Ruxience 100 mg/10 mL single-dose vial for injection: 00069-0238-xx
- Ruxience 500 mg/50 mL single-dose vial for injection: 00069-0249-xx
- Riabni 100 mg/10 mL single-dose vial for injection: 55513-0224-xx
- Riabni 500 mg/50 mL single-dose vial for injection: 55513-0326-xx

VII. References (STANDARD)

1. Rituxan [package insert]. South San Francisco, CA; Genentech, Inc; December 2021. Accessed June 2022.
2. Truxima [package insert]. Incheon, Korea; Celltrion, Inc; February 2022. Accessed June 2022.
3. Ruxience [package insert]. New York, NY; Pfizer, Inc; November 2021. Accessed June 2022.
4. Riabni [package insert]. Thousand Oaks, CA; Amgen, Inc; December 2020. Accessed June 2022.
5. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) rituximab. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2022.
6. Arnold DM, Dentali F, Crowther MA, et al. Systematic review: efficacy and safety of rituximab for adults with idiopathic thrombocytopenic purpura. *Ann Intern Med* 2007; 146:25-33.
7. Zaja F, Baccarani M, Mazza P, et al: Dexamethasone plus rituximab yields higher sustained response rates than dexamethasone monotherapy in adults with primary immune thrombocytopenia. *Blood* 2010; 115(14):2755-2762.
8. Stasi R, Pagano A, Stipa E, et al: Rituximab chimeric anti-CD10 monoclonal antibody treatment for adults with chronic idiopathic thrombocytopenic purpura. *Blood* 2001; 98(4):952-957.
9. Neunert C, Lim W, Crowther M, Cohen A, Solberg L Jr, Crowther MA. The American Society of Hematology 2011 evidence-based practice guideline for immune thrombocytopenia. *Blood*. 117(16):4190-4207.
10. Joly P, Mouquet H, Roujeau JC, et al. A single cycle of rituximab for the treatment of severe pemphigus. *N Engl J Med* 2007; 357:545-52.
11. Ahmed AR, Spigelman Z, Cavacini LA et al. Treatment of pemphigus vulgaris with rituximab and intravenous immune globulin. *N Engl J Med* 2006; 355:1772-9.
12. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res (Hoboken)*. 2015 Nov 6. doi: 10.1002/acr.22783.

13. Smolen JS, Landewé R, Bijlsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis*. 2017 Mar 6. pii: annrheumdis-2016-210715.
14. González-Barca E, Domingo-Domenech E, Capote FJ, et al. Prospective phase II trial of extended treatment with rituximab in patients with B-cell post-transplant lymphoproliferative disease. *Haematologica*. 2007 Nov; 92(11):1489-94.
15. Chamberlain MC, Johnston SK, Van Horn A, et al. Recurrent lymphomatous meningitis treated with intra-CSF rituximab and liposomal ara-C. *J Neurooncol*. 2009 Feb;91(3):271-7.
16. Scully M, Cohen H, Cavenagh J, et al. Remission in acute refractory and relapsing thrombotic thrombocytopenic purpura following rituximab is associated with a reduction in IgG antibodies to ADAMTS-13. *Br J Haematol* 2007;136:451-461.
17. Fakhouri F, Vernant JP, Veyradier A, et al. Efficiency of curative and prophylactic treatment with rituximab in ADAMTS13-deficient thrombotic thrombocytopenic purpura: a study of 11 cases. *Blood*. 2005;106:1932-37.
18. Elliott MA, Heit JA, Rajiv K, et al. Rituximab for refractory and or relapsing thrombotic thrombocytopenic purpura related to immune-mediated severe ADAMTS13-deficiency: a report of four cases and a systematic review of the literature. *Eur J Haematol* 2009. Epub ahead of print, doi:10.1111/j.1600-0609.2009.01292.
19. Scully M, McDonald V, Cavenagh J, et al. A phase 2 study of the safety and efficacy of rituximab with plasma exchange in acute acquired thrombotic thrombocytopenic purpura. *Blood*. 2011;118(7):1746-1753.
20. Tun NM, Villani GM. Efficacy of rituximab in acute refractory or chronic relapsing non-familial idiopathic thrombotic thrombocytopenic purpura: a systematic review with pooled data analysis. *J Thromb Thrombolysis*. 2012;34(3):347-359.
21. Froissart A, Buffet M, Veyradier A, et al: Efficacy and safety of first-line rituximab in severe, acquired thrombotic thrombocytopenic purpura with a suboptimal response to plasma exchange. Experience of the French Thrombotic Microangiopathies Reference Center. *Crit Care Med* 2012; 40(1):104-111.
22. van Dorp S, Resemann H, te Boome L, et al. The immunological phenotype of rituximab-sensitive chronic graft-versus-host disease: a phase II study. *Haematologica* 2011;96(9):1380-1384.
23. Kim SJ, Lee JW, Jung CW, et al. Weekly rituximab followed by monthly rituximab treatment for steroid-refractory chronic graft-versus-host disease: results from a prospective, multicenter, phase II study. *Haematologica* 2010;95(11):1935-1942.
24. Cutler C, Miklos D, Kim HT, et al. Rituximab for Steroid-Refractory Chronic Graft-Versus-Host Disease. *Blood*. 2006, 108(2):756-62.
25. Wolff D, Schleuning M, von Harsdorf S, et al. Consensus Conference on Clinical Practice in Chronic GVHD: Second-Line Treatment of Chronic Graft-versus-Host Disease. *Biol Blood Marrow Transplant*. 2011 Jan;17(1):1-17. doi: 10.1016/j.bbmt.2010.05.011.

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



26. Frame JN, Fichtner R, McDevitt PW. Rituximab for the treatment of autoimmune hemolytic anemia (AIHA) in adults: an analysis of literature reports in 92 patients. *Blood* 2004;104:Abstract 3721.
27. Birgens H, Frederiksen H, Hasselbalch HC, et al: A phase III randomized trial comparing glucocorticoid monotherapy versus glucocorticoid and rituximab in patients with autoimmune haemolytic anaemia. *Br J Haematol* 2013; 163(3):393-399.
28. Schollkopf C, Kjeldsen L, Bjerrum OW, et al: Rituximab in chronic cold agglutinin disease: a prospective study of 20 patients. *Leuk Lymphoma* 2006; 47(N2):253-260.
29. Berentsen S, Ulvestad E, Gjertsen BT, et al: Rituximab for primary chronic cold agglutinin disease: a prospective study of 37 courses of therapy in 27 patients. *Blood* 2004; 103(8):2925-2928.
30. Reynaud Q, Durieu I, Dutertre M, et al. Efficacy and safety of rituximab in auto-immune hemolytic anemia: A meta-analysis of 21 studies. *Autoimmun Rev.* 2015;14(4):304-313.
31. Barcellini W, Zaja F, Zaninoni A, et al, "Low-dose Rituximab in Adult Patients With Idiopathic Autoimmune Hemolytic Anemia: Clinical Efficacy and Biologic Studies," *Blood*, 2012, 119(16):3691-7.
32. Roumier M, Loustau V, Guillaud C, et al. Characteristics and outcome of warm autoimmune hemolytic anemia in adults: New insights based on a single-center experience with 60 patients. *Am J Hematol.* 2014;89(9):E150-E155.
33. Gobert D, Bussel JB, Cunningham-Rundles C, et al. Efficacy and safety of rituximab in common variable immunodeficiency-associated immune cytopenias: a retrospective multicentre study on 33 patients. *Br J Haematol.* 2011;155(4):498-508.
34. YW Shin, ST Lee, KI Park, et al. Treatment strategies for autoimmune encephalitis. *Ther Adv Neurol Disord.* 2017 Aug 16;11:1756285617722347. doi: 10.1177/1756285617722347. eCollection 2018. Review.
35. Murrell DF, Dick S, Ahmed AR, et al. Consensus statement on definitions of disease, end points, and therapeutic response for pemphigus. *J Am Acad Dermatol.* 2008 June; 58(6): 1043–1046. doi:10.1016/j.jaad.2008.01.012. Avail at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2829665/pdf/nihms82304.pdf>
36. Grover, S. Scoring Systems in Pemphigus. *Indian J Dermatol.* 2011 Mar-Apr; 56(2): 145–149. doi: 10.4103/0019-5154.80403
37. Daniel BS, Hertl M, Weth VP, et al. Severity score indexes for blistering diseases. *Clin Dermatol.* 2012 Jan-Feb; 30(1): 108–113. doi: 10.1016/j.clindermatol.2011.03.017
38. Joly P, Litrowski N. Pemphigus group (vulgaris, vegetans, foliaceus, herpetiformis, brasiliensis). *Clin Dermatol* 2011; 29:432.
39. Lambert MP, Gernsheimer TB. Clinical updates in adult immune thrombocytopenia. *Blood.* 2017. 129:2829-2835. doi:10.1182/blood-2017-03-754119

40. Schulz H, Pels H, Schmidt-Wolf I, et al. Intraventricular treatment of relapsed central nervous system lymphoma with the anti-CD20 antibody rituximab. *Haematologica* January 2004 89: 753-754
41. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. *J Oncol Pract*. 2018 Mar;14(3):e130-e136.
42. Hematology/Oncology Pharmacy Association (2019). *Intravenous Cancer Drug Waste Issue Brief*. Retrieved from http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug_Waste_2019.pdf
43. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. *BMJ*. 2016 Feb 29;352:i788.
44. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for B-Cell Lymphomas 3.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2022.
45. Imbruvica [package insert]. Horsham, PA; Janssen Biotech, Inc. December 2020. Accessed November 2021.
46. Keystone E, Burmester GR, Furie R, et al. Improvement in patient-reported outcomes in a rituximab trial in patients with severe rheumatoid arthritis refractory to anti-tumor necrosis factor therapy. *Arthritis Rheum*. 2008 Jun 15;59(6):785-93. doi: 10.1002/art.23715.
47. Mease PJ, Cohen S, Gaylis NB, et al. Efficacy and Safety of Retreatment in Patients with Rheumatoid Arthritis with Previous Inadequate Response to Tumor Necrosis Factor Inhibitors: Results from the SUNRISE Trial. *The Journal of Rheumatology* May 2010, 37 (5) 917-927; DOI: <https://doi.org/10.3899/jrheum.090442>
48. Tak PP, Rigby W, Rubbert-Roth A, et al. Sustained inhibition of progressive joint damage with rituximab plus methotrexate in early active rheumatoid arthritis: 2-year results from the randomised controlled trial IMAGE. *Ann Rheum Dis*. 2012 Mar;71(3):351-7. doi: 10.1136/annrheumdis-2011-200170. Epub 2011 Oct 19.
49. Emery P, Deodhar A, Rigby WF, et al. Efficacy and safety of different doses and retreatment of rituximab: a randomised, placebo-controlled trial in patients who are biological naive with active rheumatoid arthritis and an inadequate response to methotrexate (Study Evaluating Rituximab's Efficacy in MTX iNadequate rEsponders (SERENE)). *Ann Rheum Dis*. 2010 Sep;69(9):1629-35. doi: 10.1136/ard.2009.119933. Epub 2010 May 20. Erratum in: *Ann Rheum Dis*. 2011 Aug;70(8):1519.
50. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Pediatric Aggressive Mature B-Cell Lymphomas 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2022.

51. Lee KH, Lee J, Bae JS, et al. Analytical similarity assessment of rituximab biosimilar CT-P10 to reference medicinal product. *MAbs*. 2018;10(3):380-396
52. Ogura M, Sancho JM, Cho S-G, et al. Efficacy, pharmacokinetics, and safety of the biosimilar CT-P10 in comparison with rituximab in patients with previously untreated low-tumour-burden follicular lymphoma: a randomised, double-blind, parallel-group phase 3 trial. *Lancet Haematol*. 2018;5:e543-e553.
53. Gulácsi L, Brodszky V, Baji P, et al. The rituximab biosimilar CT-P10 in rheumatology and cancer: a budget impact analysis in 28 European countries. *Adv Ther*. 2017; 34: 1128-1144.
54. Yoo DH, Suh CH, Shim SC, et al. A multicentre randomised controlled trial to compare the pharmacokinetics, efficacy and safety of CT-P10 and innovator rituximab in patients with rheumatoid arthritis. *Ann Rheum Dis*. 2017; 76: 566-570.
55. Suh C, Berrocal Kasay A, Chalouhi El-Khoury E, et al. Pharmacokinetics and safety of three formulations of rituximab (CT-P10, US-sourced innovator rituximab and EU-sourced innovator rituximab) in patients with rheumatoid arthritis: results from phase 3 randomized controlled trial over 24 weeks. *Arthritis Rheumatol*. 2016; 68: 1634.
56. Kim WS, Buske C, Ogura M, et al. Efficacy, pharmacokinetics, and safety of the biosimilar CT-P10 compared with rituximab in patients with previously untreated advanced-stage follicular lymphoma: a randomised, double-blind, parallel-group, non-inferiority phase 3 trial. *Lancet Haematol*. 2017; 4: e362-e373.
57. Cohen S, Emery P, Greenwald M, et al. A phase I pharmacokinetics trial comparing PF-05280586 (a potential biosimilar) and rituximab in patients with active rheumatoid arthritis. *Br J Clin Pharmacol*. 2016 Jul;82(1):129-38.
58. Williams JH, Hutmacher MM, Zierhut ML, et al. Comparative assessment of clinical response in patients with rheumatoid arthritis between PF-05280586, a proposed rituximab biosimilar, and rituximab. *Br J Clin Pharmacol*. 2016 Dec;82(6):1568-1579.
59. Sharman JP, Liberati AM, Ishizawa K, et al. A Randomized, Double-Blind, Efficacy and Safety Study of PF-05280586 (a Rituximab Biosimilar) Compared with Rituximab Reference Product (MabThera®) in Subjects with Previously Untreated CD20-Positive, Low-Tumor-Burden Follicular Lymphoma (LTB-FL). *BioDrugs*. 2019 Dec 9. doi: 10.1007/s40259-019-00398-7.
60. Cohen SB, Burgos-Vargas R, Emery P, et al. Comparative assessment of clinical response in patients with rheumatoid arthritis between PF-05280586, a proposed rituximab biosimilar, and rituximab. *Br J Clin Pharmacol*. 2016 Dec;82(6):1568-1579.
61. Murrel DF, Peña S, Joly P, et al. Diagnosis and management of pemphigus: Recommendations of an international panel of experts. *JAAD*: Mar2020;82;3;575-585. DOI:<https://doi.org/10.1016/j.jaad.2018.02.021>.
62. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Management of Immunotherapy-Related Toxicities 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2022.

63. Neunert C, Terrell DR, Arnold DM, et al. American Society of Hematology 2019 guidelines for immune thrombocytopenia [published correction appears in *Blood Adv.* 2020 Jan 28;4(2):252]. *Blood Adv.* 2019;3(23):3829-3866. Doi:10.1182/bloodadvances.2019000966.
64. McLaughlin P, Grillo-López AJ, Link BK, et al. Rituximab chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to a four-dose treatment program. *J Clin Oncol.* 1998 Aug;16(8):2825-33.
65. Piro LD, White CA, Grillo-López AJ, et al. Extended Rituximab (anti-CD20 monoclonal antibody) therapy for relapsed or refractory low-grade or follicular non-Hodgkin's lymphoma. *Ann Oncol.* 1999;10(6):655-661. Doi:10.1023/a:1008389119525.
66. Davis TA, Grillo-López AJ, White CA, et al. Rituximab anti-CD20 monoclonal antibody therapy in non-Hodgkin's lymphoma: safety and efficacy of re-treatment. *J Clin Oncol.* 2000;18(17):3135-3143. Doi:10.1200/JCO.2000.18.17.3135.
67. Marcus R, Imrie K, Belch A, et al. CVP chemotherapy plus rituximab compared with CVP as first-line treatment for advanced follicular lymphoma. *Blood.* 2005;105(4):1417-1423. Doi:10.1182/blood-2004-08-3175.
68. Salles G, Seymour JF, Offner F, et al. Rituximab maintenance for 2 years in patients with high tumour burden follicular lymphoma responding to rituximab plus chemotherapy (PRIMA): a phase 3, randomised controlled trial [published correction appears in *Lancet.* 2011 Apr 2;377].
69. Hochster H, Weller E, Gascoyne RD, et al. Maintenance rituximab after cyclophosphamide, vincristine, and prednisone prolongs progression-free survival in advanced indolent lymphoma: results of the randomized phase III ECOG1496 Study. *J Clin Oncol.* 2009;27(10):1607-1614. Doi:10.1200/JCO.2008.17.1561.
70. Habermann TM, Weller EA, Morrison VA, et al. Rituximab-CHOP versus CHOP alone or with maintenance rituximab in older patients with diffuse large B-cell lymphoma. *J Clin Oncol.* 2006;24(19):3121-3127. Doi:10.1200/JCO.2005.05.1003,
71. Coiffier B, Thieblemont C, Van Den Neste E, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: a study by the Groupe d'Etudes des Lymphomes de l'Adulte. *Blood.* 2010;116(12):2040-2045. Doi:10.1182/blood-2010-03-276246.
72. Pfreundschuh M, Kuhnt E, Trümper L, et al. CHOP-like chemotherapy with or without rituximab in young patients with good-prognosis diffuse large-B-cell lymphoma: 6-year results of an open-label randomised study of the MabThera International Trial (MinT) Group. *Lancet Oncol.* 2011;12(11):1013-1022. Doi:10.1016/S1470-2045(11)70235-2.
73. Dakhil S, Hermann R, Schreeder MT, et al. Phase III safety study of rituximab administered as a 90-minute infusion in patients with previously untreated diffuse large B-cell and

- follicular lymphoma. *Leuk Lymphoma*. 2014;55(10):2335-2340.
Doi:10.3109/10428194.2013.877135.
74. Fischer K, Bahlo J, Fink AM, et al. Long-term remissions after FCR chemoimmunotherapy in previously untreated patients with CLL: updated results of the CLL8 trial. *Blood*. 2016;127(2):208-215. Doi:10.1182/blood-2015-06-651125.
75. Robak T, Dmoszynska A, Solal-Céligny P, et al. Rituximab plus fludarabine and cyclophosphamide prolongs progression-free survival compared with fludarabine and cyclophosphamide alone in previously treated chronic lymphocytic leukemia. *J Clin Oncol*. 2010 Apr 1;28(10):1756-65.
76. Stone JH, Merkel PA, Spiera R, et al. Rituximab versus cyclophosphamide for ANCA-associated vasculitis. *N Engl J Med*. 2010;363(3):221-232. Doi:10.1056/NEJMoa0909905.
77. Guillevin L, Pagnoux C, Karras A, et al. Rituximab versus azathioprine for maintenance in ANCA-associated vasculitis. *N Engl J Med*. 2014;371(19):1771-1780.
Doi:10.1056/NEJMoa1404231.
78. Niles JL, Merkel PA, Mertz L, et al. Long-Term Safety of Rituximab in Granulomatosis with Polyangiitis or Microscopic Polyangiitis: Results of the Four-Year Study of Rituximab in ANCA-Associated Vasculitis Registry [abstract]. *Arthritis Rheumatol*. 2018; 70 (suppl 10).
79. Brogan P, Cleary G, Hersh AO, et al. Pediatric Open-Label Clinical Study of Rituximab for the Treatment of Granulomatosis with Polyangiitis (GPA) and Microscopic Polyangiitis (MPA) [abstract]. *Arthritis Rheumatol*. 2018; 70 (suppl 10).
80. Joly P, Maho-Vaillant M, Prost-Squarcioni C, et al. First-line rituximab combined with short-term prednisone versus prednisone alone for the treatment of pemphigus (Ritux 3): a prospective, multicentre, parallel-group, open-label randomised trial. *Lancet*. 2017;389(10083):2031-2040. Doi:10.1016/S0140-6736(17)30070-3.
81. Thomas DA, O'Brien S, Faderl S, et al. Chemoimmunotherapy with a modified hyper-CVAD and rituximab regimen improves outcome in de novo Philadelphia chromosome-negative precursor B-lineage acute lymphoblastic leukemia. *J Clin Oncol*. 2010;28(24):3880-3889.
Doi:10.1200/JCO.2009.26.9456.
82. Kadia TM, Kantarjian HM, Thomas DA, et al. Phase II study of methotrexate, vincristine, pegylated-asparaginase, and dexamethasone (MopAD) in patients with relapsed/refractory acute lymphoblastic leukemia. *Am J Hematol*. 2015;90(2):120-124. Doi:10.1002/ajh.23886.
83. Goldman S, Smith L, Anderson JR, et al. Rituximab and FAB/LMB 96 chemotherapy in children with Stage III/IV B-cell non-Hodgkin lymphoma: a Children's Oncology Group report. *Leukemia*. 2013;27(5):1174-1177. doi:10.1038/leu.2012.255.
84. Griffin TC, Weitzman S, Weinstein H, et al. A study of rituximab and ifosfamide, carboplatin, and etoposide chemotherapy in children with recurrent/refractory B-cell (CD20+) non-Hodgkin lymphoma and mature B-cell acute lymphoblastic leukemia: a report from the Children's Oncology Group. *Pediatr Blood Cancer*. 2009;52(2):177-181. doi:10.1002/pbc.21753.

85. Choquet S, Leblond V, Herbrecht R, et al. Efficacy and safety of rituximab in B-cell post-transplantation lymphoproliferative disorders: results of a prospective multicenter phase 2 study. *Blood*. 2006;107(8):3053-3057. doi:10.1182/blood-2005-01-0377.
86. Trappe R, Oertel S, Leblond V, et al. Sequential treatment with rituximab followed by CHOP chemotherapy in adult B-cell post-transplant lymphoproliferative disorder (PTLD): the prospective international multicentre phase 2 PTLD-1 trial. *Lancet Oncol*. 2012;13(2):196-206.
87. Ghobrial IM, Hong F, Padmanabhan S, et al. Phase II trial of weekly bortezomib in combination with rituximab in relapsed or relapsed and refractory Waldenstrom macroglobulinemia. *J Clin Oncol*. 2010;28(8):1422-1428. doi:10.1200/JCO.2009.25.3237.
88. Advani RH, Horning SJ, Hoppe RT, et al. Mature results of a phase II study of rituximab therapy for nodular lymphocyte-predominant Hodgkin lymphoma. *J Clin Oncol*. 2014;32(9):912-918. doi:10.1200/JCO.2013.53.2069.
89. Godeau B, Porcher R, Fain O, et al. Rituximab efficacy and safety in adult splenectomy candidates with chronic immune thrombocytopenic purpura: results of a prospective multicenter phase 2 study. *Blood*. 2008;112(4):999-1004. doi:10.1182/blood-2008-01-131029.
90. Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology*. 2015 Jul;85(2):177-89. Epub 2015 Jun 19.
91. Trebst C, Jarius S, Berthele A, et al. Update on the diagnosis and treatment of neuromyelitis optica: recommendations of the Neuromyelitis Optica Study Group (NEMOS). *J Neurol* 2014; 261:1.
92. Nikoo Z, Badihian S, Shaygannejad V, et al. Comparison of the efficacy of azathioprine and rituximab in neuromyelitis optica spectrum disorder: a randomized clinical trial. *J Neurol*. 2017;264(9):2003. Epub 2017 Aug 22.
93. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Acute Lymphoblastic Leukemia 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2022.
94. Thomas DA, O'Brien S, Bueso-Ramos C, et al. Rituximab in relapsed or refractory hairy cell leukemia. *Blood*. 2003 Dec 1;102(12):3906-11. doi: 10.1182/blood-2003-02-0630.
95. Nieva J, Bethel K, Saven A. Phase 2 study of rituximab in the treatment of cladribine-failed patients with hairy cell leukemia. *Blood*. 2003 Aug 1;102(3):810-3.
96. Chihara D, Kantarjian H, O'Brien S, et al. Long-term durable remission by cladribine followed by rituximab in patients with hairy cell leukaemia: update of a phase II trial. *Br J Haematol*. 2016 Sep;174(5):760-6.

97. Else M, Dearden CE, Matutes E, et al. Rituximab with pentostatin or cladribine: an effective combination treatment for hairy cell leukemia after disease recurrence. *Leuk Lymphoma*. 2011 Jun;52 Suppl 2:75-8. doi: 10.3109/10428194.2011.568650.
98. Zenhäusern R, Simcock M, Gratwohl A, et al; Swiss Group for Clinical Cancer Research (SAKK). Rituximab in patients with hairy cell leukemia relapsing after treatment with 2-chlorodeoxyadenosine (SAKK 31/98). *Haematologica*. 2008 Sep;93(9):1426-8.
99. Birgens H, Frederiksen H, Hasselbalch HC, et al. A phase III randomized trial comparing glucocorticoid monotherapy versus glucocorticoid and rituximab in patients with autoimmune haemolytic anaemia. *Br J Haematol*. 2013 Nov;163(3):393-9. doi: 10.1111/bjh.12541.
100. Niederwieser, D., Hamm, C., Cobb, P. et al. Efficacy and Safety of ABP 798: Results from the JASMINE Trial in Patients with Follicular Lymphoma in Comparison with Rituximab Reference Product. *Targ Oncol* 15, 599–611 (2020). <https://doi.org/10.1007/s11523-020-00748-4>.
101. Burmester, G., Drescher, E., Hrycaj, P. et al. Efficacy and safety results from a randomized double-blind study comparing proposed biosimilar ABP 798 with rituximab reference product in subjects with moderate-to-severe rheumatoid arthritis. *Clin Rheumatol* 39, 3341–3352 (2020). <https://doi.org/10.1007/s10067-020-05305-y>.
102. Solimando AG, Crudele L, Leone P, et al. Immune Checkpoint Inhibitor-Related Myositis: From Biology to Bedside. *Int J Mol Sci*. 2020;21(9):3054. Published 2020 Apr 26. doi:10.3390/ijms21093054.
103. Kong SS, Chen YJ, Su IC, et al; CHEESE Study Group. Immunotherapy for anti-NMDA receptor encephalitis: Experience from a single center in Taiwan. *Pediatr Neonatol*. 2019 Aug;60(4):417-422. doi: 10.1016/j.pedneo.2018.10.006.
104. Feng S, Coward J, McCaffrey E, et al. Pembrolizumab-Induced Encephalopathy: A Review of Neurological Toxicities with Immune Checkpoint Inhibitors. *J Thorac Oncol*. 2017 Nov;12(11):1626-1635. doi: 10.1016/j.jtho.2017.08.007.
105. Chamberlain MC, Johnston SK, Van Horn A, et al. Recurrent lymphomatous meningitis treated with intra-CSF rituximab and liposomal ara-C. *J Neurooncol*. 2009 Feb;91(3):271-7. doi: 10.1007/s11060-008-9707-1. Epub 2008 Sep 27.
106. Rituximab in the treatment of Rosai-Dorfman syndrome with IgG4 disease. *Journal of the American Academy of Dermatology* 2019; 81: AB269.
107. Abla O, Jacobsen E, Picarsic J, et al. Consensus recommendations for the diagnosis and clinical management of Rosai-Dorfman-Destombes disease. *Blood* (2018) 131 (26): 2877–2890.
108. Maury S, Chevret S, Thomas X, et al; for GRAALL. Rituximab in B-Lineage Adult Acute Lymphoblastic Leukemia. *N Engl J Med*. 2016 Sep 15;375(11):1044-53. doi: 10.1056/NEJMoa1605085.
109. Wieduwilt MJ, Jonas BA, Schiller GJ, et al; A Phase II Study of Pegylated Asparaginase, Cyclophosphamide, Rituximab, and Dasatinib Added to the UCSF 8707 (Linker 4-drug) Regimen with Liposomal Cytarabine CNS Prophylaxis for Adults with Newly Diagnosed

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



- Acute Lymphoblastic Leukemia (ALL) or Lymphoblastic Lymphoma (LBL): University of California Hematologic Malignancies Consortium Study (UCHMC) 1401. *Blood* 2018; 132 (Supplement 1): 4018. doi: <https://doi.org/10.1182/blood-2018-99-117469>.
110. Thomas DA, Faderl S, O'Brien S, et al. Chemoimmunotherapy with hyper-CVAD plus rituximab for the treatment of adult Burkitt and Burkitt-type lymphoma or acute lymphoblastic leukemia. *Cancer*, 106: 1569-1580. <https://doi.org/10.1002/cncr.21776>.
111. Kreitman RJ, Wilson W, Calvo KR, et al. Cladribine with immediate rituximab for the treatment of patients with variant hairy cell leukemia. *Clin Cancer Res*. 2013 Dec 15;19(24):6873-81. doi: 10.1158/1078-0432.CCR-13-1752.
112. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol*. 2021 Jul;73(7):1108-1123. doi: 10.1002/art.41752.
113. Smolen JS, Landewé RBM, Bijlsma JWJ, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Annals of the Rheumatic Diseases* 2020;79:685-699.
114. Venugopal SS, Murrell DF. Diagnosis and clinical features of pemphigus vulgaris. *Dermatol Clin*. 2011 Jul;29(3):373-80, vii. doi: 10.1016/j.det.2011.03.004. PMID: 21605802.
115. Fanouriakis A, Kostopoulou M, Cheema K, et al: 2019 update of the Joint European League Against Rheumatism and European Renal Association-European Dialysis and Transplant Association (EULAR/ERA-EDTA) recommendations for the management of lupus nephritis. *Ann Rheum Dis* 2020; 79(6):713-723.
116. Vigna-Perez M, Hernández-Castro B, Paredes-Saharopulos O, et al. Clinical and immunological effects of Rituximab in patients with lupus nephritis refractory to conventional therapy: a pilot study. *Arthritis Res Ther*. 2006;8(3):R83. doi: 10.1186/ar1954. Epub 2006 May 5.
117. Melander C, Sallée M, Trolliet P, et al. Rituximab in severe lupus nephritis: early B-cell depletion affects long-term renal outcome. *Clin J Am Soc Nephrol*. 2009 Mar;4(3):579-87. doi: 10.2215/CJN.04030808. Epub 2009 Mar 4.
118. Bird SJ. (2021). Chronic immunosuppressive therapy for myasthenia gravis. In Shefner JM, Goddeau RP (Eds.), *UptoDate*. Available from https://www.uptodate.com/contents/chronic-immunosuppressive-therapy-for-myasthenia-gravis?sectionName=REFRACTORY%20DISEASE&search=myasthenia%20gravis&topicRef=5157&anchor=H2442874626&source=see_link#H5661854.
119. Topakian R, Zimprich F, Iglseider S, et al. High efficacy of rituximab for myasthenia gravis: a comprehensive nationwide study in Austria. *J Neurol*. 2019;266(3):699-706. doi:10.1007/s00415-019-09191-6.
120. Li T, Zhang GQ, Li Y, et al. Efficacy and safety of different dosages of rituximab for refractory generalized AChR myasthenia gravis: a meta-analysis. *J Clin Neurosci*. 2021;85:6-12. Doi:10.1016/j.jocn.2020.11.043.

121. Colin V, Auperin A, Pillon M, et al. Rituximab for High-Risk, Mature B-Cell Non-Hodgkin's Lymphoma in Children. *Clinical Trial. N Engl J Med.* 2020 Jun 4;382(23):2207-2219. doi: 10.1056/NEJMoa1915315.
122. Furman RR, Sharman JP, Coutre SE, et al. Idelalisib and rituximab in relapsed chronic lymphocytic leukemia. *N Engl J Med.* 2014;370(11):997-1007. doi:10.1056/NEJMoa1315226.
123. Greil R, Obrtlíková P, Smolej L, et al. Rituximab maintenance versus observation alone in patients with chronic lymphocytic leukaemia who respond to first-line or second-line rituximab-containing chemoimmunotherapy: final results of the AGMT CLL-8a Maintenance randomised trial. *Lancet Haematol.* 2016 Jul;3(7):e317-29. doi: 10.1016/S2352-3026(16)30045-X.
124. Dartigeas C, Van Den Neste E, Léger J, et al. Rituximab maintenance versus observation following abbreviated induction with chemoimmunotherapy in elderly patients with previously untreated chronic lymphocytic leukaemia (CLL 2007 SA): an open-label, randomised phase 3 study. *Lancet Haematol.* 2018 Feb;5(2):e82-e94. doi: 10.1016/S2352-3026(17)30235-1.
125. Chung SA, Langford CA, Maz M, et al. 2021 American College of Rheumatology/Vasculitis Foundation Guideline for the Management of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. *Arthritis Care Res (Hoboken).* 2021;73(8):1088. Epub 2021 Jul 8.
126. Zheng XL, Vesely SK, Cataland SR, et al. ISTH guidelines for treatment of thrombotic thrombocytopenic purpura. *J Thromb Haemost.* 2020;18(10):2496-2502. doi:10.1111/jth.15010.
127. Michel M, Chanet V, Dechartres A, et al. The spectrum of Evans syndrome in adults: new insight into the disease based on the analysis of 68 cases. *Blood.* 2009 Oct 8;114(15):3167-72. doi: 10.1182/blood-2009-04-215368.
128. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Pediatric Hodgkin Lymphoma 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2022.
129. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Hematopoietic Cell Transplantation 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed June 2022.
130. Khouri IF, Saliba RM, Giralt SA, et al. Nonablative allogeneic hematopoietic transplantation as adoptive immunotherapy for indolent lymphoma: low incidence of toxicity,

acute graft-versus-host disease, and treatment-related mortality. *Blood*. 2001 Dec 15;98(13):3595-9. doi: 10.1182/blood.v98.13.3595.

131. National Government Services, Inc. Local Coverage Article: Billing and Coding: Rituximab biosimilars and Rituximab and hyaluronidase human (Rituxan Hycela™) (A52452). Centers for Medicare & Medicaid Services, Inc. Updated on 09/24/2021 with effective date of 10/01/2021. Accessed June 2022.
132. Wisconsin Physicians Service Insurance Corp. Local Coverage Article: Billing and Coding: Chemotherapy Agents for Non-Oncologic Conditions (A55639). Centers for Medicare & Medicaid Services, Inc. Updated on 11/16/2021 with effective date 11/25/2021. Accessed June 2022.
133. Palmetto GBA. Local Coverage Article: Billing and Coding: Rituximab (A56380). Centers for Medicare & Medicaid Services, Inc. Updated on 11/03/2021 with effective date of 10/01/2021. Accessed June 2022.
134. CGS Administrators, LLC. Local Coverage Article: Billing and Coding: Immune Thrombocytopenia (ITP) Therapy (A57160). Centers for Medicare & Medicaid Services, Inc. Updated on 02/23/2022 with effective date 03/03/2022. Accessed June 2022.
135. CGS Administrators, LLC. Local Coverage Article: Billing and Coding: Off-label Use of Rituximab and Rituximab Biosimilars (A58582). Centers for Medicare & Medicaid Services, Inc. Updated on 10/18/2021 with effective date 10/01/2021. Accessed June 2022.

VIII. References (ENHANCED)

- 1e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Central Nervous System Cancers. Version 1.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2022.
- 2e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Hodgkin Lymphoma. Version 2.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2022.
- 3e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma. Version 3.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2022.

- 4e. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Waldenström Macroglobulinemia/Lymphoplasmacytic Lymphoma. Version 3.2022. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed June 2022.
- 5e. Maury S, Chevret S, Thomas X, et al. Rituximab in B-Lineage Adult Acute Lymphoblastic Leukemia. *N Engl J Med* 2016; 375:1044-1053.
- 6e. Kantarjian H, Stein A, Gökbuget N, et al. Blinatumomab versus Chemotherapy for Advanced Acute Lymphoblastic Leukemia. *N Engl J Med* 2017; 376:836-847.
- 7e. Kantarjian HM, DeAngelo DJ, Stelljes M, et al. Inotuzumab Ozogamicin versus Standard Therapy for Acute Lymphoblastic Leukemia. *N Engl J Med*. 2016 Aug 25;375(8):740-53.
- 8e. Maude S, Laetsch T, Buechner J, et al. Tisagenlecleucel in Children and Young Adults with B-Cell Lymphoblastic Leukemia. *N Engl J Med* 2018; 378:439-448.
- 9e. Chamberlain MC, Johnston SK, Van Horn A, Glantz MJ. Recurrent lymphomatous meningitis treated with intra-CSF rituximab and liposomal ara-C. *J Neurooncol*. 2009 Feb;91(3):271-7.
- 10e. Rubenstein JL, Li J, Chen L, et al. Multicenter phase 1 trial of intraventricular immunochemotherapy in recurrent CNS lymphoma. *Blood*. 2013;121(5):745–751.
- 11e. Grossman SA, Finkelstein DM, Ruckdeschel JC, et al. Randomized prospective comparison of intraventricular methotrexate and thiotepa in patients with previously untreated neoplastic meningitis. Eastern Cooperative Oncology Group. *J Clin Oncol*. 1993 Mar;11(3):561-9.
- 12e. Glantz MJ, Jaeckle KA, Chamberlain MC, et al. A randomized controlled trial comparing intrathecal sustained-release cytarabine (DepoCyt) to intrathecal methotrexate in patients with neoplastic meningitis from solid tumors. *Clin Cancer Res*. 1999 Nov;5(11):3394-402.
- 13e. Chamberlain MC, Johnston SK. High-dose methotrexate and rituximab with deferred radiotherapy for newly diagnosed primary B-cell CNS lymphoma. *Neuro Oncol*. 2010;12(7):736–744.
- 14e. Glass J, Won M, Schultz CJ, et al. Phase I and II Study of Induction Chemotherapy With Methotrexate, Rituximab, and Temozolomide, Followed By Whole-Brain Radiotherapy and Postirradiation Temozolomide for Primary CNS Lymphoma: NRG Oncology RTOG 0227. *J Clin Oncol*. 2016;34(14):1620–1625.
- 15e. Gregory G, Arumugaswamy A, Leung T, et al. Rituximab is associated with improved survival for aggressive B cell CNS lymphoma. *Neuro Oncol*. 2013;15(8):1068–1073.

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

- 16e. Holdhoff M, Ambady P, Abdelaziz A, et al. High-dose methotrexate with or without rituximab in newly diagnosed primary CNS lymphoma. *Neurology*. 2014;83(3):235–239.
- 17e. Bromberg JEC, Issa S, Bakunina K, et al. Rituximab in patients with primary CNS lymphoma (HOVON 105/ALLG NHL 24): a randomised, open-label, phase 3 intergroup study. *Lancet Oncol*. 2019 Feb;20(2):216-228.
- 18e. Plotkin SR, Betensky RA, Hochberg FH, et al. Treatment of relapsed central nervous system lymphoma with high-dose methotrexate. *Clin Cancer Res*. 2004 Sep 1;10(17):5643-6.
- 19e. Enting RH, Demopoulos A, DeAngelis LM, Abrey LE. Salvage therapy for primary CNS lymphoma with a combination of rituximab and temozolomide. *Neurology*. 2004 Sep 14;63(5):901-3.
- 20e. Nayak L, Abrey LE, Drappatz J, et al. Multicenter phase II study of rituximab and temozolomide in recurrent primary central nervous system lymphoma. *Leuk Lymphoma*. 2013;54(1):58–61.
- 21e. Batchelor TT, Grossman SA, Mikkelsen T, Ye X, Desideri S, Lesser GJ. Rituximab monotherapy for patients with recurrent primary CNS lymphoma. *Neurology*. 2011;76(10):929–930.
- 22e. Rubenstein JL, Geng H, Fraser EJ, et al. Phase 1 investigation of lenalidomide/rituximab plus outcomes of lenalidomide maintenance in relapsed CNS lymphoma. *Blood Adv*. ;2(13):1595–1607.
- 23e. Grommes C, Pastore A, Palaskas N, et al. Ibrutinib Unmasks Critical Role of Bruton Tyrosine Kinase in Primary CNS Lymphoma. *Cancer Discov*. 2017;7(9):1018–1029.
- 24e. Makino K, Nakamura H, Hide T, Kuratsu J, et al. Salvage treatment with temozolomide in refractory or relapsed primary central nervous system lymphoma and assessment of the MGMT status. *J Neurooncol*. 2012 Jan;106(1):155-60.
- 25e. Fischer L, Thiel E, Klasen HA, et al. Prospective trial on topotecan salvage therapy in primary CNS lymphoma. *Ann Oncol*. 2006 Jul;17(7):1141-5.
- 26e. Raizer JJ, Rademaker A, Evens AM, et al. Pemetrexed in the treatment of relapsed/refractory primary central nervous system lymphoma. *Cancer*. 2012 Aug 1;118(15):3743-8.
- 27e. Tun HW, Johnston PB, DeAngelis LM, et al. Phase 1 study of pomalidomide and dexamethasone for relapsed/refractory primary CNS or vitreoretinal lymphoma. *Blood*. 2018;132(21):2240–2248.
- 28e. Ekstrand BC, Lucas JB, Horwitz SM, et al. Rituximab in lymphocyte-predominant Hodgkin disease: results of a phase 2 trial. *Blood*. 2003 Jun 1;101(11):4285-9.
- 29e. Eichenauer DA, Fuchs M, Pluetschow A, et al. Phase 2 study of rituximab in newly diagnosed stage IA nodular lymphocyte-predominant Hodgkin lymphoma: a report from the German Hodgkin Study Group. *Blood*. 2011 Oct 20;118(16):4363-5.

- 30e. Schulz H, Rehwald U, Morschhauser F, et al. Rituximab in relapsed lymphocyte-predominant Hodgkin lymphoma: long-term results of a phase 2 trial by the German Hodgkin Lymphoma Study Group (GHSG). *Blood*. 2008 Jan 1;111(1):109-11.
- 31e. Burger JA, Tedeschi A, Barr PM, et al. Ibrutinib as Initial Therapy for Patients with Chronic Lymphocytic Leukemia. *N Engl J Med*. 2015;373(25):2425–2437.
- 32e. Woyach JA, Ruppert AS, Heerema NA, et al. Ibrutinib Regimens versus Chemoimmunotherapy in Older Patients with Untreated CLL. *N Engl J Med*. 2018 Dec 27;379(26):2517-2528.
- 33e. Fischer K, Cramer P, Busch R, et al. Bendamustine in combination with rituximab for previously untreated patients with chronic lymphocytic leukemia: a multicenter phase II trial of the German Chronic Lymphocytic Leukemia Study Group. *J Clin Oncol*. 2012 Sep 10;30(26):3209-16.
- 34e. Michallet AS, Aktan M, Hiddemann W, et al. Rituximab plus bendamustine or chlorambucil for chronic lymphocytic leukemia: primary analysis of the randomized, open-label MABLE study. *Haematologica*. 2018;103(4):698–706.
- 35e. Hillmen P, Robak T, Janssens A, et al. Chlorambucil plus ofatumumab versus chlorambucil alone in previously untreated patients with chronic lymphocytic leukaemia (COMPLEMENT 1): a randomised, multicentre, open-label phase 3 trial. *Lancet*. 2015 May 9;385(9980):1873-83.
- 36e. Goede V, Fischer K, Busch R, et al. Obinutuzumab plus chlorambucil in patients with CLL and coexisting conditions. *N Engl J Med*. 2014 Mar 20;370(12):1101-10.
- 37e. Shanafelt TD, Wang V, Kay NE, et al. A Randomized Phase III Study of Ibrutinib (PCI-32765)-Based Therapy Vs. Standard Fludarabine, Cyclophosphamide, and Rituximab (FCR) Chemoimmunotherapy in Untreated Younger Patients with Chronic Lymphocytic Leukemia (CLL): A Trial of the ECOG-ACRIN Cancer Research Group (E1912). *Blood*. 2018;132:LBA-4.
- 38e. Eichhorst B, Fink AM, Bahlo J, et al. First-line chemoimmunotherapy with bendamustine and rituximab versus fludarabine, cyclophosphamide, and rituximab in patients with advanced chronic lymphocytic leukaemia (CLL10): an international, open-label, randomised, phase 3, non-inferiority trial. *Lancet Oncol*. 2016 Jul;17(7):928-942.
- 39e. Woyach JA, Ruppert AS, Heerema NA, et al. Chemoimmunotherapy with fludarabine and rituximab produces extended overall survival and progression-free survival in chronic lymphocytic leukemia: long-term follow-up of CALGB study 9712. *J Clin Oncol*. 2011;29(10):1349–1355.
- 40e. Flinn IW, Panayiotidis P, Afanasyev B, et al. A phase 2, multicenter study investigating ofatumumab and bendamustine combination in patients with untreated or relapsed CLL. *Am J Hematol*. 2016 Sep;91(9):900-6.
- 41e. Sharman JP, Yimer HA, Boxer M, et al. Results of a phase II multicenter study of obinutuzumab plus bendamustine in pts with previously untreated chronic lymphocytic leukemia (CLL). *J Clin Oncol*. 2017;35(15_suppl):7523-7523.

- 42e. Ahn IE, Farooqui MZH, Tian X, et al. Depth and durability of response to ibrutinib in CLL: 5-year follow-up of a phase 2 study. *Blood*. 2018 May 24;131(21):2357-2366.
- 43e. Hillmen P, Skotnicki AB, Robak T, et al. Alemtuzumab compared with chlorambucil as first-line therapy for chronic lymphocytic leukemia. *J Clin Oncol*. 2007 Dec 10;25(35):5616-23.
- 44e. Castro JE, James DF, Sandoval-Sus JD, et al. Rituximab in combination with high-dose methylprednisolone for the treatment of chronic lymphocytic leukemia [published correction appears in *Leukemia*. 2009 Dec;23(12):2326]. *Leukemia*. 2009;23(10):1779–1789.
- 45e. Byrd JC, Flynn JM, Kipps TJ, et al. Randomized phase 2 study of obinutuzumab monotherapy in symptomatic, previously untreated chronic lymphocytic leukemia. *Blood*. 2016;127(1):79–86.
- 46e. Seymour JF, Kipps TJ, Eichhorst B, et al. Venetoclax–Rituximab in Relapsed or Refractory Chronic Lymphocytic Leukemia. *N Engl J Med* 2018; 378:1107-1120.
- 47e. Byrd JC, Brown JR, O'Brien S, et al. Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med*. 2014;371(3):213–223.
- 48e. Byrd JC, Hillmen P, O'Brien SM, et al. Long-term efficacy and safety with ibrutinib (ibr) in previously treated chronic lymphocytic leukemia (CLL): Up to four years follow-up of the RESONATE study. *J Clin Oncol*. 2017;35(15_suppl):7510-7510.
- 49e. Furman RR, Sharman JP, Coutre SE, et al. Idelalisib and rituximab in relapsed chronic lymphocytic leukemia. *N Engl J Med*. 2014;370(11):997–1007.
- 50e. Flinn IW, Hillmen P, Montillo M, et al. The phase 3 DUO trial: duvelisib vs ofatumumab in relapsed and refractory CLL/SLL. *Blood*. 2018;132(23):2446–2455.
- 51e. Keating MJ, Flinn I, Jain V, et al. Therapeutic role of alemtuzumab (Campath-1H) in patients who have failed fludarabine: results of a large international study. *Blood*. 2002 May 15;99(10):3554-61.
- 52e. Faderl S, Ferrajoli A, Wierda W, O'Brien S, Lerner S, Keating MJ. Alemtuzumab by continuous intravenous infusion followed by subcutaneous injection plus rituximab in the treatment of patients with chronic lymphocytic leukemia recurrence [published correction appears in *Cancer*. 2010 Aug 15;116(16):3982. Dosage error in article text]. *Cancer*. 2010;116(10):2360–2365.
- 53e. Robak T, Warzocha K, Govind Babu K, et al. Ofatumumab plus fludarabine and cyclophosphamide in relapsed chronic lymphocytic leukemia: results from the COMPLEMENT 2 trial. *Leuk Lymphoma*. 2017 May;58(5):1084-1093.
- 54e. Castro JE, Sandoval-Sus JD, Bole J, Rassenti L, Kipps TJ. Rituximab in combination with high-dose methylprednisolone for the treatment of fludarabine refractory high-risk chronic lymphocytic leukemia. *Leukemia*. 2008;22(11):2048–2053.
- 55e. Badoux XC, Keating MJ, Wen S, et al. Phase II study of lenalidomide and rituximab as salvage therapy for patients with relapsed or refractory chronic lymphocytic leukemia. *J Clin Oncol*. 2012;31(5):584–591.

- 56e. Bühler A, Wendtner CM, Kipps TJ, et al. Lenalidomide treatment and prognostic markers in relapsed or refractory chronic lymphocytic leukemia: data from the prospective, multicenter phase-II CLL-009 trial. *Blood Cancer J.* 2016;6(3):e404. Published 2016 Mar 11.
- 57e. Byrd JC, Wierda WG, Schuh A, et al. Acalabrutinib Monotherapy in Patients with Relapsed/Refractory Chronic Lymphocytic Leukemia: Updated Results from the Phase 1/2 ACE-CL-001 Study. *Blood.* 2017;130:498.
- 58e. Gopal AK, Davies AJ, Flinn IW, et al. Idelalisib Monotherapy and Durable Responses in Patients with Relapsed or Refractory Small Lymphocytic Lymphoma (SLL). *Blood.* 2015;126:2743.
- 59e. Cartron G, de Guibert S, Dillhuydy MS, et al. Obinutuzumab (GA101) in relapsed/refractory chronic lymphocytic leukemia: final data from the phase 1/2 GAUGUIN study. *Blood.* 2014: 2196-2202.
- 60e. Österborg A, Jewell RC, Padmanabhan-Iyer S, et al. Ofatumumab monotherapy in fludarabine-refractory chronic lymphocytic leukemia: final results from a pivotal study. *Haematologica.* 2015;100(8):e311–e314.
- 61e. Lamanna N, Kalaycio M, Maslak P, et al. Pentostatin, cyclophosphamide, and rituximab is an active, well-tolerated regimen for patients with previously treated chronic lymphocytic leukemia. *J Clin Oncol.* 2006 Apr 1;24(10):1575-81.
- 62e. Jones JA, Mato AR, Wierda WG, et al. Venetoclax for chronic lymphocytic leukaemia progressing after ibrutinib: an interim analysis of a multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2017;19(1):65–75.
- 63e. Zelenetz AD, Barrientos JC, Brown JR, et al. Idelalisib or placebo in combination with bendamustine and rituximab in patients with relapsed or refractory chronic lymphocytic leukaemia: interim results from a phase 3, randomised, double-blind, placebo-controlled trial. *Lancet Oncol.* 2017;18(3):297–311.
- 64e. Chanan-Khan A, Cramer P, Demirkan F, et al. Ibrutinib combined with bendamustine and rituximab compared with placebo, bendamustine, and rituximab for previously treated chronic lymphocytic leukaemia or small lymphocytic lymphoma (HELIOS): a randomised, double-blind, phase 3 study. *Lancet Oncol.* 2016 Feb;17(2):200-211.
- 65e. O'Brien S, Jones JA2, Coutre SE, et al. Ibrutinib for patients with relapsed or refractory chronic lymphocytic leukaemia with 17p deletion (RESONATE-17): a phase 2, open-label, multicentre study. *Lancet Oncol.* 2016 Oct;17(10):1409-1418.
- 66e. Brown JR, Hillmen P, O'Brien S, et al. Extended follow-up and impact of high-risk prognostic factors from the phase 3 RESONATE study in patients with previously treated CLL/SLL. *Leukemia.* 2017;32(1):83–91.
- 67e. Sharman JP, Coutre SE, Furman RR, et al. Second Interim Analysis of a Phase 3 Study of Idelalisib (ZYDELIG®) Plus Rituximab (R) for Relapsed Chronic Lymphocytic Leukemia (CLL): Efficacy Analysis in Patient Subpopulations with Del(17p) and Other Adverse Prognostic Factors. *Blood.* 2014;124:330.

- 68e. Stilgenbauer S, Eichhorst B, Schetelig J, et al. Venetoclax in relapsed or refractory chronic lymphocytic leukaemia with 17p deletion: a multicentre, open-label, phase 2 study. *Lancet Oncol.* 2016 Jun;17(6):768-778.
- 69e. Stilgenbauer S, Zenz T, Winkler D, et al. Subcutaneous alemtuzumab in fludarabine-refractory chronic lymphocytic leukemia: clinical results and prognostic marker analyses from the CLL2H study of the German Chronic Lymphocytic Leukemia Study Group. *J Clin Oncol.* 2009 Aug 20;27(24):3994-4001.
- 70e. Bowen DA, Call TG, Jenkins GD, et al. Methylprednisolone-rituximab is an effective salvage therapy for patients with relapsed chronic lymphocytic leukemia including those with unfavorable cytogenetic features. *Leuk Lymphoma.* 2007 Dec;48(12):2412-7.
- 71e. Brown JR, Byrd JC, Coutre SE, et al. Idelalisib, an inhibitor of phosphatidylinositol 3-kinase p110δ, for relapsed/refractory chronic lymphocytic leukemia. *Blood.* 2014;123(22):3390–3397.
- 72e. Wierda WG, Kipps TJ, Mayer J, et al. Ofatumumab as single-agent CD20 immunotherapy in fludarabine-refractory chronic lymphocytic leukemia [published correction appears in *J Clin Oncol.* 2010 Aug 1;28(22):3670]. *J Clin Oncol.* 2010;28(10):1749–1755.
- 73e. van Oers MH, Kuliczowski K, Smolej L, et al. Ofatumumab maintenance versus observation in relapsed chronic lymphocytic leukaemia (PROLONG): an open-label, multicentre, randomised phase 3 study. *Lancet Oncol.* 2015 Oct;16(13):1370-9.
- 74e. Langerbeins P, Busch R, Anheier N, et al. Poor efficacy and tolerability of R-CHOP in relapsed/refractory chronic lymphocytic leukemia and Richter transformation. *Am J Hematol.* 2014 Dec;89(12):E239-43.
- 75e. Rogers KA, Huang Y, Ruppert AS, et al. A single-institution retrospective cohort study of first-line R-EPOCH chemoimmunotherapy for Richter syndrome demonstrating complex chronic lymphocytic leukaemia karyotype as an adverse prognostic factor. *Br J Haematol.* 2018 Jan;180(2):259-266.
- 76e. Tsimberidou AM, Kantarjian HM, Cortes J, et al. Fractionated cyclophosphamide, vincristine, liposomal daunorubicin, and dexamethasone plus rituximab and granulocyte-macrophage-colony stimulating factor (GM-CSF) alternating with methotrexate and cytarabine plus rituximab and GM-CSF in patients with Richter syndrome or fludarabine-refractory chronic lymphocytic leukemia. *Cancer.* 2003 Apr 1;97(7):1711-20.
- 77e. Tsimberidou AM, Wierda WG, Plunkett W, et al. Phase I-II study of oxaliplatin, fludarabine, cytarabine, and rituximab combination therapy in patients with Richter's syndrome or fludarabine-refractory chronic lymphocytic leukemia. *J Clin Oncol.* 2008 Jan 10;26(2):196-203.
- 78e. Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet.* 2013 Apr 6;381(9873):1203-10.

- 79e. Dimopoulos MA, García-Sanz R, Gavriatopoulou M, et al. Primary therapy of Waldenstrom macroglobulinemia (WM) with weekly bortezomib, low-dose dexamethasone, and rituximab (BDR): long-term results of a phase 2 study of the European Myeloma Network (EMN). *Blood*. 2013 Nov 7;122(19):3276-82.
- 80e. Dimopoulos MA, Anagnostopoulos A, Kyrtsolis MC, et al. Primary treatment of Waldenström macroglobulinemia with dexamethasone, rituximab, and cyclophosphamide. *J Clin Oncol*. 2007 Aug 1;25(22):3344-9.
- 81e. Treon SP, Hanzis C, Tripsas C, et al. Bendamustine therapy in patients with relapsed or refractory Waldenström's macroglobulinemia. *Clin Lymphoma Myeloma Leuk*. 2011 Feb;11(1):133-5.
- 82e. Furman RR, Eradat H, Switzky JC, et al. A Phase II Trial of Ofatumumab In Subjects with Waldenstrom's Macroglobulinemia. *Blood*. 2010;116:1795.
- 83e. Paludo J, Abeykoon JP, Shreders A, et al. Bendamustine and rituximab (BR) versus dexamethasone, rituximab, and cyclophosphamide (DRC) in patients with Waldenström macroglobulinemia. *Ann Hematol*. 2018 Aug;97(8):1417-1425.
- 84e. Treon SP, Hunter ZR, Matous J, et al. Multicenter clinical trial of bortezomib in relapsed/refractory Waldenstrom's macroglobulinemia: results of WMCTG Trial 03-248. *Clin Cancer Res*. 2007 Jun 1;13(11):3320-5.
- 85e. Ghobrial IM, Witzig TE, Gertz M, et al. Long-term results of the phase II trial of the oral mTOR inhibitor everolimus (RAD001) in relapsed or refractory Waldenstrom Macroglobulinemia. *Am J Hematol*. 2014 Mar;89(3):237-42.
- 86e. Barta SK, Xue X, Wang D, et al. Treatment factors affecting outcomes in HIV-associated non-Hodgkin lymphomas: a pooled analysis of 1546 patients. *Blood*. 2013;122(19):3251–3262.
- 87e. Kaplan LD, Lee JY, Ambinder RF, et al. Rituximab does not improve clinical outcome in a randomized phase 3 trial of CHOP with or without rituximab in patients with HIV-associated non-Hodgkin lymphoma: AIDS-Malignancies Consortium Trial 010. *Blood*. 2005;106(5):1538–1543.
- 88e. Boué F, Gabarre J, Gisselbrecht C, et al. Phase II trial of CHOP plus rituximab in patients with HIV-associated non-Hodgkin's lymphoma. *J Clin Oncol*. 2006 Sep 1;24(25):4123-8.
- 89e. Barnes JA, Lacasce AS, Feng Y, et al. Evaluation of the addition of rituximab to CODOX-M/IVAC for Burkitt's lymphoma: a retrospective analysis. *Ann Oncol*. 2011 Aug;22(8):1859-64.
- 90e. Sparano JA, Lee JY, Kaplan LD, et al. Rituximab plus concurrent infusional EPOCH chemotherapy is highly effective in HIV-associated B-cell non-Hodgkin lymphoma. *Blood*. 2010;115(15):3008–3016.
- 91e. Barta SK, Lee JY, Kaplan LD, Noy A, Sparano JA. Pooled analysis of AIDS malignancy consortium trials evaluating rituximab plus CHOP or infusional EPOCH chemotherapy in HIV-associated non-Hodgkin lymphoma. *Cancer*. 2012;118(16):3977–3983.

- 92e. Cortes J, Thomas D, Rios A, et al. Hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone and highly active antiretroviral therapy for patients with acquired immunodeficiency syndrome-related Burkitt lymphoma/leukemia. *Cancer*. 2002 Mar 1;94(5):1492-9.
- 93e. Ribrag V, Koscielny S, Bosq J, et al. Rituximab and dose-dense chemotherapy for adults with Burkitt's lymphoma: a randomised, controlled, open-label, phase 3 trial. *Lancet*. 2016 Jun 11;387(10036):2402-11.
- 94e. Maruyama D, Watanabe T, Maeshima AM, et al. Modified cyclophosphamide, vincristine, doxorubicin, and methotrexate (CODOX-M)/ifosfamide, etoposide, and cytarabine (IVAC) therapy with or without rituximab in Japanese adult patients with Burkitt lymphoma (BL) and B cell lymphoma, unclassifiable, with features intermediate between diffuse large B cell lymphoma and BL. *Int J Hematol*. 2010 Dec;92(5):732-43.
- 95e. Thomas DA, Faderl S, O'Brien S, et al. Chemoimmunotherapy with hyper-CVAD plus rituximab for the treatment of adult Burkitt and Burkitt-type lymphoma or acute lymphoblastic leukemia. *Cancer*. 2006 Apr 1;106(7):1569-80.
- 96e. Hoelzer D, Walewski J, Döhner H, et al. Improved outcome of adult Burkitt lymphoma/leukemia with rituximab and chemotherapy: report of a large prospective multicenter trial. *Blood*. 2014;124(26):3870–3879.
- 97e. Gérard L, Bérezné A, Galicier L, et al. Prospective study of rituximab in chemotherapy-dependent human immunodeficiency virus associated multicentric Castleman's disease: ANRS 117 CastlemaB Trial. *J Clin Oncol*. 2007 Aug 1;25(22):3350-6.
- 98e. Hoffmann C, Schmid H, Müller M, et al. Improved outcome with rituximab in patients with HIV-associated multicentric Castleman disease. *Blood*. 2011 Sep 29;118(13):3499-503.
- 99e. Petrich AM, Gandhi M, Jovanovic B, et al. Impact of induction regimen and stem cell transplantation on outcomes in double-hit lymphoma: a multicenter retrospective analysis. *Blood*. 2014 Oct 9;124(15):2354-61.
- 100e. Howlett C, Snedecor SJ, Landsburg DJ, et al. Front-line, dose-escalated immunochemotherapy is associated with a significant progression-free survival advantage in patients with double-hit lymphomas: a systematic review and meta-analysis. *Br J Haematol*. 2015 Aug;170(4):504-14.
- 101e. Dunleavy K, Fanale MA, Abramson JS, et al. Dose-adjusted EPOCH-R (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin, and rituximab) in untreated aggressive diffuse large B-cell lymphoma with MYC rearrangement: a prospective, multicentre, single-arm phase 2 study. *Lancet Haematol*. 2018 Dec;5(12):e609-e617.
- 102e. Gisselbrecht C, Schmitz N, Mounier N, et al. Rituximab maintenance therapy after autologous stem-cell transplantation in patients with relapsed CD20(+) diffuse large B-cell lymphoma: final analysis of the collaborative trial in relapsed aggressive lymphoma. *J Clin Oncol*. 2012;30(36):4462–4469. doi:10.1200/JCO.2012.41.9416

- 103e. Ohmachi K, Niitsu N, Uchida T, et al. Multicenter phase II study of bendamustine plus rituximab in patients with relapsed or refractory diffuse large B-cell lymphoma. *J Clin Oncol*. 2013 Jun 10;31(17):2103-9.
- 104e. Jacobsen ED, Sharman JP, Oki Y, et al. Brentuximab vedotin demonstrates objective responses in a phase 2 study of relapsed/refractory DLBCL with variable CD30 expression. *Blood*. 2015 Feb 26;125(9):1394-402.
- 105e. Wilson WH, Young RM, Schmitz R, et al. Targeting B cell receptor signaling with ibrutinib in diffuse large B cell lymphoma. *Nat Med*. 2015 Aug;21(8):922-6.
- 106e. van Imhoff GW, McMillan A, Matasar MJ, et al. Ofatumumab Versus Rituximab Salvage Chemoimmunotherapy in Relapsed or Refractory Diffuse Large B-Cell Lymphoma: The ORCHARRD Study. *J Clin Oncol*. 2017 Feb 10;35(5):544-551.
- 107e. Marcus R, Imrie K, Solal-Celigny P, et al. Phase III study of R-CVP compared with cyclophosphamide, vincristine, and prednisone alone in patients with previously untreated advanced follicular lymphoma. *J Clin Oncol*. 2008 Oct 1;26(28):4579-86.
- 108e. Federico M, Luminari S, Dondi A, et al. R-CVP versus R-CHOP versus R-FM for the initial treatment of patients with advanced-stage follicular lymphoma: results of the FOLL05 trial conducted by the Fondazione Italiana Linfomi. *J Clin Oncol*. 2013 Apr 20;31(12):1506-13.
- 109e. Marcus R, Davies A, Ando K, et al. Obinutuzumab for the First-Line Treatment of Follicular Lymphoma. *N Engl J Med* 2017; 377:1331-1344.
- 110e. Rosenbaum CA, Jung SH, Pitcher B, et al. Phase 2 multicentre study of single-agent ofatumumab in previously untreated follicular lymphoma: CALGB 50901 (Alliance). *Br J Haematol*. 2019 Apr;185(1):53-64.
- 111e. Schulz H, Bohlius JF, Trelle S, et al. Immunochemotherapy with rituximab and overall survival in patients with indolent or mantle cell lymphoma: a systematic review and meta-analysis. *J Natl Cancer Inst*. 2007 May 2;99(9):706-14.
- 112e. Morschhauser F, Fowler NH, Feugier P, et al. Rituximab plus Lenalidomide in Advanced Untreated Follicular Lymphoma. *N Engl J Med* 2018; 379:934-947.
- 113e. Sehn LH, Chua N, Mayer J, et al. Obinutuzumab plus bendamustine versus bendamustine monotherapy in patients with rituximab-refractory indolent non-Hodgkin lymphoma (GADOLIN): a randomised, controlled, open-label, multicentre, phase 3 trial. *Lancet Oncol*. 2016 Jun 23. pii: S1470-2045(16)30097-3.
- 114e. Rummel M, Kaiser U, Balsler C, et al. Bendamustine plus rituximab versus fludarabine plus rituximab for patients with relapsed indolent and mantle-cell lymphomas: a multicentre, randomised, open-label, non-inferiority phase 3 trial. *Lancet Oncol*. 2016 Jan;17(1):57-66.
- 115e. Dreyling M, Santoro A, Mollica L, et al. Long-Term Efficacy and Safety from the Copanlisib CHRONOS-1 Study in Patients with Relapsed or Refractory Indolent B-Cell Lymphoma. *Blood*. 2018; 132:1595.

- 116e. Radford J, Davies A, Cartron G, et al. Obinutuzumab (GA101) plus CHOP or FC in relapsed/refractory follicular lymphoma: results of the GAUDI study (BO21000). *Blood*. 2013 Aug 15;122(7):1137-43.
- 117e. Czuczman MS, Fayad L, Delwail V, et al. Ofatumumab monotherapy in rituximab-refractory follicular lymphoma: results from a multicenter study. *Blood*. 2012 Apr 19;119(16):3698-704.
- 118e. Sehn LH, Goy A, Offner FC, et al. Randomized Phase II Trial Comparing Obinutuzumab (GA101) With Rituximab in Patients With Relapsed CD20+ Indolent B-Cell Non-Hodgkin Lymphoma: Final Analysis of the GAUSS Study. *J Clin Oncol*. 2015;33(30):3467–3474.
- 119e. Cheson BD, Chua N, Mayer J, et al. Overall Survival Benefit in Patients With Rituximab-Refractory Indolent Non-Hodgkin Lymphoma Who Received Obinutuzumab Plus Bendamustine Induction and Obinutuzumab Maintenance in the GADOLIN Study. *J Clin Oncol*. 2018 36:22, 2259-2266.
- 120e. Martinelli G, Laszlo D, Ferreri AJ, et al. Clinical activity of rituximab in gastric marginal zone non-Hodgkin's lymphoma resistant to or not eligible for anti-*Helicobacter pylori* therapy. *J Clin Oncol*. 2005 Mar 20;23(9):1979-83.
- 121e. Conconi A, Martinelli G, Thiéblemont C, et al. Clinical activity of rituximab in extranodal marginal zone B-cell lymphoma of MALT type. *Blood*. 2003 Oct 15;102(8):2741-5.
- 122e. Raderer M, Wohrer S, Streubel B, et al. Activity of rituximab plus cyclophosphamide, doxorubicin/mitoxantrone, vincristine and prednisone in patients with relapsed MALT lymphoma. *Oncology*. 2006;70(6):411-7.
- 123e. Salar A, Domingo-Domenech E, Estany C, et al. Combination therapy with rituximab and intravenous or oral fludarabine in the first-line, systemic treatment of patients with extranodal marginal zone B-cell lymphoma of the mucosa-associated lymphoid tissue type. *Cancer*. 2009 Nov 15;115(22):5210-7.
- 124e. Zucca E, Conconi A, Laszlo D, et al. Addition of rituximab to chlorambucil produces superior event-free survival in the treatment of patients with extranodal marginal-zone B-cell lymphoma: 5-year analysis of the IELSG-19 Randomized Study. *J Clin Oncol*. 2013 Feb 10;31(5):565-72.
- 125e. Flinn IW, van der Jagt R, Kahl BS, et al. Randomized trial of bendamustine-rituximab or R-CHOP/R-CVP in first-line treatment of indolent NHL or MCL: the BRIGHT study. *Blood*. 2014;123(19):2944–2952.
- 126e. Salar A, Domingo-Domenech E, Panizo C, et al. Final Results of a Multicenter Phase II Trial with Bendamustine and Rituximab As First Line Treatment for Patients with MALT Lymphoma (MALT-2008–01). *Blood*. 2012;120:3691.
- 127e. Conconi A, Martinelli G, Thiéblemont C, et al. Clinical activity of rituximab in extranodal marginal zone B-cell lymphoma of MALT type. *Blood*. 2003 Oct 15;102(8):2741-5.

- 128e. Kiesewetter B, Neuper O1, Mayerhoefer ME, et al. A pilot phase II study of ofatumumab monotherapy for extranodal marginal zone B-cell lymphoma of the mucosa-associated lymphoid tissue (MALT) lymphoma. *Hematol Oncol*. 2018 Feb;36(1):49-55.
- 129e. Noy A, de Vos S, Thieblemont C, et al. Targeting Bruton tyrosine kinase with ibrutinib in relapsed/refractory marginal zone lymphoma. *Blood*. 2017;129(16):2224–2232.
- 130e. Leonard JP, Trněný M, Izutsu K, et al. AUGMENT: A Phase III Randomized Study of Lenalidomide Plus Rituximab (R2) Vs Rituximab/Placebo in Patients with Relapsed/Refractory Indolent Non-Hodgkin Lymphoma. *Blood*. 2018;132:445.
- 131e. Tsimberidou AM, Catovsky D, Schlette E, et al. Outcomes in patients with splenic marginal zone lymphoma and marginal zone lymphoma treated with rituximab with or without chemotherapy or chemotherapy alone. *Cancer*. 2006 Jul 1;107(1):125-35.
- 132e. Else M, Marín-Niebla A, de la Cruz F, et al. Rituximab, used alone or in combination, is superior to other treatment modalities in splenic marginal zone lymphoma. *Br J Haematol*. 2012 Nov;159(3):322-8.
- 133e. Goodman GR, Burian C, Koziol JA, Saven A. Extended follow-up of patients with hairy cell leukemia after treatment with cladribine. *J Clin Oncol*. 2003 Mar 1;21(5):891-6.
- 134e. Jones J, Andritsos L, Kreitman RJ, et al. Efficacy and Safety of the Bruton Tyrosine Kinase Inhibitor Ibrutinib in Patients with Hairy Cell Leukemia: Stage 1 Results of a Phase 2 Study. *Blood*. 2016;128:1215.
- 135e. Park JH, Lee JO, Stone RM, et al. Acquired Resistance to BRAF Inhibition in Hcl Is Rare and Retreatment with Vemurafenib at Relapse Can Induce High Response Rates: Final Results of a Phase II Trial of Vemurafenib in Relapsed Hcl. *Blood*. 2018;132:392.
- 136e. Tiacchi E, De Carolis L, Zaja F, et al. The Chemotherapy-Free Combination of Vemurafenib and Rituximab Produces Deep and Durable Responses in Relapsed or Refractory Hairy Cell Leukemia (HCL) Patients. *Blood*. 2017;130:409.
- 137e. Kreitman RJ, Dearden C, Zingani PL, et al. Moxetumomab pasudotox in relapsed/refractory hairy cell leukemia. *Leukemia*. 2018; 32(8): 1768–1777.
- 138e. Le Guill S, Thieblemont C, Oberic L, et al. Rituximab after Autologous Stem-Cell Transplantation in Mantle-Cell Lymphoma. *N Engl J Med* 2017; 377:1250-1260
- 139e. Hermine O, Hoster E, Walewski J, et al. Addition of high-dose cytarabine to immunochemotherapy before autologous stem-cell transplantation in patients aged 65 years or younger with mantle cell lymphoma (MCL Younger): a randomised, open-label, phase 3 trial of the European Mantle Cell Lymphoma Network. *Lancet*. 2016 Aug 6;388(10044):565-75.
- 140e. Romaguera JE, Fayad L, Rodriguez MA, et al. High rate of durable remissions after treatment of newly diagnosed aggressive mantle-cell lymphoma with rituximab plus hyper-CVAD alternating with rituximab plus high-dose methotrexate and cytarabine. *J Clin Oncol*. 2005 Oct 1;23(28):7013-23.

- 141e. Lenz G, Dreyling M, Hoster E, et al. Immunochemotherapy with rituximab and cyclophosphamide, doxorubicin, vincristine, and prednisone significantly improves response and time to treatment failure, but not long-term outcome in patients with previously untreated mantle cell lymphoma: results of a prospective randomized trial of the German Low Grade Lymphoma Study Group (GLSG). *J Clin Oncol*. 2005 Mar 20;23(9):1984-92.
- 142e. Schulz H, Bohlius JF, Trelle S, et al. Immunochemotherapy with rituximab and overall survival in patients with indolent or mantle cell lymphoma: a systematic review and meta-analysis. *J Natl Cancer Inst*. 2007 May 2;99(9):706-14.
- 143e. Casulo C, Iannotta A, Walkley J, et al. Ofatumumab-Bendamustine As First Line Treatment for Elderly Patients with Mantle Cell Lymphoma: A Phase II Risk Adapted Design with Comprehensive Geriatric Assessment. *Blood*. 2014;124:1751.
- 144e. Cavalli F, Rooney B, Pei L, et al. Randomized phase 3 study of rituximab, cyclophosphamide, doxorubicin, and prednisone plus vincristine (R-CHOP) or bortezomib (VR-CAP) in newly diagnosed mantle cell lymphoma (MCL) patients (pts) ineligible for bone marrow transplantation (BMT). *J Clin Oncol*. 2014;32(15_suppl):8500-8500.
- 145e. Wang M, Rule S, Zinzani PL, et al. Acalabrutinib in relapsed or refractory mantle cell lymphoma (ACE-LY-004): a single-arm, multicentre, phase 2 trial. *Lancet*. 2018 Feb 17;391(10121):659-667.
- 146e. Rule S, Jurczak W, Jerkeman M, et al. Ibrutinib versus temsirolimus: 3-year follow-up of patients with previously treated mantle cell lymphoma from the phase 3, international, randomized, open-label RAY study. *Leukemia*. 2018;32(8):1799–1803.
- 147e. Jain P, Romaguera J, Srour SA, et al. Four-year follow-up of a single arm, phase II clinical trial of ibrutinib with rituximab (IR) in patients with relapsed/refractory mantle cell lymphoma (MCL). *Br J Haematol*. 2018 Aug;182(3):404-411.
- 148e. Goy A, Sinha R, Williams ME, et al. Single-agent lenalidomide in patients with mantle-cell lymphoma who relapsed or progressed after or were refractory to bortezomib: phase II MCL-001 (EMERGE) study. *J Clin Oncol*. 2013;31(29):3688–3695.
- 149e. Wang M, Fayad L, Wagner-Bartak N, et al. Lenalidomide in combination with rituximab for patients with relapsed or refractory mantle-cell lymphoma: a phase 1/2 clinical trial. *Lancet Oncol*. 2012 Jul;13(7):716-23.
- 150e. Davids MS, Roberts AW, Seymour JF, et al. Phase I First-in-Human Study of Venetoclax in Patients With Relapsed or Refractory Non-Hodgkin Lymphoma. *J Clin Oncol*. 2017;35(8):826–833.
- 151e. Goy A, Bernstein SH, Kahl BS, et al. Bortezomib in patients with relapsed or refractory mantle cell lymphoma: updated time-to-event analyses of the multicenter phase 2 PINNACLE study. *Ann Oncol*. 2009;20(3):520–525. doi:10.1093/annonc/mdn656
- 152e. Baiocchi RA, Alinari L, Lustberg ME, et al. Phase 2 trial of rituximab and bortezomib in patients with relapsed or refractory mantle cell and follicular lymphoma. *Cancer*. 2011;117(11):2442–2451. doi:10.1002/cncr.25792

- 153e. Furtado M, Dyer MJ, Johnson R, et al. Ofatumumab monotherapy in relapsed/refractory mantle cell lymphoma—a phase II trial. *Br J Haematol*. 2014 May;165(4):575-8.
- 154e. Evens AM, David KA, Helenowski I, et al. Multicenter analysis of 80 solid organ transplantation recipients with post-transplantation lymphoproliferative disease: outcomes and prognostic factors in the modern era. *J Clin Oncol*. 2010;28(6):1038–1046.
- 155e. Morales AV, Advani R, Horwitz SM, et al. Indolent primary cutaneous B-cell lymphoma: experience using systemic rituximab. *J Am Acad Dermatol*. 2008 Dec;59(6):953-7.
- 156e. Peñate Y, Hernández-Machín B, Pérez-Méndez LI, et al. Intralesional rituximab in the treatment of indolent primary cutaneous B-cell lymphomas: an epidemiological observational multicentre study. The Spanish Working Group on Cutaneous Lymphoma. *Br J Dermatol*. 2012 Jul;167(1):174-9.
- 157e. Brahmer JR, Lacchetti C, Schneider BJ, et al. Management of Immune-Related Adverse Events in Patients Treated With Immune Checkpoint Inhibitor Therapy: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol*. 2018;36(17):1714–1768.
- 158e. Williams TJ, Benavides DR1, Patrice KA, et al. Association of Autoimmune Encephalitis With Combined Immune Checkpoint Inhibitor Treatment for Metastatic Cancer. *JAMA Neurol*. 2016 Aug 1;73(8):928-33.
- 159e. Sehn LH, Herrera AF, Matasar MJ, et al. Polatuzumab Vedotin (Pola) Plus Bendamustine (B) with Rituximab (R) or Obinutuzumab (G) in Relapsed/Refractory (R/R) Diffuse Large B-Cell Lymphoma (DLBCL): Updated Results of a Phase (Ph) Ib/II Study. *Blood* 2018;132:Abstract 1683.
- 160e. Wang M, Fowler N, Wagner-Bartak N, et al. Oral lenalidomide with rituximab in relapsed or refractory diffuse large cell, follicular and transformed lymphoma: a phase II clinical trial. *Leukemia*. 2013 Sep;27(9):1902-9. doi: 10.1038/leu.2013.95.
- 161e. Fischer K, Al-Sawaf O, Bahlo J, et al. Venetoclax and Obinutuzumab in Patients with CLL and Coexisting Conditions. *N Engl J Med*. 2019;380(23):2225-2236. doi:10.1056/NEJMoa1815281.
- 162e. Sharman JP, Egyed M, Jurczak W, et al. Acalabrutinib with or without obinutuzumab versus chlorambucil and obinutuzumab for treatment-naive chronic lymphocytic leukaemia (ELEVATE TN): a randomised, controlled, phase 3 trial [published correction appears in *Lancet*. 2020 May 30;395.
- 163e. Minard-Colin V, Auperin A, Pillon M, et al. Results of the randomized Intergroup trial Inter-B-NHL Ritux 2010 for children and adolescents with high-risk B-cell non-Hodgkin lymphoma (B-NHL) and mature acute leukemia (B-AL): Evaluation of rituximab (R) efficacy in addition to standard LMB chemotherapy (CT) regimen. *J Clin Oncol*. 2016 May;34(15_suppl):10507-10507.
- 164e. Wieduwilt MJ, Jonas BA, Schiller GJ, et al. A Phase II Study of Pegylated Asparaginase, Cyclophosphamide, Rituximab, and Dasatinib Added to the UCSF 8707 (Linker 4-drug) Regimen with Liposomal Cytarabine CNS Prophylaxis for Adults with Newly Diagnosed

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



Acute Lymphoblastic Leukemia (ALL) or Lymphoblastic Lymphoma (LBL): University of California Hematologic Malignancies Consortium Study (UCHMC) 1401. Blood 2018;132:4018.

165e. Namoglu EC, Hughes ME, Plastaras JP, et al. Management and outcomes of sinus histiocytosis with massive lymphadenopathy (Rosai Dorfman Disease). Leuk Lymphoma. 2020 Apr;61(4):905-911. doi: 10.1080/10428194.2019.1703971.

166e. Goekbuget N, Beck J, Brueggemann M, et al. Moderate Intensive Chemotherapy Including CNS-Prophylaxis with Liposomal Cytarabine Is Feasible and effective in Older Patients with Ph-Negative Acute Lymphoblastic Leukemia (ALL): Results of a Prospective Trial From the German Multicenter Study Group for Adult ALL (GMALL). Blood 2012;120:1493.

167e. Magellan Health, Magellan Rx Management. Rituximab Clinical Literature Review Analysis. Last updated June 2022. Accessed June 2022.

Appendix 1 – Covered Diagnosis Codes

ICD-10	Description
C79.32	Secondary malignant neoplasm of cerebral meninges
C81.00	Nodular lymphocyte predominant Hodgkin lymphoma, unspecified site
C81.01	Nodular lymphocyte predominant Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.02	Nodular lymphocyte predominant Hodgkin lymphoma, intrathoracic lymph nodes
C81.03	Nodular lymphocyte predominant Hodgkin lymphoma, intra-abdominal lymph nodes
C81.04	Nodular lymphocyte predominant Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.05	Nodular lymphocyte predominant Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.06	Nodular lymphocyte predominant Hodgkin lymphoma, intrapelvic lymph nodes
C81.07	Nodular lymphocyte predominant Hodgkin lymphoma, spleen
C81.08	Nodular lymphocyte predominant Hodgkin lymphoma, lymph nodes of multiple sites
C81.09	Nodular lymphocyte predominant Hodgkin lymphoma, extranodal and solid organ sites
C82.00	Follicular lymphoma grade I, unspecified site
C82.01	Follicular lymphoma grade I, lymph nodes of head, face and neck
C82.02	Follicular lymphoma, grade I, intrathoracic lymph nodes
C82.03	Follicular lymphoma grade I, intra-abdominal lymph nodes
C82.04	Follicular lymphoma grade I, lymph nodes of axilla and upper limb
C82.05	Follicular lymphoma grade I, lymph nodes of inguinal regional and lower limb
C82.06	Follicular lymphoma grade I, intrapelvic lymph nodes
C82.07	Follicular lymphoma grade I, spleen
C82.08	Follicular lymphoma grade I, lymph nodes of multiple sites
C82.09	Follicular lymphoma grade I, extranodal and solid organ sites
C82.10	Follicular lymphoma grade II, unspecified site
C82.11	Follicular lymphoma grade II, lymph nodes of head, face and neck

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



C82.12	Follicular lymphoma, grade II, intrathoracic lymph nodes
C82.13	Follicular lymphoma grade II, intra-abdominal lymph nodes
C82.14	Follicular lymphoma grade II, lymph nodes of axilla and upper limb
C82.15	Follicular lymphoma grade II, lymph nodes of inguinal region and lower limb
C82.16	Follicular lymphoma grade II, intrapelvic lymph nodes
C82.17	Follicular lymphoma grade II, spleen
C82.18	Follicular lymphoma grade II, lymph nodes of multiple sites
C82.19	Follicular lymphoma grade II, extranodal and solid organ sites
C82.20	Follicular lymphoma grade III, unspecified, unspecified site
C82.21	Follicular lymphoma grade III, unspecified, lymph nodes of head, face and neck
C82.22	Follicular lymphoma, grade III, unspecified, intrathoracic lymph nodes
C82.23	Follicular lymphoma grade III, unspecified, intra-abdominal lymph nodes
C82.24	Follicular lymphoma grade III, unspecified, lymph nodes of axilla and upper limb
C82.25	Follicular lymphoma grade III, unspecified, lymph nodes of inguinal region and lower limb
C82.26	Follicular lymphoma grade III, unspecified, intrapelvic lymph nodes
C82.27	Follicular lymphoma grade III, unspecified, spleen
C82.28	Follicular lymphoma grade III, unspecified, lymph nodes of multiple sites
C82.29	Follicular lymphoma grade III, unspecified, extranodal and solid organ sites
C82.30	Follicular lymphoma grade IIIa, unspecified site
C82.31	Follicular lymphoma grade IIIa, lymph nodes of head, face and neck
C82.32	Follicular lymphoma, grade IIIa, intrathoracic lymph nodes
C82.33	Follicular lymphoma grade IIIa, intra-abdominal lymph nodes
C82.34	Follicular lymphoma grade IIIa, lymph nodes of axilla and upper limb
C82.35	Follicular lymphoma grade IIIa, lymph nodes of inguinal region and lower limb
C82.36	Follicular lymphoma grade IIIa, intrapelvic lymph nodes
C82.37	Follicular lymphoma grade IIIa, spleen
C82.38	Follicular lymphoma grade IIIa, lymph nodes of multiple sites
C82.39	Follicular lymphoma grade IIIa, extranodal and solid organ sites
C82.40	Follicular lymphoma grade IIIb, unspecified site
C82.41	Follicular lymphoma grade IIIb, lymph nodes of head, face and neck
C82.42	Follicular lymphoma, grade IIIb, intrathoracic lymph nodes
C82.43	Follicular lymphoma grade IIIb, intra-abdominal lymph nodes
C82.44	Follicular lymphoma grade IIIb, lymph nodes of axilla and upper limb
C82.45	Follicular lymphoma grade IIIb, lymph nodes of inguinal region and lower limb
C82.46	Follicular lymphoma grade IIIb, intrapelvic lymph nodes
C82.47	Follicular lymphoma grade IIIb, spleen
C82.48	Follicular lymphoma grade IIIb, lymph nodes of multiple sites

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

C82.49	Follicular lymphoma grade IIIb, extranodal and solid organ sites
C82.50	Diffuse follicle center lymphoma, unspecified site
C82.51	Diffuse follicle center lymphoma, lymph nodes of head, face and neck
C82.52	Diffuse follicle center lymphoma, intrathoracic lymph nodes
C82.53	Diffuse follicle center lymphoma, intra-abdominal lymph nodes
C82.54	Diffuse follicle center lymphoma, lymph nodes of axilla and upper limb
C82.55	Diffuse follicle center lymphoma, lymph nodes of inguinal region and lower limb
C82.56	Diffuse follicle center lymphoma, intrapelvic lymph nodes
C82.57	Diffuse follicle center lymphoma, spleen
C82.58	Diffuse follicle center lymphoma, lymph nodes of multiple sites
C82.59	Diffuse follicle center lymphoma, extranodal and solid organ sites
C82.60	Cutaneous follicle center lymphoma, unspecified site
C82.61	Cutaneous follicle center lymphoma, lymph nodes of head, face and neck
C82.62	Cutaneous follicle center lymphoma, intrathoracic lymph nodes
C82.63	Cutaneous follicle center lymphoma, intra-abdominal lymph nodes
C82.64	Cutaneous follicle center lymphoma, lymph nodes of axilla and upper limb
C82.65	Cutaneous follicle center lymphoma, lymph nodes of inguinal region and lower limb
C82.66	Cutaneous follicle center lymphoma, intrapelvic lymph nodes
C82.67	Cutaneous follicle center lymphoma, spleen
C82.68	Cutaneous follicle center lymphoma, lymph nodes of multiple sites
C82.69	Cutaneous follicle center lymphoma, extranodal and solid organ sites
C82.80	Other types of follicular lymphoma, unspecified site
C82.81	Other types of follicular lymphoma, lymph nodes of head, face and neck
C82.82	Other types of follicular lymphoma, intrathoracic lymph nodes
C82.83	Other types of follicular lymphoma, intra-abdominal lymph nodes
C82.84	Other types of follicular lymphoma, lymph nodes of axilla and upper limb
C82.85	Other types of follicular lymphoma, lymph nodes of inguinal region and lower limb
C82.86	Other types of follicular lymphoma, intrapelvic lymph nodes
C82.87	Other types of follicular lymphoma, spleen
C82.88	Other types of follicular lymphoma, lymph nodes of multiple sites
C82.89	Other types of follicular lymphoma, extranodal and solid organ sites
C82.90	Follicular lymphoma, unspecified, unspecified site
C82.91	Follicular lymphoma, unspecified, lymph nodes of head, face and neck
C82.92	Follicular lymphoma, unspecified, intrathoracic lymph nodes
C82.93	Follicular lymphoma, unspecified, intra-abdominal lymph nodes
C82.94	Follicular lymphoma, unspecified, lymph nodes of axilla and upper limb
C82.95	Follicular lymphoma, unspecified lymph nodes of inguinal region and lower limb

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

C82.96	Follicular lymphoma, unspecified, intrapelvic lymph nodes
C82.97	Follicular lymphoma, unspecified, spleen
C82.98	Follicular lymphoma, unspecified, lymph nodes of multiple sites
C82.99	Follicular lymphoma, unspecified, extranodal and solid organ sites
C83.00	Small cell B-cell lymphoma, unspecified site
C83.01	Small cell B-cell lymphoma, lymph nodes of head, face and neck
C83.02	Small cell B-cell lymphoma, intrathoracic lymph nodes
C83.03	Small cell B-cell lymphoma, intra-abdominal lymph nodes
C83.04	Small cell B-cell lymphoma, lymph nodes of axilla and upper limb
C83.05	Small cell B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.06	Small cell B-cell lymphoma, intrapelvic lymph nodes
C83.07	Small cell B-cell lymphoma, spleen
C83.08	Small cell B-cell lymphoma, lymph nodes of multiple sites
C83.09	Small cell B-cell lymphoma, extranodal and solid organ sites
C83.10	Mantle cell lymphoma, unspecified site
C83.11	Mantle cell lymphoma, lymph nodes of head, face and neck
C83.12	Mantle cell lymphoma, intrathoracic lymph nodes
C83.13	Mantle cell lymphoma, intra-abdominal lymph nodes
C83.14	Mantle cell lymphoma, lymph nodes of axilla and upper limb
C83.15	Mantle cell lymphoma, lymph nodes of inguinal region and lower limb
C83.16	Mantle cell lymphoma, intrapelvic lymph nodes
C83.17	Mantle cell lymphoma, spleen
C83.18	Mantle cell lymphoma, lymph nodes of multiple sites
C83.19	Mantle cell lymphoma, extranodal and solid organ sites
C83.30	Diffuse large B-cell lymphoma unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck
C83.32	Diffuse large B-cell lymphoma intrathoracic lymph nodes
C83.33	Diffuse large B-cell lymphoma intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma extranodal and solid organ sites
C83.50	Lymphoblastic (diffuse) lymphoma, unspecified site
C83.51	Lymphoblastic (diffuse) lymphoma, lymph nodes of head, face, and neck
C83.52	Lymphoblastic (diffuse) lymphoma, intrathoracic lymph nodes

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

C83.53	Lymphoblastic (diffuse) lymphoma, intra-abdominal lymph nodes
C83.54	Lymphoblastic (diffuse) lymphoma, lymph nodes of axilla and upper limb
C83.55	Lymphoblastic (diffuse) lymphoma, lymph nodes of inguinal region and lower limb
C83.56	Lymphoblastic (diffuse) lymphoma, intrapelvic lymph nodes
C83.57	Lymphoblastic (diffuse) lymphoma, spleen
C83.58	Lymphoblastic (diffuse) lymphoma, lymph nodes of multiple sites
C83.59	Lymphoblastic (diffuse) lymphoma, extranodal and solid organ sites
C83.70	Burkitt lymphoma, unspecified site
C83.71	Burkitt lymphoma, lymph nodes of head, face, and neck
C83.72	Burkitt lymphoma, intrathoracic lymph nodes
C83.73	Burkitt lymphoma, intra-abdominal lymph nodes
C83.74	Burkitt lymphoma, lymph nodes of axilla and upper limb
C83.75	Burkitt lymphoma, lymph nodes of inguinal region and lower limb
C83.76	Burkitt lymphoma, intrapelvic lymph nodes
C83.77	Burkitt lymphoma, spleen
C83.78	Burkitt lymphoma, lymph nodes of multiple sites
C83.79	Burkitt lymphoma, extranodal and solid organ sites
C83.80	Other non-follicular lymphoma, unspecified site
C83.81	Other non-follicular lymphoma, lymph nodes of head, face and neck
C83.82	Other non-follicular lymphoma, intrathoracic lymph nodes
C83.83	Other non-follicular lymphoma, intra-abdominal lymph nodes
C83.84	Other non-follicular lymphoma, lymph nodes of axilla and upper limb
C83.85	Other non-follicular lymphoma, lymph nodes of inguinal region and lower limb
C83.86	Other non-follicular lymphoma, intrapelvic lymph nodes
C83.87	Other non-follicular lymphoma, spleen
C83.88	Other non-follicular lymphoma, lymph nodes of multiple sites
C83.89	Other non-follicular lymphoma, extranodal and solid organ sites
C83.90	Non-follicular (diffuse) lymphoma, unspecified site
C83.91	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of head, face, and neck
C83.92	Non-follicular (diffuse) lymphoma, unspecified intrathoracic lymph nodes
C83.93	Non-follicular (diffuse) lymphoma, unspecified intra-abdominal lymph nodes
C83.94	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of axilla and upper limb
C83.95	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of inguinal region and lower limb
C83.96	Non-follicular (diffuse) lymphoma, unspecified intrapelvic lymph nodes
C83.97	Non-follicular (diffuse) lymphoma, unspecified spleen
C83.98	Non-follicular (diffuse) lymphoma, unspecified lymph nodes of multiple sites
C83.99	Non-follicular (diffuse) lymphoma, unspecified extranodal and solid organ sites

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

C85.10	Unspecified B-cell lymphoma, unspecified site
C85.11	Unspecified B-cell lymphoma, lymph nodes of head, face, and neck
C85.12	Unspecified B-cell lymphoma, intrathoracic lymph nodes
C85.13	Unspecified B-cell lymphoma, intra-abdominal lymph nodes
C85.14	Unspecified B-cell lymphoma, lymph nodes of axilla and upper limb
C85.15	Unspecified B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.16	Unspecified B-cell lymphoma, intrapelvic lymph nodes
C85.17	Unspecified B-cell lymphoma, spleen
C85.18	Unspecified B-cell lymphoma, lymph nodes of multiple sites
C85.19	Unspecified B-cell lymphoma, extranodal and solid organ sites
C85.20	Mediastinal (thymic) large B-cell lymphoma, unspecified site
C85.21	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck
C85.22	Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes
C85.23	Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes
C85.24	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb
C85.25	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.26	Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes
C85.27	Mediastinal (thymic) large B-cell lymphoma, spleen
C85.28	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites
C85.29	Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites
C85.80	Other specified types of non-Hodgkin lymphoma, unspecified site
C85.81	Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face and neck
C85.82	Other specified types of non-Hodgkin lymphoma, intrathoracic lymph nodes
C85.83	Other specified types of non-Hodgkin lymphoma, intra-abdominal lymph nodes
C85.84	Other specified types of non-Hodgkin lymphoma, lymph nodes of axilla and upper limb
C85.85	Other specified types of non-Hodgkin lymphoma, lymph nodes of inguinal region of lower limb
C85.86	Other specified types of non-Hodgkin lymphoma, intrapelvic lymph nodes
C85.87	Other specified types of non-Hodgkin lymphoma, spleen
C85.88	Other specified types of non-Hodgkin lymphoma, lymph nodes of multiple sites
C85.89	Other specified types of non-Hodgkin lymphoma, extranodal and solid organ sites
C85.99	Non-Hodgkin lymphoma, unspecified, extranodal and solid organ sites
C88.0	Waldenström macroglobulinemia
C88.4	Extranodal marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue (MALT-lymphoma)
C91.00	Acute lymphoblastic leukemia not having achieved remission
C91.01	Acute lymphoblastic leukemia, in remission
C91.02	Acute lymphoblastic leukemia, in relapse
C91.10	Chronic lymphocytic leukemia of B-cell type not having achieved remission

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

C91.12	Chronic lymphocytic leukemia of B-cell type in relapse
C91.40	Hairy cell leukemia not having achieved remission
C91.42	Hairy cell leukemia, in relapse
D47.Z1	Post-transplant lymphoproliferative disorder (PTLD)
D47.Z2	Other neoplasms of uncertain behavior of lymphoid, hematopoietic and related tissue-Castleman
D59.10	Autoimmune hemolytic anemia, unspecified
D59.11	Warm autoimmune hemolytic anemia
D59.12	Cold autoimmune hemolytic anemia
D59.13	Mixed type autoimmune hemolytic anemia
D59.19	Other autoimmune hemolytic anemia
D69.3	Immune thrombocytopenic purpura
D69.41	Evans Syndrome
D69.42	Congenital and hereditary thrombocytopenia purpura
D69.49	Other primary thrombocytopenia
D89.811	Chronic graft-versus-host disease
D89.812	Acute on chronic graft-versus-host disease
D89.813	Graft-versus-host disease unspecified
G04.81	Other encephalitis and encephalomyelitis
G04.89	Other myelitis
G04.90	Encephalitis and encephalomyelitis, unspecified
G36.0	Neuromyelitis optica [Devic]
G70.0	Myasthenia gravis
G70.00	Myasthenia gravis without (acute) exacerbation
G70.01	Myasthenia gravis with (acute) exacerbation
L10.0	Pemphigus vulgaris
L13.8	Other specified bullous disorders
L13.9	Bullous disorder, unspecified
M05.10	Rheumatoid lung disease with rheumatoid arthritis of unspecified site
M05.111	Rheumatoid lung disease with rheumatoid arthritis of right shoulder
M05.112	Rheumatoid lung disease with rheumatoid arthritis of left shoulder
M05.119	Rheumatoid lung disease with rheumatoid arthritis of unspecified shoulder
M05.121	Rheumatoid lung disease with rheumatoid arthritis of right elbow
M05.122	Rheumatoid lung disease with rheumatoid arthritis of left elbow
M05.129	Rheumatoid lung disease with rheumatoid arthritis of unspecified elbow
M05.131	Rheumatoid lung disease with rheumatoid arthritis of right wrist
M05.132	Rheumatoid lung disease with rheumatoid arthritis of left wrist
M05.139	Rheumatoid lung disease with rheumatoid arthritis of unspecified wrist

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

M05.141	Rheumatoid lung disease with rheumatoid arthritis of right hand
M05.142	Rheumatoid lung disease with rheumatoid arthritis of left hand
M05.149	Rheumatoid lung disease with rheumatoid arthritis of unspecified hand
M05.151	Rheumatoid lung disease with rheumatoid arthritis of right hip
M05.152	Rheumatoid lung disease with rheumatoid arthritis of left hip
M05.159	Rheumatoid lung disease with rheumatoid arthritis of unspecified hip
M05.161	Rheumatoid lung disease with rheumatoid arthritis of right knee
M05.162	Rheumatoid lung disease with rheumatoid arthritis of left knee
M05.169	Rheumatoid lung disease with rheumatoid arthritis of unspecified knee
M05.171	Rheumatoid lung disease with rheumatoid arthritis of right ankle and foot
M05.172	Rheumatoid lung disease with rheumatoid arthritis of left ankle and foot
M05.179	Rheumatoid lung disease with rheumatoid arthritis of unspecified ankle and foot
M05.19	Rheumatoid lung disease with rheumatoid arthritis of multiple sites
M05.20	Rheumatoid vasculitis with rheumatoid arthritis of unspecified site
M05.211	Rheumatoid vasculitis with rheumatoid arthritis of right shoulder
M05.212	Rheumatoid vasculitis with rheumatoid arthritis of left shoulder
M05.219	Rheumatoid vasculitis with rheumatoid arthritis of unspecified shoulder
M05.221	Rheumatoid vasculitis with rheumatoid arthritis of right elbow
M05.222	Rheumatoid vasculitis with rheumatoid arthritis of left elbow
M05.229	Rheumatoid vasculitis with rheumatoid arthritis of unspecified elbow
M05.231	Rheumatoid vasculitis with rheumatoid arthritis of right wrist
M05.232	Rheumatoid vasculitis with rheumatoid arthritis of left wrist
M05.239	Rheumatoid vasculitis with rheumatoid arthritis of unspecified wrist
M05.241	Rheumatoid vasculitis with rheumatoid arthritis of right hand
M05.242	Rheumatoid vasculitis with rheumatoid arthritis of left hand
M05.249	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hand
M05.251	Rheumatoid vasculitis with rheumatoid arthritis of right hip
M05.252	Rheumatoid vasculitis with rheumatoid arthritis of left hip
M05.259	Rheumatoid vasculitis with rheumatoid arthritis of unspecified hip
M05.261	Rheumatoid vasculitis with rheumatoid arthritis of right knee
M05.262	Rheumatoid vasculitis with rheumatoid arthritis of left knee
M05.269	Rheumatoid vasculitis with rheumatoid arthritis of unspecified knee
M05.271	Rheumatoid vasculitis with rheumatoid arthritis of right ankle and foot
M05.272	Rheumatoid vasculitis with rheumatoid arthritis of left ankle and foot
M05.279	Rheumatoid vasculitis with rheumatoid arthritis of unspecified ankle and foot
M05.29	Rheumatoid vasculitis with rheumatoid arthritis of multiple sites
M05.30	Rheumatoid heart disease with rheumatoid arthritis of unspecified site

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

M05.311	Rheumatoid heart disease with rheumatoid arthritis of right shoulder
M05.312	Rheumatoid heart disease with rheumatoid arthritis of left shoulder
M05.319	Rheumatoid heart disease with rheumatoid arthritis of unspecified shoulder
M05.321	Rheumatoid heart disease with rheumatoid arthritis of right elbow
M05.322	Rheumatoid heart disease with rheumatoid arthritis of left elbow
M05.329	Rheumatoid heart disease with rheumatoid arthritis of unspecified elbow
M05.331	Rheumatoid heart disease with rheumatoid arthritis of right wrist
M05.332	Rheumatoid heart disease with rheumatoid arthritis of left wrist
M05.339	Rheumatoid heart disease with rheumatoid arthritis of unspecified wrist
M05.341	Rheumatoid heart disease with rheumatoid arthritis of right hand
M05.342	Rheumatoid heart disease with rheumatoid arthritis of left hand
M05.349	Rheumatoid heart disease with rheumatoid arthritis of unspecified hand
M05.351	Rheumatoid heart disease with rheumatoid arthritis of right hip
M05.352	Rheumatoid heart disease with rheumatoid arthritis of left hip
M05.359	Rheumatoid heart disease with rheumatoid arthritis of unspecified hip
M05.361	Rheumatoid heart disease with rheumatoid arthritis of right knee
M05.362	Rheumatoid heart disease with rheumatoid arthritis of left knee
M05.369	Rheumatoid heart disease with rheumatoid arthritis of unspecified knee
M05.371	Rheumatoid heart disease with rheumatoid arthritis of right ankle and foot
M05.372	Rheumatoid heart disease with rheumatoid arthritis of left ankle and foot
M05.379	Rheumatoid heart disease with rheumatoid arthritis of unspecified ankle and foot
M05.39	Rheumatoid heart disease with rheumatoid arthritis of multiple sites
M05.40	Rheumatoid myopathy with rheumatoid arthritis of unspecified site
M05.411	Rheumatoid myopathy with rheumatoid arthritis of right shoulder
M05.412	Rheumatoid myopathy with rheumatoid arthritis of left shoulder
M05.419	Rheumatoid myopathy with rheumatoid arthritis of unspecified shoulder
M05.421	Rheumatoid myopathy with rheumatoid arthritis of right elbow
M05.422	Rheumatoid myopathy with rheumatoid arthritis of left elbow
M05.429	Rheumatoid myopathy with rheumatoid arthritis of unspecified elbow
M05.431	Rheumatoid myopathy with rheumatoid arthritis of right wrist
M05.432	Rheumatoid myopathy with rheumatoid arthritis of left wrist
M05.439	Rheumatoid myopathy with rheumatoid arthritis of unspecified wrist
M05.441	Rheumatoid myopathy with rheumatoid arthritis of right hand
M05.442	Rheumatoid myopathy with rheumatoid arthritis of left hand
M05.449	Rheumatoid myopathy with rheumatoid arthritis of unspecified hand
M05.451	Rheumatoid myopathy with rheumatoid arthritis of right hip
M05.452	Rheumatoid myopathy with rheumatoid arthritis of left hip

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

M05.459	Rheumatoid myopathy with rheumatoid arthritis of unspecified hip
M05.461	Rheumatoid myopathy with rheumatoid arthritis of right knee
M05.462	Rheumatoid myopathy with rheumatoid arthritis of left knee
M05.469	Rheumatoid myopathy with rheumatoid arthritis of unspecified knee
M05.471	Rheumatoid myopathy with rheumatoid arthritis of right ankle and foot
M05.472	Rheumatoid myopathy with rheumatoid arthritis of left ankle and foot
M05.479	Rheumatoid myopathy with rheumatoid arthritis of unspecified ankle and foot
M05.49	Rheumatoid myopathy with rheumatoid arthritis of multiple sites
M05.50	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified site
M05.511	Rheumatoid polyneuropathy with rheumatoid arthritis of right shoulder
M05.512	Rheumatoid polyneuropathy with rheumatoid arthritis of left shoulder
M05.519	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified shoulder
M05.521	Rheumatoid polyneuropathy with rheumatoid arthritis of right elbow
M05.522	Rheumatoid polyneuropathy with rheumatoid arthritis of left elbow
M05.529	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified elbow
M05.531	Rheumatoid polyneuropathy with rheumatoid arthritis of right wrist
M05.532	Rheumatoid polyneuropathy with rheumatoid arthritis of left wrist
M05.539	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified wrist
M05.541	Rheumatoid polyneuropathy with rheumatoid arthritis of right hand
M05.542	Rheumatoid polyneuropathy with rheumatoid arthritis of left hand
M05.549	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified hand
M05.551	Rheumatoid polyneuropathy with rheumatoid arthritis of right hip
M05.552	Rheumatoid polyneuropathy with rheumatoid arthritis of left hip
M05.559	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified hip
M05.561	Rheumatoid polyneuropathy with rheumatoid arthritis of right knee
M05.562	Rheumatoid polyneuropathy with rheumatoid arthritis of left knee
M05.569	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified knee
M05.571	Rheumatoid polyneuropathy with rheumatoid arthritis of right ankle and foot
M05.572	Rheumatoid polyneuropathy with rheumatoid arthritis of left ankle and foot
M05.579	Rheumatoid polyneuropathy with rheumatoid arthritis of unspecified ankle and foot
M05.59	Rheumatoid polyneuropathy with rheumatoid arthritis of multiple sites
M05.60	Rheumatoid arthritis of unspecified site with involvement of other organs and systems
M05.611	Rheumatoid arthritis of right shoulder with involvement of other organs and systems
M05.612	Rheumatoid arthritis of left shoulder with involvement of other organs and systems
M05.619	Rheumatoid arthritis of unspecified shoulder with involvement of other organs and systems
M05.621	Rheumatoid arthritis of right elbow with involvement of other organs and systems
M05.622	Rheumatoid arthritis of left elbow with involvement of other organs and systems

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

M05.629	Rheumatoid arthritis of unspecified elbow with involvement of other organs and systems
M05.631	Rheumatoid arthritis of right wrist with involvement of other organs and systems
M05.632	Rheumatoid arthritis of left wrist with involvement of other organs and systems
M05.639	Rheumatoid arthritis of unspecified wrist with involvement of other organs and systems
M05.641	Rheumatoid arthritis of right hand with involvement of other organs and systems
M05.642	Rheumatoid arthritis of left hand with involvement of other organs and systems
M05.649	Rheumatoid arthritis of unspecified hand with involvement of other organs and systems
M05.651	Rheumatoid arthritis of right hip with involvement of other organs and systems
M05.652	Rheumatoid arthritis of left hip with involvement of other organs and systems
M05.659	Rheumatoid arthritis of unspecified hip with involvement of other organs and systems
M05.661	Rheumatoid arthritis of right knee with involvement of other organs and systems
M05.662	Rheumatoid arthritis of left knee with involvement of other organs and systems
M05.669	Rheumatoid arthritis of unspecified knee with involvement of other organs and systems
M05.671	Rheumatoid arthritis of right ankle and foot with involvement of other organs and systems
M05.672	Rheumatoid arthritis of left ankle and foot with involvement of other organs and systems
M05.679	Rheumatoid arthritis of unspecified ankle and foot with involvement of other organs and systems
M05.69	Rheumatoid arthritis of multiple sites with involvement of other organs and systems
M05.7A	Rheumatoid arthritis with rheumatoid factor of other specified site without organ or systems
M05.711	Rheumatoid arthritis with rheumatoid factor of right shoulder without organ or systems involvement
M05.712	Rheumatoid arthritis with rheumatoid factor of left shoulder without organ or systems involvement
M05.719	Rheumatoid arthritis with rheumatoid factor of unspecified shoulder without organ or systems
M05.721	Rheumatoid arthritis with rheumatoid factor of right elbow without organ or systems involvement
M05.722	Rheumatoid arthritis with rheumatoid factor of left elbow without organ or systems involvement
M05.729	Rheumatoid arthritis with rheumatoid factor of unspecified elbow without organ or systems
M05.731	Rheumatoid arthritis with rheumatoid factor of right wrist without organ or systems involvement
M05.732	Rheumatoid arthritis with rheumatoid factor of left wrist without organ or systems involvement
M05.739	Rheumatoid arthritis with rheumatoid factor of unspecified wrist without organ or systems
M05.741	Rheumatoid arthritis with rheumatoid factor of right hand without organ or systems involvement
M05.742	Rheumatoid arthritis with rheumatoid factor of left hand without organ or systems involvement
M05.749	Rheumatoid arthritis with rheumatoid factor of unspecified hand without organ or systems
M05.751	Rheumatoid arthritis with rheumatoid factor of right hip without organ or systems involvement
M05.752	Rheumatoid arthritis with rheumatoid factor of left hip without organ or systems involvement
M05.759	Rheumatoid arthritis with rheumatoid factor of unspecified hip without organ or systems involvement
M05.761	Rheumatoid arthritis with rheumatoid factor of right knee without organ or systems involvement
M05.762	Rheumatoid arthritis with rheumatoid factor of left knee without organ or systems involvement
M05.769	Rheumatoid arthritis with rheumatoid factor of unspecified knee without organ or systems
M05.771	Rheumatoid arthritis with rheumatoid factor of right ankle and foot without organ or systems

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

M05.772	Rheumatoid arthritis with rheumatoid factor of left ankle and foot without organ or systems
M05.779	Rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot without organ or systems
M05.79	Rheumatoid arthritis with rheumatoid factor of multiple sites without organ or systems involvement
M05.8A	Other rheumatoid arthritis with rheumatoid factor of other specified site
M05.811	Other rheumatoid arthritis with rheumatoid factor of right shoulder
M05.812	Other rheumatoid arthritis with rheumatoid factor of left shoulder
M05.819	Other rheumatoid arthritis with rheumatoid factor of unspecified shoulder
M05.821	Other rheumatoid arthritis with rheumatoid factor of right elbow
M05.822	Other rheumatoid arthritis with rheumatoid factor of left elbow
M05.829	Other rheumatoid arthritis with rheumatoid factor of unspecified elbow
M05.831	Other rheumatoid arthritis with rheumatoid factor of right wrist
M05.832	Other rheumatoid arthritis with rheumatoid factor of left wrist
M05.839	Other rheumatoid arthritis with rheumatoid factor of unspecified wrist
M05.841	Other rheumatoid arthritis with rheumatoid factor of right hand
M05.842	Other rheumatoid arthritis with rheumatoid factor of left hand
M05.849	Other rheumatoid arthritis with rheumatoid factor of unspecified hand
M05.851	Other rheumatoid arthritis with rheumatoid factor of right hip
M05.852	Other rheumatoid arthritis with rheumatoid factor of left hip
M05.859	Other rheumatoid arthritis with rheumatoid factor of unspecified hip
M05.861	Other rheumatoid arthritis with rheumatoid factor of right knee
M05.862	Other rheumatoid arthritis with rheumatoid factor of left knee
M05.869	Other rheumatoid arthritis with rheumatoid factor of unspecified knee
M05.871	Other rheumatoid arthritis with rheumatoid factor of right ankle and foot
M05.872	Other rheumatoid arthritis with rheumatoid factor of left ankle and foot
M05.879	Other rheumatoid arthritis with rheumatoid factor of unspecified ankle and foot
M05.89	Other rheumatoid arthritis with rheumatoid factor of multiple sites
M05.9	Rheumatoid arthritis with rheumatoid factor, unspecified
M06.0A	Rheumatoid arthritis without rheumatoid factor, other specified site
M06.011	Rheumatoid arthritis without rheumatoid factor, right shoulder
M06.012	Rheumatoid arthritis without rheumatoid factor, left shoulder
M06.019	Rheumatoid arthritis without rheumatoid factor, unspecified shoulder
M06.021	Rheumatoid arthritis without rheumatoid factor, right elbow
M06.022	Rheumatoid arthritis without rheumatoid factor, left elbow
M06.029	Rheumatoid arthritis without rheumatoid factor, unspecified elbow
M06.031	Rheumatoid arthritis without rheumatoid factor, right wrist
M06.032	Rheumatoid arthritis without rheumatoid factor, left wrist
M06.039	Rheumatoid arthritis without rheumatoid factor, unspecified wrist

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

M06.041	Rheumatoid arthritis without rheumatoid factor, right hand
M06.042	Rheumatoid arthritis without rheumatoid factor, left hand
M06.049	Rheumatoid arthritis without rheumatoid factor, unspecified hand
M06.051	Rheumatoid arthritis without rheumatoid factor, right hip
M06.052	Rheumatoid arthritis without rheumatoid factor, left hip
M06.059	Rheumatoid arthritis without rheumatoid factor, unspecified hip
M06.061	Rheumatoid arthritis without rheumatoid factor, right knee
M06.062	Rheumatoid arthritis without rheumatoid factor, left knee
M06.069	Rheumatoid arthritis without rheumatoid factor, unspecified knee
M06.071	Rheumatoid arthritis without rheumatoid factor, right ankle and foot
M06.072	Rheumatoid arthritis without rheumatoid factor, left ankle and foot
M06.079	Rheumatoid arthritis without rheumatoid factor, unspecified ankle and foot
M06.08	Rheumatoid arthritis without rheumatoid factor, vertebrae
M06.09	Rheumatoid arthritis without rheumatoid factor, multiple sites
M06.8A	Other specified rheumatoid arthritis, other specified site
M06.811	Other specified rheumatoid arthritis, right shoulder
M06.812	Other specified rheumatoid arthritis, left shoulder
M06.819	Other specified rheumatoid arthritis, unspecified shoulder
M06.821	Other specified rheumatoid arthritis, right elbow
M06.822	Other specified rheumatoid arthritis, left elbow
M06.829	Other specified rheumatoid arthritis, unspecified elbow
M06.831	Other specified rheumatoid arthritis, right wrist
M06.832	Other specified rheumatoid arthritis, left wrist
M06.839	Other specified rheumatoid arthritis, unspecified wrist
M06.841	Other specified rheumatoid arthritis, right hand
M06.842	Other specified rheumatoid arthritis, left hand
M06.849	Other specified rheumatoid arthritis, unspecified hand
M06.851	Other specified rheumatoid arthritis, right hip
M06.852	Other specified rheumatoid arthritis, left hip
M06.859	Other specified rheumatoid arthritis, unspecified hip
M06.861	Other specified rheumatoid arthritis, right knee
M06.862	Other specified rheumatoid arthritis, left knee
M06.869	Other specified rheumatoid arthritis, unspecified knee
M06.871	Other specified rheumatoid arthritis, right ankle and foot
M06.872	Other specified rheumatoid arthritis, left ankle and foot
M06.879	Other specified rheumatoid arthritis, unspecified ankle and foot
M06.88	Other specified rheumatoid arthritis, vertebrae

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

M06.89	Other specified rheumatoid arthritis, multiple sites
M06.9	Rheumatoid arthritis, unspecified
M32.14	Glomerular disease in systemic lupus erythematosus
M31.1	Thrombotic microangiopathy
M31.30	Wegener's granulomatosis without renal involvement
M31.31	Wegener's granulomatosis with renal involvement
M31.7	Microscopic polyangiitis
M79.10	Myalgia, unspecified site
M79.11	Myalgia of mastication muscle
M79.12	Myalgia of auxiliary muscles, head and neck
M79.18	Myalgia, other site
T86.09	Other complications of bone marrow transplant
Z94.81	Bone marrow transplant status
Z94.89	Other transplanted organ and tissue status
Z94.9	Transplanted organ and tissue status, unspecified

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at:

<https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

Jurisdiction(s): 6, K	NCD/LCD/LCA Document (s): A52452
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a52452&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	
Jurisdiction(s): 5,8	NCD/LCD/LCA Document (s): A55639
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a55639&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Jurisdiction(s): J,M	NCD/LCD/LCA Document (s): A56380
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56380&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	
Jurisdiction(s): 15	NCD/LCD/LCA Document (s): A57160
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a57160&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	
Jurisdiction(s): 15	NCD/LCD/LCA Document (s): A58582
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a58582&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

RITUXIMAB -E- (Rituxan®, Truxima®, Ruxience®, Riabni™)
Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management