

Halaven® (eribulin) (Intravenous)

-E-

Document Number: IC-0403

Last Review Date: 04/04/2022

Date of Origin: 01/07/2019

Dates Reviewed: 01/2019, 04/2019, 07/2019, 10/2019, 01/2020, 04/2020, 07/2020, 10/2020, 05/2021, 04/2022

I. Length of Authorization

Coverage will be provided for six months and may be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Halaven 1 mg/2 mL solution in a single-dose vial for injection: 8 vials every 21 days

B. Max Units (per dose and over time) [HCPCS Unit]:

- 80 billable units every 21 days

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

Breast Cancer † ‡ 1-3,7e,9e,14e,17e,18e

- Patient has metastatic disease †; **AND**
 - Used as a single agent for patients who have previously received at least two chemotherapy regimens for the treatment of metastatic disease; **AND**
 - Prior therapy includes treatment with an anthracycline and a taxane in either the adjuvant or metastatic setting; **OR**
- Patient has recurrent unresectable or metastatic disease; **AND**
 - Used as a single agent for HER2-negative disease in patients who have previously received therapy with an anthracycline and a taxane; **AND**
 - Patient has hormone-receptor negative disease; **OR**
 - Patient has hormone receptor-positive disease with visceral crisis or is endocrine therapy refractory; **OR**
 - Used in combination with margetuximab-cmkb for HER2-positive disease as third-line therapy and beyond; **AND**

- Patient has previously been treated with at least two (2) prior HER2 regimens, at least one of which was used for metastatic disease; **OR**
- o Used in combination with trastuzumab for HER2-positive disease as first-line therapy; **AND**

- Use of eribulin will be restricted to patients with a contraindication or intolerance to pertuzumab/trastuzumab/docetaxel (or paclitaxel)

Liposarcoma † 1,2,4,20e

- Used as a single agent; **AND**
- Patient has unresectable or metastatic or recurrent disease; **AND**
- Patient has received prior anthracycline-based therapy (e.g., doxorubicin, etc.)

Preferred therapies and recommendations are determined by review of clinical evidence. NCCN category of recommendation is taken into account as a component of this review. Regimens deemed equally efficacious (i.e., those having the same NCCN categorization) are considered to be therapeutically equivalent.

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Ⓢ Orphan Drug

IV. Renewal Criteria ¹

Coverage can be renewed based on the following criteria:

- Patient continues to meet indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: QT-prolongation, severe neutropenia (ANC < 500/mm³), peripheral neuropathy (Grade 3 or 4), etc.

V. Dosage/Administration ^{1,6}

Indication	Dose
All Indications	Administer 1.4 mg/m ² , intravenously, on days 1 and 8, repeated every 21 days until disease progression or unacceptable toxicity

VI. Billing Code/Availability Information

HCPSC Code:

- J9179 – Injection, eribulin mesylate, 0.1 mg; 1 billable unit = 0.1mg

NDC:

- Halaven 1 mg/2 mL solution in a single-dose vial for injection: 62856-0389-xx

VII. References (STANDARD)

1. Halaven [package insert]. Woodcliff Lake, NJ; Eisai Inc; December 2021. Accessed March 2022.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) eribulin. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed March 2022.
3. Cortes J, O'Shaughnessy J, Loesch D, et al; EMBRACE (Eisai Metastatic Breast Cancer Study Assessing Physician's Choice Versus E7389) investigators. Eribulin monotherapy versus treatment of physician's choice in patients with metastatic breast cancer (EMBRACE): a phase 3 open-label randomised study. *Lancet*. 2011;377(9769):914-923.
4. Schöffski P, Chawla S, Maki RG, et al. Eribulin versus dacarbazine in previously treated patients with advanced liposarcoma or leiomyosarcoma: a randomised, open-label, multicentre, phase 3 trial. *Lancet*. 2016 Apr 16;387(10028):1629-37. doi: 10.1016/S0140-6736(15)01283-0. Epub 2016 Feb 10.
5. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Soft Tissue Sarcoma, Version 2.2021. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed March 2022.
6. Schöffski P, Ray-Coquard IL, Cioffi A, et al. Activity of eribulin mesylate in patients with soft-tissue sarcoma: a phase 2 study in four independent histological subtypes. *Lancet Oncol*. 2011;12(11):1045-1052.

VIII. References (ENHANCED)

- 1e. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) Breast Cancer, Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. Accessed March 2022.
- 2e. Sledge GW, Neuberg D, Bernardo P, et al. Phase III trial of doxorubicin, paclitaxel, and the combination of doxorubicin and paclitaxel as front-line chemotherapy for metastatic breast cancer: an intergroup trial (E1193). *J Clin Oncol*. 2003 Feb 15;21(4):588-92.

- 3e. Chan S, Friedrichs K, Noel D, et al. Prospective randomized trial of docetaxel versus doxorubicin in patients with metastatic breast cancer. *J Clin Oncol*. 1999 Aug;17(8):2341-54.
- 4e. Seidman AD, Berry D, Cirrincione C, et al. Randomized Phase III Trial of Weekly Compared With Every-3-Weeks Paclitaxel for Metastatic Breast Cancer, With Trastuzumab for all HER-2 Overexpressors and Random Assignment to Trastuzumab or Not in HER-2 Nonoverexpressors: Final Results of Cancer and Leukemia Group B Protocol 9840. *Journal of Clinical Oncology* 2008 26:10, 1642-1649.
- 5e. Seidman AD, Tiersten A, Hudis C, et al. Phase II trial of paclitaxel by 3-hour infusion as initial and salvage chemotherapy for metastatic breast cancer. *J Clin Oncol*. 1995 Oct;13(10):2575-81.
- 6e. Bajetta E, Procopio G, Celio L, et al. Safety and Efficacy of Two Different Doses of Capecitabine in the Treatment of Advanced Breast Cancer in Older Women. *Journal of Clinical Oncology* 2005 23:10, 2155-2161.
- 7e. Kaufman PA, Awada A, Twelves C, et al. Phase III open-label randomized study of eribulin mesylate versus capecitabine in patients with locally advanced or metastatic breast cancer previously treated with an anthracycline and a taxane. *J Clin Oncol*. 2015 Feb 20;33(6):594-601. doi: 10.1200/JCO.2013.52.4892. Epub 2015 Jan 20.
- 8e. Twelves C, Awada A, Cortes J, et al. Subgroup Analyses from a Phase 3, Open-Label, Randomized Study of Eribulin Mesylate Versus Capecitabine in Pretreated Patients with Advanced or Metastatic Breast Cancer. *Breast Cancer (Auckl)*. 2016 Jun 28;10:77-84. doi: 10.4137/BCBCR.S39615. eCollection 2016.
- 9e. Baselga J, Cortés J, Kim SB, et al. Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer. *N Engl J Med*. 2012 Jan 12;366(2):109-19. doi: 10.1056/NEJMoa1113216. Epub 2011 Dec 7.
- 10e. Swain SM, Baselga J, Kim SB, et al. Pertuzumab, trastuzumab, and docetaxel in HER2-positive metastatic breast cancer. *N Engl J Med*. 2015 Feb 19;372(8):724-34. doi: 10.1056/NEJMoa1413513.
- 11e. Datko FM, D'Andrea G, Dickler M, et al. Phase II study of pertuzumab, trastuzumab, and weekly paclitaxel in patients with HER2-overexpressing metastatic breast cancer (MBC). *Journal of Clinical Oncology* 2012 30:27_suppl, 134-134.
- 12e. Smyth LM, Iyengar NM, Chen MF, et al. Weekly paclitaxel with trastuzumab and pertuzumab in patients with HER2-overexpressing metastatic breast cancer: overall survival and updated progression-free survival results from a phase II study. *Breast Cancer Res Treat*. 2016 Jul;158(1):91-97. doi: 10.1007/s10549-016-3851-7. Epub 2016 Jun 15.
- 13e. Ellis PA, Barrios CH, Eiermann W, et al. Phase III, randomized study of trastuzumab emtansine (T-DM1) ± pertuzumab (P) vs trastuzumab + taxane (HT) for first-line treatment of HER2-positive MBC: Primary results from the MARIANNE study. *Journal of Clinical Oncology* 2015 33:15_suppl, 507-507.

- 14e. Verma S, Miles D, Gianni L, et al. Trastuzumab emtansine for HER2-positive advanced breast cancer. *N Engl J Med*. 2012 Nov 8;367(19):1783-91. doi: 10.1056/NEJMoa1209124. Epub 2012 Oct 1.
- 15e. Cameron D, Casey M, Oliva C, et al. Lapatinib plus capecitabine in women with HER-2-positive advanced breast cancer: final survival analysis of a phase III randomized trial. *Oncologist*. 2010;15(9):924-34. doi: 10.1634/theoncologist.2009-0181. Epub 2010 Aug 24.
- 16e. Blackwell KL, Burstein HJ, Storniolo AM, et al. Overall survival benefit with lapatinib in combination with trastuzumab for patients with human epidermal growth factor receptor 2-positive metastatic breast cancer: final results from the EGF104900 Study. *J Clin Oncol*. 2012 Jul 20;30(21):2585-92. doi: 10.1200/JCO.2011.35.6725. Epub 2012 Jun 11.
- 17e. von Minckwitz G, du Bois A, Schmidt M, et al. Trastuzumab beyond progression in human epidermal growth factor receptor 2-positive advanced breast cancer: a german breast group 26/breast international group 03-05 study. *J Clin Oncol*. 2009 Apr 20;27(12):1999-2006. doi: 10.1200/JCO.2008.19.6618. Epub 2009 Mar 16.
- 18e. Wilks S, Puhalla S, O'Shaughnessy J, et al. Phase 2, multicenter, single-arm study of eribulin mesylate with trastuzumab as first-line therapy for locally recurrent or metastatic HER2-positive breast cancer. *Clin Breast Cancer*. 2014 Dec;14(6):405-12. doi: 10.1016/j.clbc.2014.04.004. Epub 2014 Jun 2.
- 19e. Demetri GD, von Mehren M, Jones RL, et al. Efficacy and Safety of Trabectedin or Dacarbazine for Metastatic Liposarcoma or Leiomyosarcoma After Failure of Conventional Chemotherapy: Results of a Phase III Randomized Multicenter Clinical Trial. *J Clin Oncol*. 2016 Mar 10;34(8):786-93.
- 20e. Demetri GD, Schöffski P, Grignani G, et al. Activity of Eribulin in Patients With Advanced Liposarcoma Demonstrated in a Subgroup Analysis From a Randomized Phase III Study of Eribulin Versus Dacarbazine. *J Clin Oncol*. 2017 Oct 20;35(30):3433-3439. doi: 10.1200/JCO.2016.71.6605. Epub 2017 Aug 30.
- 21e. Bui-Nguyen B, Butrynski JE, Penel N, et al. A phase IIb multicentre study comparing the efficacy of trabectedin to doxorubicin in patients with advanced or metastatic untreated soft tissue sarcoma: the TRUSTS trial. *Eur J Cancer*. 2015 Jul;51(10):1312-20.
- 22e. Judson I, Verweij J, Gelderblom H, et al. Doxorubicin alone versus intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic soft-tissue sarcoma: a randomised controlled phase 3 trial. *Lancet Oncol*. 2014 Apr;15(4):415-23.
- 23e. Seddon B, Strauss SJ, Whelan J, et al. Gemcitabine and docetaxel versus doxorubicin as first-line treatment in previously untreated advanced unresectable or metastatic soft-tissue sarcomas (GeDDiS): a randomised controlled phase 3 trial. *Lancet Oncol*. 2017 Oct;18(10):1397-1410.
- 24e. Le Cesne A, Blay J-Y, Cupissol D, Italiano A, Delcambre C, Penel N, et al. Results of a prospective randomized phase III T-SAR trial comparing trabectedin vs best supportive care (BSC) in patients with pretreated advanced soft tissue sarcoma (ASTS) *Ann Oncol*. 2016;27(suppl 6):1396O.

- 25e. Le Cesne A, Blay JY, Judson I, et al. Phase II study of ET-743 in advanced soft tissue sarcomas: a European Organisation for the Research and Treatment of Cancer (EORTC) soft tissue and bone sarcoma group trial. *J Clin Oncol*. 2005 Jan 20;23(3):576-84.
- 26e. Kawai A, Araki N, Sugiura H, et al. Trabectedin monotherapy after standard chemotherapy versus best supportive care in patients with advanced, translocation-related sarcoma: a randomised, open-label, phase 2 study. *Lancet Oncol*. 2015 Apr;16(4):406-16. doi: 10.1016/S1470-2045(15)70098-7. Epub 2015 Mar 18.
- 27e. Blay JY, Leahy MG, Nguyen BB, et al. Randomised phase III trial of trabectedin versus doxorubicin-based chemotherapy as first-line therapy in translocation-related sarcomas. *Eur J Cancer*. 2014 Apr;50(6):1137-47. doi: 10.1016/j.ejca.2014.01.012. Epub 2014 Feb 7.
- 28e. Baruchel S, Pappo A, Krailo M, et al. A phase 2 trial of trabectedin in children with recurrent rhabdomyosarcoma, Ewing sarcoma and non-rhabdomyosarcoma soft tissue sarcomas: a report from the Children's Oncology Group. *Eur J Cancer*. 2012 Mar;48(4):579-85. doi: 10.1016/j.ejca.2011.09.027. Epub 2011 Nov 14.
- 29e. Arndt CA, Stoner JA, Hawkins DS, et al. Vincristine, actinomycin, and cyclophosphamide compared with vincristine, actinomycin, and cyclophosphamide alternating with vincristine, topotecan, and cyclophosphamide for intermediate-risk rhabdomyosarcoma: children's oncology group study D9803. *J Clin Oncol*. 2009 Nov 1;27(31):5182-8. doi: 10.1200/JCO.2009.22.3768. Epub 2009 Sep 21.
- 30e. Penel N, Bui BN, Bay JO, et al. Phase II trial of weekly paclitaxel for unresectable angiosarcoma: the ANGIOTAX Study. *J Clin Oncol*. 2008 Nov 10;26(32):5269-74. doi: 10.1200/JCO.2008.17.3146. Epub 2008 Sep 22.
- 31e. Tap WD, Jones RL, Van Tine BA, et al. Olaratumab and doxorubicin versus doxorubicin alone for treatment of soft-tissue sarcoma: an open-label phase 1b and randomised phase 2 trial. *Lancet*. 2016 Jul 30;388(10043):488-97. doi: 10.1016/S0140-6736(16)30587-6. Epub 2016 Jun 9.
- 32e. Agulnik M, Yarber JL, Okuno SH, et al. An open-label, multicenter, phase II study of bevacizumab for the treatment of angiosarcoma and epithelioid hemangioendotheliomas. *Ann Oncol*. 2013 Jan;24(1):257-63. doi: 10.1093/annonc/mds237. Epub 2012 Aug 21.
- 33e. Hawkins DS, Chi YY, Anderson JR, et al. Addition of Vincristine and Irinotecan to Vincristine, Dactinomycin, and Cyclophosphamide Does Not Improve Outcome for Intermediate-Risk Rhabdomyosarcoma: A Report From the Children's Oncology Group. *J Clin Oncol*. 2018 Sep 20;36(27):2770-2777. doi: 10.1200/JCO.2018.77.9694. Epub 2018 Aug 9.
- 34e. Kawai A, Araki N, Naito Y, et al. Phase 2 study of eribulin in patients with previously treated advanced or metastatic soft tissue sarcoma. *Jpn J Clin Oncol*. 2017 Feb 1;47(2):137-144. doi: 10.1093/jjco/hyw175.
- 35e. Pautier P, Floquet A, Penel N, et al. Randomized multicenter and stratified phase II study of gemcitabine alone versus gemcitabine and docetaxel in patients with metastatic or relapsed leiomyosarcomas: a Federation Nationale des Centres de Lutte Contre le Cancer

(FNCLCC) French Sarcoma Group Study (TAXOGEM study). *Oncologist*. 2012;17(9):1213-20. Epub 2012 Aug 20.

36e. Rugo HS, Im SA, Shaw-Wright GL, et al. SOPHIA primary analysis: A phase 3 (P3) study of margetuximab (M) + chemotherapy (C) versus trastuzumab (T) + C in patients (pts) with HER2+ metastatic (met) breast cancer (MBC) after prior anti-HER2 therapies (Tx). *Journal of Clinical Oncology* 2019 37:15_suppl, 1000-1000.

37e. Saura C, Oliveira M, Feng YH, et al. NALA Investigators. Neratinib Plus Capecitabine Versus Lapatinib Plus Capecitabine in HER2-Positive Metastatic Breast Cancer Previously Treated With ≥ 2 HER2-Directed Regimens: Phase III NALA Trial. *J Clin Oncol*. 2020 Sep 20;38(27):3138-3149. doi: 10.1200/JCO.20.00147. Epub 2020 Jul 17.

38e. Magellan Health, Magellan Rx Management. Halaven Clinical Literature Review Analysis. Last updated March 2022. Accessed March 2022.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C47.0	Malignant neoplasm of peripheral nerves of head, face and neck
C47.10	Malignant neoplasm of peripheral nerves of unspecified upper limb, including shoulder
C47.11	Malignant neoplasm of peripheral nerves of right upper limb, including shoulder
C47.12	Malignant neoplasm of peripheral nerves of left upper limb, including shoulder
C47.20	Malignant neoplasm of peripheral nerves of unspecified lower limb, including hip
C47.21	Malignant neoplasm of peripheral nerves of right lower limb, including hip
C47.22	Malignant neoplasm of peripheral nerves of left lower limb, including hip
C47.3	Malignant neoplasm of peripheral nerves of thorax
C47.4	Malignant neoplasm of peripheral nerves of abdomen
C47.5	Malignant neoplasm of peripheral nerves of pelvis
C47.6	Malignant neoplasm of peripheral nerves of trunk, unspecified
C47.8	Malignant neoplasm of overlapping sites of peripheral nerves and autonomic nervous system
C47.9	Malignant neoplasm of peripheral nerves and autonomic nervous system, unspecified
C48.0	Malignant neoplasm of retroperitoneum
C48.1	Malignant neoplasm of specified parts of peritoneum
C48.2	Malignant neoplasm of peritoneum, unspecified
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum
C49.0	Malignant neoplasm of connective and soft tissue of head, face and neck
C49.10	Malignant neoplasm of connective and soft tissue of unspecified upper limb, including shoulder
C49.11	Malignant neoplasm of connective and soft tissue of right upper limb
C49.12	Malignant neoplasm of connective and soft tissue of left lower limb
C49.20	Malignant neoplasm of connective and soft tissue of unspecified lower limb, including hip
C49.21	Malignant neoplasm of connective and soft tissue of right lower limb

ICD-10	ICD-10 Description
C49.22	Malignant neoplasm of connective and soft tissue of left lower limb
C49.3	Malignant neoplasm of connective and soft tissue of thorax
C49.4	Malignant neoplasm of connective and soft tissue of abdomen
C49.5	Malignant neoplasm of connective and soft tissue of pelvis
C49.6	Malignant neoplasm of connective and soft tissue of trunk, unspecified
C49.8	Malignant neoplasm of overlapping sites of connective and soft tissue
C49.9	Malignant neoplasm of connective and soft tissue, unspecified
C50.011	Malignant neoplasm of nipple and areola, right female breast
C50.012	Malignant neoplasm of nipple and areola, left female breast
C50.019	Malignant neoplasm of nipple and areola, unspecified female breast
C50.021	Malignant neoplasm of nipple and areola, right male breast
C50.022	Malignant neoplasm of nipple and areola, left male breast
C50.029	Malignant neoplasm of nipple and areola, unspecified male breast
C50.111	Malignant neoplasm of central portion of right female breast
C50.112	Malignant neoplasm of central portion of left female breast
C50.119	Malignant neoplasm of central portion of unspecified female breast
C50.121	Malignant neoplasm of central portion of right male breast
C50.122	Malignant neoplasm of central portion of left male breast
C50.129	Malignant neoplasm of central portion of unspecified male breast
C50.211	Malignant neoplasm of upper-inner quadrant of right female breast
C50.212	Malignant neoplasm of upper-inner quadrant /of left female breast
C50.219	Malignant neoplasm of upper-inner quadrant of unspecified female breast
C50.221	Malignant neoplasm of upper-inner quadrant of right male breast
C50.222	Malignant neoplasm of upper-inner quadrant of left male breast
C50.229	Malignant neoplasm of upper-inner quadrant of unspecified male breast
C50.311	Malignant neoplasm of lower-inner quadrant of right female breast
C50.312	Malignant neoplasm of lower-inner quadrant of left female breast
C50.319	Malignant neoplasm of lower-inner quadrant of unspecified female breast
C50.321	Malignant neoplasm of lower-inner quadrant of right male breast
C50.322	Malignant neoplasm of lower-inner quadrant of left male breast
C50.329	Malignant neoplasm of lower-inner quadrant of unspecified male breast
C50.411	Malignant neoplasm of upper-outer quadrant of right female breast
C50.412	Malignant neoplasm of upper-outer quadrant of left female breast
C50.419	Malignant neoplasm of upper-outer quadrant of unspecified female breast
C50.421	Malignant neoplasm of upper-outer quadrant of right male breast
C50.422	Malignant neoplasm of upper-outer quadrant of left male breast

ICD-10	ICD-10 Description
C50.429	Malignant neoplasm of upper-outer quadrant of unspecified male breast
C50.511	Malignant neoplasm of lower-outer quadrant of right female breast
C50.512	Malignant neoplasm of lower-outer quadrant of left female breast
C50.519	Malignant neoplasm of lower-outer quadrant of unspecified female breast
C50.521	Malignant neoplasm of lower-outer quadrant of right male breast
C50.522	Malignant neoplasm of lower-outer quadrant of left male breast
C50.529	Malignant neoplasm of lower-outer quadrant of unspecified male breast
C50.611	Malignant neoplasm of axillary tail of right female breast
C50.612	Malignant neoplasm of axillary tail of left female breast
C50.619	Malignant neoplasm of axillary tail of unspecified female breast
C50.621	Malignant neoplasm of axillary tail of right male breast
C50.622	Malignant neoplasm of axillary tail of left male breast
C50.629	Malignant neoplasm of axillary tail of unspecified male breast
C50.811	Malignant neoplasm of overlapping sites of right female breast
C50.812	Malignant neoplasm of overlapping sites of left female breast
C50.819	Malignant neoplasm of overlapping sites of unspecified female breast
C50.821	Malignant neoplasm of overlapping sites of right male breast
C50.822	Malignant neoplasm of overlapping sites of left male breast
C50.829	Malignant neoplasm of overlapping sites of unspecified male breast
C50.911	Malignant neoplasm of unspecified site of right female breast
C50.912	Malignant neoplasm of unspecified site of left female breast
C50.919	Malignant neoplasm of unspecified site of unspecified female breast
C50.921	Malignant neoplasm of unspecified site of right male breast
C50.922	Malignant neoplasm of unspecified site of left male breast
C50.929	Malignant neoplasm of unspecified site of unspecified male breast

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions

Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC