

Vabysmo™ (faricimab-svoa) (Intravitreal)

Document Number: SHP-0659

Last Review Date: 07/01/2022

Date of Origin: 03/01/2022

Dates Reviewed: 03/2022, 07/2022

I. Length of Authorization

Coverage will be provided annually and may be renewed.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC unit]:

- 2 mg injection: 1 vial per eye every 28 days

B. Max Units (per dose and over time) [HCPCS Unit]:

Diagnosis	MU for Initial Dosing	MU for Maintenance Dosing
Neovascular age-related macular degeneration (AMD)	60 billable units (6 mg) every 28 days x 4 doses	60 billable units (6 mg) every 28 days
Diabetic Macular Edema (DME)	60 billable units (6 mg) every 28 days x 6 doses	60 billable units (6 mg) every 28 days

(Max units are based on administration to both eyes)

III. Initial Approval Criteria ¹⁻⁸

Coverage is provided in the following conditions:

- Patient must try and have an inadequate response, contraindication, or intolerance to an adequate trial of bevacizumab in either eye prior to consideration of a non-preferred product; **AND**
- Patient is at least 18 years of age; **AND**

Universal Criteria

- Patient is free of ocular and/or peri-ocular infections; **AND**
- Patient does not have active intraocular inflammation; **AND**
- Therapy will not be used with other ophthalmic VEGF inhibitors (i.e., aflibercept, brolocizumab-dbl, ranibizumab, pegaptanib, bevacizumab, etc.); **AND**
- Patient's best corrected visual acuity (BCVA) is measured at baseline and periodically during treatment; **AND**

- Patient has a definitive diagnosis of one of the following:

Neovascular (Wet) Age-Related Macular Degeneration (nAMD) †*

Diabetic Macular Edema (DME) †*

**Patients with an insufficient response during initial therapy administered every 4 weeks for at least 4 doses may continue with dosing every 4 weeks. Patients with loss of response to maintenance therapy administered at less frequent intervals may increase the dosing frequency in a step-wise manner until response is regained. (Refer to Section V)*

† FDA Approved Indication(s)

IV. Renewal Criteria ¹⁻⁸

Coverage can be renewed based upon the following criteria:

- Patient continues to meet the universal and indication-specific relevant criteria as identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include the following: endophthalmitis and retinal detachments, increase in intraocular pressure, arterial thromboembolic events (ATE), etc.; **AND**
- Patient has had a beneficial response to therapy (e.g., improvement in the baseline best corrected visual acuity (BCVA), etc.) and continued administration is necessary for the maintenance treatment of the condition

V. Dosage/Administration ¹

Indication	Dose
nAMD	<p><u>Initiation:</u></p> <ul style="list-style-type: none"> • 6 mg intravitreally per affected eye once every 4 weeks (approximately every 28 days \pm7 days, monthly) for the first four doses (16 weeks or 4 months) <p><u>Maintenance:</u></p> <ul style="list-style-type: none"> • Follow the initial four doses with optical coherence tomography and visual acuity evaluations 8 and 12 weeks later to ascertain whether to give a 6 mg dose via intravitreal injection on one of the following three regimens: <ul style="list-style-type: none"> – Weeks 28 and 44 (every 16 weeks); or – Weeks 24, 36 and 48 (every 12 weeks); or – Weeks 20, 28, 36 and 44 (every 8 weeks) <p><u>Note:</u> Additional efficacy was not demonstrated in most patients when VABYSMO was dosed every 4 weeks compared to every 8 weeks. Some patients may need every 4 week (approximately monthly) dosing after the first four doses (16 weeks or 4 months).</p>
Diabetic macular edema	<ul style="list-style-type: none"> • 6 mg intravitreally per affected eye once every 4 weeks (approximately every 28 days \pm7 days, monthly) for at least four doses. If after at least 4 doses, resolution of edema based on the central subfield thickness (CST) of the macula as measured by optical coherence

	<p>tomography is achieved, then the interval of dosing may be modified by extensions of up to 4 week interval increments or reductions of up to 8 week interval increments based on CST and visual acuity evaluations through week 52; OR</p> <ul style="list-style-type: none"> 6 mg intravitreally per affected eye once every 4 weeks (approximately every 28 days \pm7 days, monthly) for the first six doses, followed by 6 mg dose via intravitreal injection at intervals of every 8 weeks (2 months) over the next 28 weeks. <p><u>Note:</u> Additional efficacy was not demonstrated in most patients when VABYSMO was dosed every 4 weeks compared to every 8 weeks. Some patients may need every 4 week (monthly) dosing after the first four doses.</p>
--	--

VI. Billing Code/Availability Information

HCPCS Code:

- J3590 – Unclassified biologics
- C9097 – Injection, faricimab-svoa, 0.1 mg; 1 billable unit = 0.1 mg (*Effective 07/01/2022*)
- C9399 – Unclassified drugs or biologicals (*Discontinue use on 07/01/2022*)

NDC:

- Vabysmo 6 mg/0.05 mL (concentration 120 mg/mL) solution for injection, overfilled single-use vial: 50242 -0096-xx

VII. References

- Vabysmo [package insert]. South San Francisco, CA; Genentech, Inc.; January 2022. Accessed January 2022.
- Solomon SD, Chew E, Duh EJ, et al. Diabetic Retinopathy: A Position Statement by the American Diabetes Association. *Diabetes Care*. 2017 Mar;40(3):412-418.
- American Academy of Ophthalmology-Preferred Practice Patterns (AAO-PPP) Retina/Vitreous Panel, Hoskins Center for Quality Eye Care. Diabetic Retinopathy PPP – Update 2017. Nov 2017.
- American Academy of Ophthalmology-Preferred Practice Patterns (AAO-PPP) Retina/Vitreous Panel, Hoskins Center for Quality Eye Care. Retinal Vein Occlusions PPP – Update 2017. Nov 2017.
- American Academy of Ophthalmology-Preferred Practice Patterns (AAO-PPP) Retina/Vitreous Panel, Hoskins Center for Quality Eye Care. Age-Related Macular Degeneration PPP – Update 2017. Nov 2017.
- Royal College of Ophthalmologists. Clinical Guidelines – Retinal Vein Occlusion (RVO) Guidelines – July 2015. Accessed at <https://www.rcophth.ac.uk/standards-publications-research/clinical-guidelines>.
- Heier JS, Khanani AM, Quezada et al; TENAYA and LUCERNE Investigators. Efficacy, durability, and safety of intravitreal faricimab up to every 16 weeks for neovascular age-related macular degeneration (TENAYA and LUCERNE): two randomised, double-masked,

- phase 3, non-inferiority trials. *Lancet*. 2022 Jan 21. pii: S0140-6736(22)00010-1. doi: 10.1016/S0140-6736(22)00010-1.
8. Wykoff CC, Abreu F, Adamis AP, et al; YOSEMITE and RHINE Investigators. Efficacy, durability, and safety of intravitreal faricimab with extended dosing up to every 16 weeks in patients with diabetic macular oedema (YOSEMITE and RHINE): two randomised, double-masked, phase 3 trials. *Lancet*. 2022 Jan 21. pii: S0140-6736(22)00018-6. doi: 10.1016/S0140-6736(22)00018-6.
 9. American Academy of Ophthalmology Retina/Vitreous Panel. Age-related macular degeneration. San Francisco (CA): American Academy of Ophthalmology (AAO); 2008. 37 p.
 10. Intravitreal bevacizumab (Avastin®) treatment of macular edema in central retinal vein occlusion: a short-term study. *Retina*. 2006 Mar; 26(3): 279-84
 11. Epstein DL, Algvere PV, von Wendt G, et al: Bevacizumab for macular edema in central retinal vein occlusion: a prospective, randomized, double-masked clinical study. *Ophthalmology* 2012; 119(6):1184-1189.
 12. Cekic O, Cakir M, Yazici AT, et al: A comparison of three different intravitreal treatment modalities of macular edema due to branch retinal vein occlusion. *Curr Eye Res* 2010; 35(10):925-929.
 13. Moradian S, Ahmadieh H, Malihi M, et al. Intravitreal bevacizumab in active progressive proliferative diabetic retinopathy. *Graefes Arch Clin Exp Ophthalmol* 2008;246:1699-1705.
 14. Short-term safety and efficacy of intravitreal bevacizumab (Avastin®) for neovascular age-related macular degeneration. *Retina*. 2006 May-Jun; 26(5): 495-511
 15. Rich RM, Rosenfeld PJ, Puliakit CA, et al. Short-term safety and efficacy of intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration. *Retina* 2006;26:495-511.
 16. Avery RL, Pieramici DJ, Rabena MD, et al. Intravitreal bevacizumab (Avastin) for neovascular age-related macular degeneration. *Ophthalmol* 2006;113:363-72.
 17. Ruiz-Moreno JM, Montero JA, Araiz J, et al. Intravitreal anti-vascular endothelial growth factor therapy for choroidal neovascularization secondary to pathologic myopia: six years outcome. *Retina*. 2015 Dec;35(12):2450-6.
 18. Gharbiya M, Giustolisi R, Allievi F, et al. Choroidal neovascularization in pathologic myopia: intravitreal ranibizumab versus bevacizumab--a randomized controlled trial. *Am J Ophthalmol*. 2010 Mar;149(3):458-64.e1.
 19. Iacono P, Parodi MB, Papayannis A, et al. Intravitreal ranibizumab versus bevacizumab for treatment of myopic choroidal neovascularization. *Retina*. 2012 Sep;32(8):1539-46.
 20. The CATT Research Group. Ranibizumab and Bevacizumab for Neovascular Age-Related Macular Degeneration. *N Engl J Med* 2011; 364:1897-1908
 21. Wang E & Chen Y: Intravitreal anti-vascular endothelial growth factor for choroidal neovascularization secondary to pathologic myopia: systematic review and meta-analysis. *Retina* 2013; 33(7):1375-1392.

22. Scott IU, Bressler NM, Bressler SB, Browning DJ, Chan CK, Danis RP, Davis MD, Kollman C, Qin H; Diabetic Retinopathy Clinical Research Network Study Group. Agreement between clinician and reading center gradings of diabetic retinopathy severity level at baseline in a phase 2 study of intravitreal bevacizumab for diabetic macular edema. *Retina*.2008 Jan;28(1):36-40.
23. Diabetic Retinopathy Clinical Research Network, Scott IU, Edwards AR, Beck RW, Bressler NM, Chan CK, Elman MJ, Friedman SM, Greven CM, Maturi RK, Pieramici DJ, Shami M, Singerman LJ, Stockdale CR. A phase II randomized clinical trial of intravitreal bevacizumab for diabetic macular edema. *Ophthalmology*.2007 Oct;114(10):1860-7
24. Diabetic Retinopathy Clinical Research Network, Wells JA, Glassman AR, Ayala AR, Jampol LM, Aiello LP, Antoszyk AN, Arnold-Bush B, Baker CW, Bressler NM, Browning DJ, Elman MJ, Ferris FL, Friedman SM, Melia M, Pieramici DJ, Sun JK, Beck RW. Aflibercept, Bevacizumab, or Ranibizumab for Diabetic Macular Edema. *N Engl J Med*. 2015 Mar 26; 372(13): 1193–1203.
25. Wells JA, Glassman AR, Jampol LM, et al., for the Diabetic Retinopathy Clinical Research Network. Association of Baseline Visual Acuity and Retinal Thickness with 1-year Efficacy of Aflibercept, Bevacizumab, and Ranibizumab for Diabetic Macular Edema. *JAMA Ophthalmol*. 2016 Feb;134(2):127-34.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
E08.311	Diabetes mellitus due to underlying condition with unspecified diabetic retinopathy with macular edema
E08.3211	Diabetes mellitus due to underlying condition with mild nonproliferative diabetic retinopathy with macular edema, right eye
E08.3212	Diabetes mellitus due to underlying condition with mild nonproliferative diabetic retinopathy with macular edema, left eye
E08.3213	Diabetes mellitus due to underlying condition with mild nonproliferative diabetic retinopathy with macular edema, bilateral
E08.3219	Diabetes mellitus due to underlying condition with mild nonproliferative diabetic retinopathy with macular edema, unspecified eye
E08.3311	Diabetes mellitus due to underlying condition with moderate nonproliferative diabetic retinopathy with macular edema, right eye
E08.3312	Diabetes mellitus due to underlying condition with moderate nonproliferative diabetic retinopathy with macular edema, left eye
E08.3313	Diabetes mellitus due to underlying condition with moderate nonproliferative diabetic retinopathy with macular edema, bilateral
E08.3319	Diabetes mellitus due to underlying condition with moderate nonproliferative diabetic retinopathy with macular edema, unspecified eye
E08.3411	Diabetes mellitus due to underlying condition with severe nonproliferative diabetic retinopathy with macular edema, right eye
E08.3412	Diabetes mellitus due to underlying condition with severe nonproliferative diabetic retinopathy with macular edema, left eye
E08.3413	Diabetes mellitus due to underlying condition with severe nonproliferative diabetic retinopathy with macular edema, bilateral
E08.3419	Diabetes mellitus due to underlying condition with severe nonproliferative diabetic retinopathy with macular edema, unspecified eye

ICD-10	ICD-10 Description
E08.3511	Diabetes mellitus due to underlying condition with proliferative diabetic retinopathy with macular edema, right eye
E08.3512	Diabetes mellitus due to underlying condition with proliferative diabetic retinopathy with macular edema, left eye
E08.3513	Diabetes mellitus due to underlying condition with proliferative diabetic retinopathy with macular edema, bilateral
E08.3519	Diabetes mellitus due to underlying condition with proliferative diabetic retinopathy with macular edema, unspecified eye
E09.311	Drug or chemical induced diabetes mellitus with unspecified diabetic retinopathy with macular edema
E09.3211	Drug or chemical induced diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, right eye
E09.3212	Drug or chemical induced diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, left eye
E09.3213	Drug or chemical induced diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, bilateral
E09.3219	Drug or chemical induced diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, unspecified eye
E09.3311	Drug or chemical induced diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, right eye
E09.3312	Drug or chemical induced diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, left eye
E09.3313	Drug or chemical induced diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, bilateral
E09.3319	Drug or chemical induced diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, unspecified eye
E09.3411	Drug or chemical induced diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, right eye
E09.3412	Drug or chemical induced diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, left eye
E09.3413	Drug or chemical induced diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, bilateral
E09.3419	Drug or chemical induced diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, unspecified eye
E09.3511	Drug or chemical induced diabetes mellitus with proliferative diabetic retinopathy with macular edema, right eye
E09.3512	Drug or chemical induced diabetes mellitus with proliferative diabetic retinopathy with macular edema, left eye
E09.3513	Drug or chemical induced diabetes mellitus with proliferative diabetic retinopathy with macular edema, bilateral
E09.3519	Drug or chemical induced diabetes mellitus with proliferative diabetic retinopathy with macular edema, unspecified eye
E10.311	Type 1 diabetes mellitus with unspecified diabetic retinopathy with macular edema
E10.3211	Type 1 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, right eye
E10.3212	Type 1 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, left eye
E10.3213	Type 1 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, bilateral
E10.3219	Type 1 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, unspecified eye
E10.3311	Type 1 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, right eye

ICD-10	ICD-10 Description
E10.3312	Type 1 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, left eye
E10.3313	Type 1 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, bilateral
E10.3319	Type 1 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, unspecified eye
E10.3411	Type 1 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, right eye
E10.3412	Type 1 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, left eye
E10.3413	Type 1 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, bilateral
E10.3419	Type 1 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, unspecified eye
E10.3511	Type 1 diabetes mellitus with proliferative diabetic retinopathy with macular edema, right eye
E10.3512	Type 1 diabetes mellitus with proliferative diabetic retinopathy with macular edema, left eye
E10.3513	Type 1 diabetes mellitus with proliferative diabetic retinopathy with macular edema, bilateral
E10.3519	Type 1 diabetes mellitus with proliferative diabetic retinopathy with macular edema, unspecified eye
E11.311	Type 2 diabetes mellitus with unspecified diabetic retinopathy with macular edema
E11.3211	Type 2 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, right eye
E11.3212	Type 2 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, left eye
E11.3213	Type 2 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, bilateral
E11.3219	Type 2 diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, unspecified eye
E11.3311	Type 2 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, right eye
E11.3312	Type 2 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, left eye
E11.3313	Type 2 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, bilateral
E11.3319	Type 2 diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, unspecified eye
E11.3411	Type 2 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, right eye
E11.3412	Type 2 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, left eye
E11.3413	Type 2 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, bilateral
E11.3419	Type 2 diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, unspecified eye
E11.3511	Type 2 diabetes mellitus with proliferative diabetic retinopathy with macular edema, right eye
E11.3512	Type 2 diabetes mellitus with proliferative diabetic retinopathy with macular edema, left eye
E11.3513	Type 2 diabetes mellitus with proliferative diabetic retinopathy with macular edema, bilateral
E11.3519	Type 2 diabetes mellitus with proliferative diabetic retinopathy with macular edema, unspecified eye
E13.311	Other specified diabetes mellitus with unspecified diabetic retinopathy with macular edema

ICD-10	ICD-10 Description
E13.3211	Other specified diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, right eye
E13.3212	Other specified diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, left eye
E13.3213	Other specified diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, bilateral
E13.3219	Other specified diabetes mellitus with mild nonproliferative diabetic retinopathy with macular edema, unspecified eye
E13.3311	Other specified diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, right eye
E13.3312	Other specified diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, left eye
E13.3313	Other specified diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, bilateral
E13.3319	Other specified diabetes mellitus with moderate nonproliferative diabetic retinopathy with macular edema, unspecified eye
E13.3411	Other specified diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, right eye
E13.3412	Other specified diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, left eye
E13.3413	Other specified diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, bilateral
E13.3419	Other specified diabetes mellitus with severe nonproliferative diabetic retinopathy with macular edema, unspecified eye
E13.3511	Other specified diabetes mellitus with proliferative diabetic retinopathy with macular edema, right eye
E13.3512	Other specified diabetes mellitus with proliferative diabetic retinopathy with macular edema, left eye
E13.3513	Other specified diabetes mellitus with proliferative diabetic retinopathy with macular edema, bilateral
E13.3519	Other specified diabetes mellitus with proliferative diabetic retinopathy with macular edema, unspecified eye
H35.3210	Exudative age-related macular degeneration, right eye, stage unspecified
H35.3211	Exudative age-related macular degeneration, right eye, with active choroidal neovascularization
H35.3212	Exudative age-related macular degeneration, right eye, with inactive choroidal neovascularization
H35.3213	Exudative age-related macular degeneration, right eye, with inactive scar
H35.3220	Exudative age-related macular degeneration, left eye, stage unspecified
H35.3221	Exudative age-related macular degeneration, left eye, with active choroidal neovascularization
H35.3222	Exudative age-related macular degeneration, left eye, with inactive choroidal neovascularization
H35.3223	Exudative age-related macular degeneration, left eye, with inactive scar
H35.3230	Exudative age-related macular degeneration, bilateral, stage unspecified
H35.3231	Exudative age-related macular degeneration, bilateral, with active choroidal neovascularization
H35.3232	Exudative age-related macular degeneration, bilateral, with inactive choroidal neovascularization
H35.3233	Exudative age-related macular degeneration, bilateral, with inactive scar
H35.3290	Exudative age-related macular degeneration, unspecified eye, stage unspecified

ICD-10	ICD-10 Description
H35.3291	Exudative age-related macular degeneration, unspecified eye, with active choroidal neovascularization
H35.3292	Exudative age-related macular degeneration, unspecified eye, with inactive choroidal neovascularization
H35.3293	Exudative age-related macular degeneration, unspecified eye, with inactive scar

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC