

## Aduhelm™ (aducanumab-avwa) (Intravenous)

Document Number: IC-0610

Last Review Date: 06/06/2022

Date of Origin: 07/01/2021

Dates Reviewed: 07/2021, 06/2022

### I. Length of Authorization

- Coverage will be provided for 6 months and may be renewed.

### II. Dosing Limits

#### A. Quantity Limit (max daily dose) [NDC Unit]:

- Aduhelm 170 mg/1.7 mL (100 mg/mL) solution in a single-dose vial: 1 vial every 28 days
- Aduhelm 300 mg/3 mL (100 mg/mL) solution in a single-dose vial: 4 vials every 28 days

#### B. Max Units (per dose and over time) [HCPCS Unit]:

- Infusions 1 & 2: 57 billable units (115 mg) every 28 days
- Infusions 3 & 4: 172 billable units (345 mg) every 28 days
- Infusions 5 & 6: 345 billable units (690 mg) every 28 days
- Infusion 7 and beyond: 575 billable units (1,150 mg) every 28 days

### III. Initial Approval Criteria <sup>1,5,6,9</sup>

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**
- Physician has assessed baseline disease severity utilizing an objective measure/tool (i.e., Mini-Mental Status Exam [MMSE], Alzheimer's Disease Assessment Scale-Cognitive Subscale [ADAS-Cog-13], Alzheimer's Disease Cooperative Study-Activities of Daily Living Inventory-Mild Cognitive Impairment version [ADCS-ADL-MCI], Clinical Dementia Rating-Sum of Boxes [CDR-SB], etc.); **AND**
- Patient does not have any of the following within 1 year of treatment initiation: pre-treatment localized superficial siderosis, 10 or more brain microhemorrhages, or brain hemorrhage >1 cm; **AND**

#### Universal Criteria <sup>1,5,6,9</sup>

- Must be prescribed by, or in consultation with, a specialist in neurology or gerontology; **AND**

- Patient has received a baseline brain magnetic resonance imaging (MRI) prior to initiating treatment (within one year prior – unless the patient has a more recent exacerbation, traumatic event [e.g., falls, etc.], or co-morbidity necessitating an evaluation within one month preceding initiation) and periodically throughout therapy (*see prescribing information for schedule of MRI scans*); **AND**
- Patient has not had a stroke or transient ischemic attack (TIA) or unexplained loss of consciousness in the past 12 months; **AND**
- Patient does not have any relevant brain hemorrhage, bleeding disorder, cerebrovascular abnormalities, or recent (within the prior year) cardiovascular condition (e.g., unstable angina, myocardial infarction, advanced CHF, or clinically significant conduction abnormalities); **AND**
- Patient does not have a clinically significant and unstable psychiatric illness in the past 6 months; **AND**
- Patient is not currently receiving anti-platelet agents (with the exception of prophylactic aspirin), anticoagulants (e.g., Factor Xa inhibitors), or anti-thrombins (e.g., heparin); **AND**
- Patient does not have a history of alcohol or substance abuse in the preceding year; **AND**

#### **Alzheimer’s Disease (AD) †<sup>1,2,5,6</sup>**

- Patient has mild cognitive impairment (MCI) due to AD or has mild Alzheimer’s dementia (there is insufficient evidence in moderate or severe AD) as evidenced by all of the following:
  - Clinical Dementia Rating (CDR)-Global Score of 0.5
  - Objective evidence of cognitive impairment at screening
  - MMSE score between 24 and 30 (inclusive)
  - Positron Emission Tomography (PET) scan is positive for amyloid beta plaque
- Other conditions mimicking, but of non-Alzheimer’s Dementia etiology, have been ruled out (e.g., vascular dementia, dementia with Lewy bodies [DLB], frontotemporal dementia [FTD], normal pressure hydrocephalus, etc.)

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); ◻ Orphan Drug

#### **IV. Renewal Criteria<sup>1,5,6</sup>**

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the universal and other indication-specific relevant criteria identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: amyloid related imaging abnormalities-edema (ARIA-E), severe hypersensitivity reactions, etc.; **AND**
- Patient has responded to therapy compared to pretreatment baseline as evidenced by improvement, stability, or slowing in cognitive and/or functional impairment in one or more

of the following (not all-inclusive): ADAS-Cog 13; ADCS-ADL-MCI; MMSE; CDR-SB, etc.; **AND**

- Patient has not progressed to moderate or severe AD; **AND**
- Patient has received a pre- 5<sup>th</sup>, 7<sup>th</sup>, 9<sup>th</sup>, AND 12<sup>th</sup> infusion MRI for monitoring of Amyloid Related Imaging Abnormalities-edema (ARIA-E) and Amyloid Related Imaging Abnormalities-hemosiderin (ARIA-H) microhemorrhages; **AND**

**ARIA-E**

- Patient is asymptomatic or mildly symptomatic with mild radiographic severity\* on MRI; **OR**
- Patient is asymptomatic or mildly symptomatic with moderate to severe radiographic severity\* on MRI AND administration will be suspended until MRI demonstrates radiographic resolution and symptoms, if present, resolve; **OR**
- Patient has moderate to severe symptoms with mild to severe radiographic severity\* on MRI AND administration will be suspended until MRI demonstrates radiographic resolution and symptoms, if present, resolve

**ARIA-H**

- Patient is asymptomatic with mild radiographic severity\* on MRI; **OR**
- Patient is asymptomatic with moderate radiographic severity\* on MRI AND administration will be suspended until MRI demonstrates radiographic resolution and symptoms, if present, resolve; **OR**
- Patient is symptomatic with mild to moderate radiographic severity\* on MRI AND administration will be suspended until MRI demonstrates radiographic resolution and symptoms, if present, resolve; **OR**
- Patient has severe radiographic severity\* on MRI AND administration will be suspended until MRI demonstrates radiographic resolution and symptoms, if present, resolve; **AND**
  - Clinical judgment will be used in considering whether to continue treatment or permanently discontinue
- Patient must continue maintenance therapy at the recommended dosage of 10 mg/kg every four weeks (*Note: clinical efficacy was demonstrated only at the highest dose, therefore doses below 10 mg/kg are not supported and will not be approved*)

*ARIA Type <sup>1</sup>	Radiographic Severity		
	Mild	Moderate	Severe
ARIA-E	FLAIR hyperintensity confined to sulcus and or cortex/subcortical white matter in one location < 5 cm	FLAIR hyperintensity 5 to 10 cm, or more than 1 site of involvement, each measuring < 10 cm	FLAIR hyperintensity measuring > 10 cm, often with significant subcortical white matter and/or sulcal involvement. One or more separate sites of involvement may be noted

ARIA-H microhemorrhage	≤ 4 new incident microhemorrhages	5 to 9 new incident microhemorrhages	10 or more new incident microhemorrhages
ARIA-H superficial siderosis	1 focal area of superficial siderosis	2 focal areas of superficial siderosis	> 2 focal areas of superficial siderosis

## V. Dosage/Administration <sup>1</sup>

Indication	Dose										
Alzheimer's Disease (AD)	<ul style="list-style-type: none"> <li>After an initial titration (see dosing schedule below), the recommended dosage of Aduhelm is 10 mg/kg and administered as an intravenous (IV) infusion over approximately one hour every four weeks and at least 21 days apart.</li> </ul>										
	<table border="1"> <thead> <tr> <th>IV Infusion (every 4 weeks)</th> <th>Aduhelm Dosage</th> </tr> </thead> <tbody> <tr> <td>Infusions 1 &amp; 2</td> <td>1 mg/kg</td> </tr> <tr> <td>Infusions 3 &amp; 4</td> <td>3 mg/kg</td> </tr> <tr> <td>Infusions 5 &amp; 6</td> <td>6 mg/kg</td> </tr> <tr> <td>Infusion 7 &amp; beyond</td> <td>10 mg/kg</td> </tr> </tbody> </table>	IV Infusion (every 4 weeks)	Aduhelm Dosage	Infusions 1 & 2	1 mg/kg	Infusions 3 & 4	3 mg/kg	Infusions 5 & 6	6 mg/kg	Infusion 7 & beyond	10 mg/kg
	IV Infusion (every 4 weeks)	Aduhelm Dosage									
	Infusions 1 & 2	1 mg/kg									
	Infusions 3 & 4	3 mg/kg									
Infusions 5 & 6	6 mg/kg										
Infusion 7 & beyond	10 mg/kg										
<ul style="list-style-type: none"> <li>Obtain recent (within one year) brain magnetic resonance imaging (MRI) prior to initiating treatment. Obtain MRIs prior to the 5<sup>th</sup> infusion (first dose of 6 mg/kg), 7<sup>th</sup> infusion (first dose of 10 mg/kg), 9<sup>th</sup> infusion (third dose of 10 mg/kg), and 12<sup>th</sup> infusion (sixth dose of 10 mg/kg).</li> </ul>											
<ul style="list-style-type: none"> <li>If an infusion is missed, resume administration at the same dose as soon as possible, infusions are to be administered every 4 weeks and at least 21 days apart.</li> </ul>											

## VI. Billing Code/Availability Information

### HCPCS Code:

- J0172 – Injection, aducanumab-avwa, 2 mg; 1 billable unit = 2 mg

### NDC:

- Aduhelm 170 mg/1.7 mL (100 mg/mL) solution in a single-dose vial: 64406-0101-xx
- Aduhelm 300 mg/3 mL (100 mg/mL) solution in a single-dose vial: 64406-0102-xx

## VII. References

- Aduhelm [package insert]. Cambridge, MA; Biogen, Inc; April 2022. Accessed May 2022.
- McKhann GM, Knopman DS, Chertklow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement.* 2011;7(3):263. Epub 2011 Apr 21.
- Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement.* 2011;7(3):280. Epub 2011 Apr 21.

4. Sevigny J, Chiao P, Bussière T, et al. The antibody aducanumab reduces Aβ plaques in Alzheimer's disease. *Nature*. 2016 Sep 1;537(7618):50-6. doi: 10.1038/nature19323. Update in: *Nature*. 2017 Jun 21;546(7659):564.
5. Biogen. 221AD302 Phase 3 Study of Aducanumab (BIIB037) in Early Alzheimer's Disease (**EMERGE**). Available from: <https://clinicaltrials.gov/ct2/show/NCT02484547?term=NCT02484547&draw=2&rank=1>. Accessed June 9, 2021.
6. Biogen. 221AD301 Phase 3 Study of Aducanumab (BIIB037) in Early Alzheimer's Disease (**ENGAGE**). Available from: <https://clinicaltrials.gov/ct2/show/NCT02477800?term=NCT02477800&draw=2&rank=1>. Accessed June 9, 2021.
7. O'Bryant SE, Lacritz LH, Hall J, et al. Validation of the new interpretive guidelines for the clinical dementia rating scale sum of boxes score in the national Alzheimer's coordinating center. *Arch Neurol*. 2010 Jun;67(6):746-9. doi: 10.1001/archneurol.2010.115.
8. Skinner J, Carvalho, JO, Potter GG, et al. The Alzheimer's Disease Assessment Scale-Cognitive-Plus (ADAS-Cog-Plus): an expansion of the ADAS-Cog to improve responsiveness in MCI. *Brain Imaging Behav*. 2012 Dec;6(4):489-501. doi: 10.1007/s11682-012-9166-3.
9. Salloway S, Chalkias S, Barkhof F, et al. Amyloid-Related Imaging Abnormalities in 2 Phase 3 Studies Evaluating Aducanumab in Patients With Early Alzheimer Disease. *JAMA Neurol*. 2022 Jan 1;79(1):13-21. doi: 10.1001/jamaneurol.2021.4161.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
G30.0	Alzheimer's disease with early onset
G30.1	Alzheimer's disease with late onset
G30.9	Alzheimer's disease, unspecified
G31.84	Mild cognitive impairment, so stated

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Articles (LCAs), and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at:

<https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCA/LCD): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC

### Medicare Part B Administrative Contractor (MAC) Jurisdictions

Jurisdiction	Applicable State/US Territory	Contractor
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC