

## Pemetrexed:

### Alimta®; Pemfexy™; Pemetrexed™ (Intravenous)

Document Number: IC-0007

Last Review Date: 12/01/2022

Date of Origin: 07/20/2010

Dates Reviewed: 09/2010, 12/2010, 03/2011, 06/2011, 09/2011, 12/2011, 03/2012, 06/2012, 09/2012, 12/2012, 03/2013, 06/2013, 09/2013, 12/2013, 03/2014, 06/2014, 09/2014, 12/2014, 03/2015, 05/2015, 08/2015, 11/2015, 02/2016, 05/2016, 08/2016, 11/2016, 02/2017, 05/2017, 08/2017, 11/2017, 02/2018, 05/2018, 09/2018, 12/2018, 03/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020, 09/2020, 12/2020, 03/2021, 06/2021, 09/2021, 12/2021, 03/2022, 06/2022, 09/2022, 12/2022

#### I. Length of Authorization <sup>15,26,28-30</sup>

Coverage will be provided for 6 months and may be renewed unless otherwise specified.

- Thymomas/Thymic Carcinoma: Coverage will be provided for six 21-day cycles and may not be renewed.
- MPeM and MPM: Coverage will be provided for six 21-day cycles and may not be renewed when used in combination with platinum therapy and bevacizumab.

#### II. Dosing Limits

##### A. Quantity Limit (max daily dose) [NDC Unit]:

- Alimta 100 mg powder for injection in a single-use vial: 4 vials every 21 days
- Alimta 500 mg powder for injection in a single-use vial: 4 vials every 21 days
- Pemfexy 500 mg solution for injection in a multi-dose vial: 4 vials every 21 days
- Pemetrexed disodium 750mg powder for injection: 2 vials every 21 days
- Pemetrexed disodium 1000mg powder for injection: 2 vials every 21 days
- Pemetrexed disodium 100mg/4 mL solution for injection: 4 vials every 21 days
- Pemetrexed disodium 500mg/20 mL solution for injection: 4 vials every 21 days
- Pemetrexed disodium 1000mg/40 mL solution for injection: 2 vials every 21 days

##### B. Max Units (per dose and over time) [HCPCS Unit]:

- CNS Lymphoma and Ovarian Cancer: 230 billable units every 21 days
- All other indications: 130 billable units every 21 days

### III. Initial Approval Criteria <sup>1,2</sup>

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

#### **Primary Central Nervous System (CNS) Lymphoma ‡ <sup>3,16,27</sup>**

- Used as single-agent; **AND**
  - Used as induction therapy in patients unsuitable for or intolerant to high-dose methotrexate (MTX); **OR**
  - Used for relapsed or refractory disease

#### **Malignant Peritoneal\* Mesothelioma (MPeM) ‡ <sup>3,29</sup>**

- Used as first-line therapy; **AND**
  - Used in combination with bevacizumab and cisplatin followed by single-agent maintenance bevacizumab (preferred) as first-line systemic therapy for unresectable disease; **OR**
  - Used as a single agent **OR** in combination with cisplatin or carboplatin (if cisplatin ineligible) for diffuse or recurrent disease; **OR**
- Used as subsequent therapy; **AND**
  - Used in combination with cisplatin or carboplatin (if cisplatin ineligible), with or without bevacizumab, if immunotherapy was administered as first-line treatment; **OR**
  - Used as a single agent; **AND**
    - Pemetrexed was not administered first-line; **OR**
    - Used as rechallenge if pemetrexed was administered first-line with a good sustained response at the time initial chemotherapy was interrupted

#### **Malignant Pleural\* Mesothelioma (MPM) † <sup>1-6,10,26</sup>**

- Used as induction therapy; **AND**
  - Used in combination with cisplatin or carboplatin (if cisplatin ineligible) in patients with epithelioid histology; **OR**
- Used as first-line therapy; **AND**
  - Used in combination with bevacizumab and cisplatin followed by single-agent maintenance bevacizumab (preferred) as first-line systemic therapy ; **OR**
  - Used as a single agent **OR** in combination with cisplatin or carboplatin (if cisplatin ineligible) for resected or recurrent disease; **OR**
- Used as subsequent therapy; **AND**
  - Used in combination with cisplatin or carboplatin (if cisplatin ineligible), with or without bevacizumab, if immunotherapy was administered as first-line treatment; **OR**
  - Used as a single agent; **AND**

**PEMETREXED (Alimta®; Pemfexy™; Pemetrexed)**  
**Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

- Pemetrexed was not administered first-line; **OR**
- Used as rechallenge if pemetrexed was administered first-line with a good sustained response at the time initial chemotherapy was interrupted

*\* Note: May also be used for pericardial mesothelioma and tunica vaginalis testis mesothelioma.*

### Non-Squamous Non-Small Cell Lung Cancer (NS-NSCLC) † 1-3,7-9,11,12,28,30

- Used in combination with carboplatin or cisplatin-containing regimen; **OR**
- Used as single-agent therapy; **AND**
  - Patient has recurrent, advanced, or metastatic disease (excluding locoregional recurrence or symptomatic local disease without evidence of disseminated disease) or mediastinal lymph node recurrence with prior radiation therapy; **AND**
    - Used as first-line therapy for PD-L1  $\geq 1\%$  tumors that have negative actionable molecular biomarkers\*; **OR**
    - Used as first-line therapy for PD-L1  $< 1\%$  and tumors that have negative actionable molecular markers \* OR BRAF V600E-mutation, NTRK1/2/3 gene fusion, MET exon-14 skipping mutation, EGFR exon 20 mutation, KRAS G12C mutation, or RET rearrangement positive tumors; **OR**
    - Used as subsequent therapy for first progression after initial systemic therapy; **OR**
    - Used continuation or switch maintenance therapy in patients who have achieved tumor response or stable disease following initial therapy (*Note: Continuation maintenance therapy may also be given in combination with bevacizumab or pembrolizumab*)

*\* Note: Actionable molecular genomic biomarkers include EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET exon 14 skipping mutation, RET rearrangement, and ERBB2 (HER2). If there is insufficient tissue to allow testing for all of EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, and ERBB2 (HER2) repeat biopsy and/or plasma testing should be done. If these are not feasible, treatment is guided by available results and, if unknown, these patients are treated as though they do not have driver oncogenes.*

### Thymomas/Thymic Carcinoma ‡ 3,14,15,25

- Used as a single agent; **AND**
  - Used as first line therapy or postoperative treatment in patients who are unable to tolerate first-line combination regimens; **OR**
  - Used as second-line therapy for unresectable or metastatic disease

### Ovarian Cancer (Epithelial Ovarian/Fallopian Tube/Primary Peritoneal Cancer) ‡ 3,13,24

- Used as single-agent therapy; **AND**
  - Patient has recurrent or persistent disease; **AND**
    - Patient is not experiencing an immediate biochemical relapse (i.e., rising CA-125 without radiographic evidence of disease); **OR**
  - Patient has recurrent low-grade serous carcinoma

**PEMETREXED (Alimta®; Pefexy™; Pemetrexed)**  
**Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.  
 ©2022, Magellan Rx Management

§ Genomic Aberration/Mutational Driver Targeted Therapies (Note: not all inclusive, refer to guidelines for appropriate use)					
Sensitizing EGFR mutation-positive tumors	ALK rearrangement-positive tumors	ROS1 rearrangement-positive tumors	BRAF V600E-mutation positive tumors	NTRK1/2/3 gene fusion positive tumors	ERBB2 (HER2) mutation positive tumors
<ul style="list-style-type: none"> <li>- Afatinib</li> <li>- Erlotinib</li> <li>- Dacomitinib</li> <li>- Gefitinib</li> <li>- Osimertinib</li> <li>- Amivantamab (exon-20 insertion)</li> <li>- Mobocertinib (exon-20 insertion)</li> </ul>	<ul style="list-style-type: none"> <li>- Alectinib</li> <li>- Brigatinib</li> <li>- Ceritinib</li> <li>- Crizotinib</li> <li>- Lorlatinib</li> </ul>	<ul style="list-style-type: none"> <li>- Ceritinib</li> <li>- Crizotinib</li> <li>- Entrectinib</li> <li>- Lorlatinib</li> </ul>	<ul style="list-style-type: none"> <li>- Dabrafenib ± trametinib</li> <li>- Vemurafenib</li> </ul>	<ul style="list-style-type: none"> <li>- Larotrectinib</li> <li>- Entrectinib</li> </ul>	<ul style="list-style-type: none"> <li>- Fam-trastuzumab deruxtecan-nxki</li> <li>- Ado-trastuzumab emtansine</li> </ul>
PD-L1 tumor expression ≥ 1%	PD-L1 tumor expression ≥ 50%	MET exon-14 skipping mutations	RET rearrangement-positive tumors	KRAS G12C mutation positive tumors	
<ul style="list-style-type: none"> <li>- Pembrolizumab</li> <li>- Atezolizumab</li> <li>- Nivolumab + ipilimumab</li> </ul>	<ul style="list-style-type: none"> <li>- Pembrolizumab</li> <li>- Atezolizumab</li> <li>- Nivolumab + ipilimumab</li> <li>- Cemiplimab</li> </ul>	<ul style="list-style-type: none"> <li>- Capmatinib</li> <li>- Crizotinib</li> <li>- Tepotinib</li> </ul>	<ul style="list-style-type: none"> <li>- Selpercatinib</li> <li>- Cabozantinib</li> <li>- Pralsetinib</li> </ul>	<ul style="list-style-type: none"> <li>- Sotorasib</li> </ul>	

#### IV. Renewal Criteria <sup>1,2</sup>

Coverage can be renewed based upon the following criteria:

- Patient continues to meet indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: myelosuppression (e.g., neutropenia, febrile neutropenia, thrombocytopenia, anemia), renal toxicity (CrCl < 45 mL/min), bullous and exfoliative skin toxicity (e.g., Stevens-Johnson Syndrome/Toxic epidermal necrolysis), interstitial pneumonitis, radiation recall, etc.; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**

#### MPeM and MPM <sup>26,29</sup>

- May not be renewed when used in combination with platinum therapy and bevacizumab

#### Thymomas/Thymic Carcinoma <sup>15</sup>

- May not be renewed

#### V. Dosage/Administration <sup>1,2,13,15,16,26,28-30</sup>

Indication	Dose
------------	------

**PEMETREXED (Alimta®; Pefexy™; Pemetrexed)**  
**Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.  
 ©2022, Magellan Rx Management

Non-Squamous NSCLC	Administer up to 500 mg/m <sup>2</sup> intravenously every 21 days
MPM, MPeM	Administer 500 mg/m <sup>2</sup> intravenously every 21 days – For 6 cycles only when used in combination with platinum therapy and bevacizumab – All others until disease progression or unacceptable toxicity
Primary CNS Lymphoma, Ovarian Cancer	Administer 900 mg/m <sup>2</sup> intravenously every 21 days, until disease progression or unacceptable toxicity
Thymomas/Thymic Carcinoma	Administer 500 mg/m <sup>2</sup> intravenously every 21 days for a maximum of 6 cycles in absence of disease progression or unacceptable toxicity
<ul style="list-style-type: none"> <li>• Supplement with oral folic acid and intramuscular vitamin B<sub>12</sub></li> <li>• Avoid administration of ibuprofen for 2 days before, the day of, and 2 days following administration in patients with CrCl &lt;80 mL/min.</li> <li>• Do not dose in patients with CrCl &lt;45 mL/min</li> </ul>	

## VI. Billing Code/Availability Information

Product Formulation	Drug	Manufacturer	Type	HCPCS Code	NDC
Pemetrexed Disodium Lyophilisate for inj.	Alimta 100 mg powder for inj. SDV *	Lilly	Brand	J9305	00002-7640-xx
	Alimta 500 mg powder for inj. SDV *				00002-7623-xx
	Pemetrexed 100 mg powder for inj. Ψ	Hospira	Brand	J9305	00409-1060-xx
	Pemetrexed 500 mg powder for inj. Ψ				00409-1061-xx
	Pemetrexed 1000 mg powder for inj. Ψ				00409-1062-xx
Pemetrexed 750 mg powder for inj.*	N/A	Generic	J9305	N/A	
Pemetrexed 1000 mg powder for inj.*					
Pemetrexed Disodium Solution for injection	Pemetrexed 100mg/4mL inj. SDV Ψ	Sandoz	Brand	J9305	00781-3518-xx
		Accord			16729-0522-xx
	Pemetrexed 500mg/20mL inj. SDV Ψ	Hospira	Brand	J9305	00409-2188-xx
		Sandoz			00781-3519-xx
	Pemetrexed 1000mg/40mL inj. SDV Ψ	Accord	Brand	J9305	16729-0522-xx
		Hospira			00409-3532-xx
		Accord			16729-0522-xx
Pemetrexed Solution for injection	Pemfexy 500 mg/20 mL inj. MDV	Eagle.	Brand	J9304	42367-0531-xx
	Pemetrexed 100mg/4mL inj. SDV Ψ	Hospira	Brand	J9305	00409-1045-xx
		Teva		J9314	00480-4516-xx
	Pemetrexed 500mg/20mL inj. SDV Ψ	Teva	Brand	J9314	00480-4514-xx
Pemetrexed 1000mg/40mL inj. SDV Ψ	Teva	Brand	J9314	00480-4515-xx	
<b>*Multiple manufacturers produce ANDA generics</b>					
<b>Ψ Approved by the FDA as a 505(b)(2) NDA of the innovator product</b>					
J9304 – Injection, pemetrexed (pemfexy), 10 mg					
J9305 – Injection, pemetrexed, not otherwise specified, 10 mg					
J9314 – Injection, pemetrexed (teva) not therapeutically equivalent to J9305, 10 mg (Effective 1/1/2023)					

## VII. References

1. Alimta [package insert]. Indianapolis, IN; Eli Lilly; August 2022. Accessed October 2022.
2. Pemfexy [package insert]. Woodcliff Lake, NJ; Eagle Pharmaceuticals, Inc; June 2020. Accessed October 2022.

3. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for pemetrexed. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®, NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed October 2022.
4. Castagneto B, Botta M, Aitini E, et al, “Phase II Study of Pemetrexed in Combination With Carboplatin in Patients With Malignant Pleural Mesothelioma (MPM),” *Ann Oncol*, 2008, 19(2):370-3.
5. Ceresoli GL, Zucali PA, Favaretto AG, et al, “Phase II Study of Pemetrexed plus Carboplatin in Malignant Pleural Mesothelioma,” *J Clin Oncol*, 2006, 24(9):1443-8.
6. Taylor P, Castagneto B, Dark G, et al, “Single-Agent Pemetrexed for Chemonaïve and Pretreated Patients With Malignant Pleural Mesothelioma: Results of an International Expanded Access Program,” *J Thorac Oncol*, 2008, 3(7):764-71
7. Ciuleanu T, Brodowicz T, Zielinski C, et al, “Maintenance Pemetrexed Plus Best Supportive Care versus Placebo Plus Best Supportive Care for Non-Small-Cell Lung Cancer: A Randomised, Double-Blind, Phase 3 Study,” *Lancet*, 2009, 374(9699):1432-40.
8. Grønberg BH, Bremnes RM, Fløtten O, et al, “Phase III Study by the Norwegian Lung Cancer Study Group: Pemetrexed Plus Carboplatin Compared With Gemcitabine Plus Carboplatin as First-Line Chemotherapy in Advanced Non-Small-Cell Lung Cancer,” *J Clin Oncol*, 2009, 27(19):3217-24.
9. Hanna N, Shepherd FA, Fossella FV, et al, “Randomized Phase III Trial of Pemetrexed versus Docetaxel in Patients With Non-Small-Cell Lung Cancer Previously Treated With Chemotherapy,” *J Clin Oncol*, 2004, 22(9):1589-97.
10. Jassem J, Ramlau R, Santoro A, et al, “Phase III Trial of Pemetrexed Plus Best Supportive Care Compared With Best Supportive Care in Previously Treated Patients With Advanced Malignant Pleural Mesothelioma,” *J Clin Oncol*, 2008, 26(10):1698-704.
11. Scagliotti GV, Parikh P, von Pawel J, et al, “Phase III Study Comparing Cisplatin Plus Gemcitabine With Cisplatin Plus Pemetrexed in Chemotherapy-Naïve Patients With Advanced-Stage Non-Small-Cell Lung Cancer,” *J Clin Oncol*, 2008, 26(21):3543-51.
12. Langer CJ, Gadgeel SM, Borghaei H, et al. Carboplatin and pemetrexed with or without pembrolizumab for advanced, non-squamous non-small-cell lung cancer: a randomised, phase 2 cohort of the open-label KEYNOTE-021 study. *Lancet Oncol*. 2016;17(11):1497-1508.
13. Miller DS, Blessing JA, Krasner CN, et al. Phase II Evaluation of Pemetrexed in the Treatment of Recurrent or Persistent Platinum-Resistant Ovarian or Primary Peritoneal Carcinoma: A Study of the Gynecologic Oncology Group. *J Clin Oncol*, 2009, 27(16):2686-91.
14. Liang Y, Padda SK, Riess JW, et al. Pemetrexed in patients with thymic malignancies previously treated with chemotherapy. *Lung Cancer*. 2015 Jan;87(1):34-8.

15. Gbolahan OB, Porter RF, Salter JT, et al. A Phase II Study of Pemetrexed in Patients with Recurrent Thymoma and Thymic Carcinoma. *J Thorac Oncol*. 2018 Dec;13(12):1940-1948.
16. Raizer JJ, Rademaker A, Evens AM, et al. Pemetrexed in the treatment of relapsed/refractory primary central nervous system lymphoma. *Cancer*. 2012 Aug 1;118(15):3743-8.
17. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. *J Oncol Pract*. 2018 Mar;14(3):e130-e136.
18. Hematology/Oncology Pharmacy Association (2019). Intravenous Cancer Drug Waste Issue Brief. Retrieved from [http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug\\_Waste\\_2019.pdf](http://www.hoparx.org/images/hopa/advocacy/Issue-Briefs/Drug_Waste_2019.pdf)
19. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. *BMJ*. 2016 Feb 29;352:i788.
20. Gandhi L, Rodríguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer. *N Engl J Med*. 2018;378(22):2078-2092. doi:10.1056/NEJMoa1801005.
21. Wu YL, Lu S, Cheng Y, et al. Efficacy and safety of pemetrexed/cisplatin versus gemcitabine/cisplatin as first-line treatment in Chinese patients with advanced nonsquamous non-small cell lung cancer. *Lung Cancer*. 2014;85(3):401-407. doi:10.1016/j.lungcan.2014.07.007.
22. Paz-Ares L, de Marinis F, Dediu M, et al. Maintenance therapy with pemetrexed plus best supportive care versus placebo plus best supportive care after induction therapy with pemetrexed plus cisplatin for advanced non-squamous non-small-cell lung cancer (PARAMOUNT): a double-blind, phase 3, randomised controlled trial. *Lancet Oncol*. 2012;13(3):247-255. doi:10.1016/S1470-2045(12)70063-3.
23. Vogelzang NJ, Rusthoven JJ, Symanowski J, et al. Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. *J Clin Oncol*. 2003;21(14):2636-2644. doi:10.1200/JCO.2003.11.136.
24. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer Version 5.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to [NCCN.org](http://NCCN.org). Accessed October 2022.
25. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Thymomas and Thymic Carcinomas Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National

Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2022.

26. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Malignant Pleural Mesothelioma Version 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2022.
27. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Central Nervous System Cancers Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2022.
28. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Non-Small Cell Lung Cancer Version 5.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2022.
29. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Malignant Peritoneal Mesothelioma Version 2.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed October 2022.
30. Forde P, Spicer J, Provencio M, et.al. Abstract CT003: Nivolumab (NIVO) + platinum-doublet chemotherapy (chemo) vs chemo as neoadjuvant treatment (tx) for resectable (IB-III A) non-small cell lung cancer (NSCLC) in the phase 3 CheckMate 816 trial. Cancer Res (2021) 81 (13\_Supplement): CT003.<https://doi.org/10.1158/1538-7445.AM2021-CT003>.

## Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C33	Malignant neoplasm of trachea
C34.00	Malignant neoplasm of unspecified main bronchus
C34.01	Malignant neoplasm of right main bronchus
C34.02	Malignant neoplasm of left main bronchus
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung

**PEMETREXED (Alimta®; Pefexy™; Pemetrexed)**  
**Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management



ICD-10	ICD-10 Description
C34.12	Malignant neoplasm of upper lobe, left bronchus or lung
C34.2	Malignant neoplasm of middle lobe, bronchus or lung
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus or lung
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung
C37	Malignant neoplasm of thymus
C45.0	Mesothelioma of pleura
C45.1	Mesothelioma of peritoneum
C45.2	Mesothelioma of pericardium
C45.7	Mesothelioma of other sites
C45.9	Mesothelioma, unspecified
C48.1	Malignant neoplasm of specified parts of peritoneum
C48.2	Malignant neoplasm of peritoneum, unspecified
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum
C56.1	Malignant neoplasm of right ovary
C56.2	Malignant neoplasm of left ovary
C56.3	Malignant neoplasm of bilateral ovaries
C56.9	Malignant neoplasm of unspecified ovary
C57.00	Malignant neoplasm of unspecified fallopian tube
C57.01	Malignant neoplasm of right fallopian tube
C57.02	Malignant neoplasm of left fallopian tube
C57.10	Malignant neoplasm of unspecified broad ligament
C57.11	Malignant neoplasm of right broad ligament
C57.12	Malignant neoplasm of left broad ligament
C57.20	Malignant neoplasm of unspecified round ligament
C57.21	Malignant neoplasm of right round ligament
C57.22	Malignant neoplasm of left round ligament
C57.3	Malignant neoplasm of parametrium

**PEMETREXED (Alimta®; Pefexy™; Pemetrexed)  
Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

ICD-10	ICD-10 Description
C57.4	Malignant neoplasm of uterine adnexa, unspecified
C57.7	Malignant neoplasm of other specified female genital organs
C57.8	Malignant neoplasm of overlapping sites of female genital organs
C57.9	Malignant neoplasm of female genital organ, unspecified
C83.30	Diffuse large B-cell lymphoma unspecified site
C83.39	Diffuse large B-cell lymphoma extranodal and solid organ sites
C83.80	Other non-follicular lymphoma, unspecified site
C83.89	Other non-follicular lymphoma, extranodal and solid organ sites
C85.89	Other specified types of non-Hodgkin lymphoma, extranodal and solid organ sites
C85.99	Non-Hodgkin's lymphoma extranodal and solid organ sites
D15.0	Benign neoplasm of thymus
Z85.118	Personal history of other malignant neoplasm of bronchus and lung
Z85.43	Personal history of malignant neoplasm of ovary

## Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT,	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)

**PEMETREXED (Alimta®; Pemfexy™; Pemetrexed)  
Prior Auth Criteria**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2022, Magellan Rx Management

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
15	KY, OH	CGS Administrators, LLC