

Kyprolis® (carfilzomib) (Intravenous)

Document Number: IC-0157

Last Review Date: 06/06/2022

Date of Origin: 02/07/2013

Dates Reviewed: 12/2013, 02/2014, 06/2014, 09/2014, 12/2014, 05/2015, 08/2015, 11/2015, 02/2016, 05/2016, 08/2016, 11/2016, 02/2017, 05/2017, 08/2017, 11/2017, 02/2018, 05/2018, 09/2018, 12/2018, 03/2019, 06/2019, 09/2019, 12/2019, 03/2020, 06/2020, 09/2020, 12/2020, 03/2021, 06/2021, 09/2021, 12/2021, 03/2022, 06/2022

I. Length of Authorization ^{1,5,12,21,27}

Coverage will be provided for 6 months and may be renewed.

Multiple Myeloma

- Combination therapy with lenalidomide and dexamethasone is limited to eighteen (18) 28-day treatment cycles.
- Combination therapy with cyclophosphamide, thalidomide, and dexamethasone is limited to twelve (12) 28-day treatment cycles.
- Combination therapy with daratumumab, lenalidomide, and dexamethasone is limited to eight (8) 28-day treatment cycles.

Waldenström's Macroglobulinemia/Lymphoplasmacytic Lymphoma

- Treatment is limited to six (6) 21-day induction therapy treatment cycles and eight (8) 56-day maintenance therapy treatment cycles.

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Kyprolis 10 mg single-dose vial: 2 vials per 28-day cycle
- Kyprolis 30 mg single-dose vial: 1 vial per 28-day cycle
- Kyprolis 60 mg single-dose vial: 12 vials per 28-day cycle

B. Max Units (per dose and over time) [HCPCS Unit]:

- **Multiple Myeloma**
 - 720 billable units every 28 days
- **Systemic Light Chain Amyloidosis**

- 360 billable units every 28 days
- **Waldenström's Macroglobulinemia/Lymphoplasmacytic Lymphoma**
 - 320 billable units every 21 days

III. Initial Approval Criteria ¹

Coverage is provided in the following conditions:

- Patient is at least 18 years of age; **AND**

Multiple Myeloma † ⊕ 1,2,10,11,13-17,19,23,32

- Used as primary therapy for symptomatic disease; **AND**
 - Used in combination with daratumumab, lenalidomide, and dexamethasone (*transplant candidates ONLY*); **OR**
 - Used in combination with lenalidomide and dexamethasone; **OR**
 - Used in combination with dexamethasone and cyclophosphamide; **OR**
- Used for disease relapse after 6 months following primary induction therapy with the same regimen; **AND**
 - Used in combination with lenalidomide and dexamethasone; **OR**
 - Used in combination with dexamethasone and cyclophosphamide; **OR**
- Used for previously treated relapsed, progressive, or refractory disease; **AND**
 - Used as a single agent †; **OR**
 - Used in combination with dexamethasone with or without lenalidomide †; **OR**
 - Used in combination with dexamethasone and daratumumab †; **OR**
 - Used in combination with dexamethasone and daratumumab and hyaluronidase-fihj †; **OR**
 - Used in combination with dexamethasone and cyclophosphamide with or without thalidomide; **OR**
 - Used in combination with dexamethasone and isatuximab-irfc; **OR**
 - Used in combination with dexamethasone and selinexor; **OR**
 - Used in combination with pomalidomide and dexamethasone; **AND**
 - Patient has received at least two (2) prior therapies, including a proteasome inhibitor [i.e., bortezomib, etc.] and an immunomodulatory agent [i.e., lenalidomide, thalidomide, etc.]; **AND**
 - Disease has progressed on or within 60 days of completion of the last therapy

Waldenström's Macroglobulinemia/Lymphoplasmacytic Lymphoma † 2,5,18

- Used in combination with rituximab and dexamethasone (CaRD regimen); **AND**
 - Used as primary therapy; **OR**
 - Used for relapsed disease; **AND**

- CaRD regimen was previously used as primary therapy; **AND**
- Patient had a prolonged response (i.e., 24 months) to CaRD therapy

Systemic Light Chain Amyloidosis †^{2,30,31}

- Patient has relapsed or refractory non-cardiac disease; **AND**
 - Used as a single agent; **OR**
 - Used in combination with dexamethasone

† FDA Approved Indication(s); ‡ Compendia Approved Indication(s); ◊ Orphan Drug

IV. Renewal Criteria ^{1,2,6}

Coverage may be renewed based upon the following criteria:

- Patient continues to meet the indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: cardiac toxicity (including CHF, pulmonary edema, decreased ejection fraction), pulmonary toxicity (including acute respiratory distress syndrome (ARDS), acute respiratory failure), pulmonary hypertension, dyspnea, severe infusion-related reactions, tumor lysis syndrome (TLS), thrombocytopenia, hepatic toxicity/failure, thrombotic microangiopathy (including thrombotic thrombocytopenic purpura/hemolytic uremic syndrome [TTP/HUS]), acute renal failure, severe hypertension, posterior reversible encephalopathy syndrome (PRES), venous thromboembolic events (including deep venous thrombosis, pulmonary embolism), hemorrhage, progressive multifocal leukoencephalopathy (PML), etc.; **AND**

Multiple Myeloma ^{1,12,27}

- Combination therapy with lenalidomide and dexamethasone may be renewed up to a maximum of eighteen (18) 28-day treatment cycles.
- Combination therapy with cyclophosphamide, thalidomide, and dexamethasone may be renewed up to a maximum of twelve (12) 28-day treatment cycles.
- Combination therapy with daratumumab, lenalidomide, and dexamethasone may be renewed up to a maximum of eight (8) 28-day treatment cycles.

Waldenström's Macroglobulinemia/Lymphoplasmacytic Lymphoma ^{5,21}

- May be renewed up to a maximum of six (6) 21-day induction therapy treatment cycles and eight (8) 56-day maintenance therapy treatment cycles.

V. Dosage/Administration ^{1,5,7,9,12,20-22,24-30,32-34}

Indication	Dose
Multiple Myeloma (primary therapy OR disease relapse ≥6 months following primary induction therapy with the same regimen)	<p><u>Combination with daratumumab, lenalidomide and dexamethasone (20/56 regimen)</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on day 1; if tolerated, increase to 56 mg/m² on days 8 and 15 of a 28-day treatment cycle – Cycles 2 through 8: 56 mg/m² on days 1, 8, and 15 of a 28-day treatment cycle <p><u>Combination with lenalidomide and dexamethasone (20/36 regimen)</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 36 mg/m² days 8, 9, 15, and 16 of a 28-day treatment cycle – Cycles 2 through 8: 36 mg/m² days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle – Cycles 9 through 18: 36 mg/m² days 1, 2, 15, and 16 of a 28-day treatment cycle <p><u>Combination with cyclophosphamide and dexamethasone</u></p> <p><u>20/36 regimen:</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 36 mg/m² days 8, 9, 15, and 16 of a 28-day treatment cycle – Cycles 2 through 9: 36 mg/m² days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle – Cycle 10 and beyond: 36 mg/m² on days 1, 2, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity <p><u>20/70 regimen:</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on day 1; if tolerated, increase to 70 mg/m² days 8 and 15 of a 28-day treatment cycle – Cycles 2 through 9: 70 mg/m² days 1, 8, and 15 of a 28-day treatment cycle – Cycle 10 and beyond: 70 mg/m² on days 1 and 15 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity
Multiple Myeloma (relapsed, progressive, or refractory disease)	<p><u>Single agent</u></p> <p><u>20/27 regimen:</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 27 mg/m² on days 8, 9, 15, and 16 of a 28-day treatment cycle – Cycles 2 through 12: 27 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle – Cycle 13 and beyond: 27 mg/m² on days 1, 2, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity <p><u>20/56 regimen:</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 56 mg/m² on days 8, 9, 15, and 16 of a 28-day treatment cycle. – Cycles 2 through 12: 56 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle – Cycle 13 and beyond: 56 mg/m² on days 1, 2, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity <p><u>Combination with lenalidomide and dexamethasone (20/27 regimen)</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 27 mg/m² on days 8, 9, 15, and 16 of a 28-day treatment cycle – Cycles 2 through 12: 27 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle – Cycles 13 through 18: 27 mg/m² on days 1, 2, 15, and 16 of a 28-day treatment cycle; beginning with cycle 19, lenalidomide and dexamethasone may be continued (until disease progression or unacceptable toxicity) without carfilzomib <p><u>Combination with dexamethasone</u></p> <p><u>20/56 regimen:</u></p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 56 mg/m² on days 8, 9, 15, and 16 of a 28-day treatment cycle – Cycle 2 and beyond: 56 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity

20/70 regimen:

- Cycle 1: 20 mg/m² on day 1; if tolerated, increase to 70 mg/m² on day 8 and 15 of a 28-day treatment cycle
- Cycle 2 and beyond: 70 mg/m² on days 1, 8, and 15 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity

Combination with daratumumab (or daratumumab and hyaluronidase-fihj) and dexamethasone

20/56 regimen:

- Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 56 mg/m² on days 8, 9, 15 and 16 of a 28-day treatment cycle
- Cycle 2 and beyond: 56 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity

20/70 regimen:

- Cycle 1: 20 mg/m² on day 1; if tolerated, increase to 70 mg/m² on day 8 and 15 of a 28-day treatment cycle
- Cycle 2 and beyond: 70 mg/m² on days 1, 8, and 15 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity

Combination with cyclophosphamide, thalidomide, and dexamethasone (20/36 regimen)

- Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 36 mg/m² days 8, 9, 15, and 16 of a 28-day treatment cycle
- Cycles 2 through 4: 36 mg/m² days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle
- Patients who achieve stable disease or better may continue treatment for up to 8 additional cycles

Combination with cyclophosphamide and dexamethasone (20/36 regimen)

Induction

- Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 36 mg/m² days 8, 9, 15, and 16 of a 28-day treatment cycle
- Cycles 2 through 6: 36 mg/m² days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle

Maintenance

- Cycles 7 through 12: 36 mg/m² on days 1, 2, 15, and 16 of a 28-day treatment cycle
- Cycle 13 and beyond: 36 mg/m² on days 1 and 2 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity

Combination with isatuximab-irfc and dexamethasone (20/56 regimen)

- Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 56 mg/m² on days 8, 9, 15 and 16 of a 28-day treatment cycle
- Cycle 2 and beyond: 56 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity

Combination with selinexor and dexamethasone (20/56 regimen)

- Cycle 1: 20 mg/m² on day 1; if tolerated, increase to 56 mg/m² on days 8 and 15 of a 28-day treatment cycle
- Cycle 2 and beyond: 56 mg/m² on days 1, 8, and 15 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity

Combination with pomalidomide and dexamethasone

20/27 regimen:

- Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 27 mg/m² on days 8, 9, 15, and 16 of a 28-day treatment cycle
- Cycles 2 through 6: 27 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle
- Cycle 7 and beyond: 27 mg/m² on days 1, 2, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity
- **NOTE:** If disease progression occurs while on maintenance dosing, resume full dosing of 27 mg/m² on days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle

20/36 regimen:

	<ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1 and 2; if tolerated, increase to 36 mg/m² days 8, 9, 15, and 16 of a 28-day treatment cycle – Cycles 2 through 8: 36 mg/m² days 1, 2, 8, 9, 15, and 16 of a 28-day treatment cycle – Cycle 9 and beyond: 36 mg/m² days 1, 2, 15, and 16 of a 28-day treatment cycle; continue until disease progression or unacceptable toxicity
Waldenström's Macroglobulinemia/ Lymphoplasmacytic Lymphoma	<p>CaRD regimen (carfilzomib, rituximab, dexamethasone)</p> <p>Induction</p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on days 1, 2, 8 and 9 of a 21-day treatment cycle – Cycles 2 through 6: 36 mg/m² on days 1, 2, 8 and 9 of a 21-day treatment; begin maintenance 8 weeks later <p>Maintenance</p> <ul style="list-style-type: none"> – 36 mg/m² on days 1 and 2 every 8 weeks for 8 cycles
Systemic Light Chain Amyloidosis	<p>Single agent or combination with dexamethasone</p> <ul style="list-style-type: none"> – Cycle 1: 20 mg/m² on day 1; if tolerated, increase to 27 mg/m² days 8 and 15 of a 28-day treatment cycle – Cycles 2 and beyond: Up to 56 mg/m² days 1, 8, and 15 of a 28-day treatment cycle
<p><i>Note: Calculate the Kyprolis dose using the patient's actual body surface area at baseline. In patients with a body surface area greater than 2.2 m², calculate the dose based upon a body surface area of 2.2 m².</i></p>	

VI. Billing Code/Availability Information

HCPCS Code:

- J9047 – Injection, carfilzomib, 1 mg; 1mg = 1 billable unit

NDC(s):

- Kyprolis 10 mg single-dose vial for injection: 76075-0103-xx
- Kyprolis 30 mg single-dose vial for injection: 76075-0102-xx
- Kyprolis 60 mg single-dose vial for injection: 76075-0101-xx

VII. References

1. Kyprolis [package insert]. Thousand Oaks, CA; Onyx Pharmaceuticals, Inc.; November 2021. Accessed May 2022.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) for Carfilzomib. National Comprehensive Cancer Network, 2022. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed May 2022.
3. BGM Durie, J-L Harousseau, J S Miguel, et al on behalf of the International Myeloma Working Group. International uniform response criteria for multiple myeloma. *Leukemia*. Sep; 20(9):1467-73.
4. Dimopoulos MA, Kastiris E, Owen RG, et al. Treatment recommendations for patients with Waldenström's macroglobulinemia (WM) and related disorders: IWWM-7 consensus. *Blood*. 2014; 124(9):1404–1411.

5. Treon SP, Tripsas CK, Meid K, et al. Carfilzomib, rituximab, and dexamethasone (CaRD) treatment offers a neuropathy-sparing approach for treating Waldenström's macroglobulinemia. *Blood*. 2014;124(4):503–510.
6. UpToDate. Hudson (OH): Lexicomp Inc.: Carfilzomib: Drug information. Topic 86042 Version 135.0, 2018 Accessed November 2018.
7. Shah JJ, Stadtmauer EA, Abonour R, et al. Carfilzomib, pomalidomide and dexamethasone for relapsed or refractory myeloma. *Blood* 2015; 126: 2284-2290.
8. Berdeja JG, Hart LL, Mace JR, et al. Phase I/II study of the combination of panobinostat and carfilzomib in patients with relapsed/refractory multiple myeloma. *Haematologica* 2015; 100: 670-676.
9. Bringhen S, Petrucci MT, Larocca A, et al. Carfilzomib, cyclophosphamide, and dexamethasone in patients with newly diagnosed multiple myeloma: a multicenter, phase 2 study. *Blood*. 2014 Jul 3;124(1):63-9.
10. Moreau P, Mateos MV, Berenson JR, et al. Once weekly versus twice weekly carfilzomib dosing in patients with relapsed and refractory multiple myeloma (A.R.R.O.W.): interim analysis results of a randomised, phase 3 study. *Lancet Oncol* 2018;19(7):953-964.
11. Chari A, Martinez-Lopez J, Mateos MV, et al. Daratumumab plus carfilzomib and dexamethasone in patients with relapsed or refractory multiple myeloma. *Blood* 2019. Aug 1;134(5):421-431. doi: 10.1182/blood.2019000722. Epub 2019 May 21.
12. Mikhael JR, Reeder CB, Libby EN, et al. Phase Ib/II trial of CYKLONE (cyclophosphamide, carfilzomib, thalidomide and dexamethasone) for newly diagnosed myeloma. *Br J Haematol*. 2015 Apr; 169(2): 219–227. Published online 2015 Feb 13.
13. Stewart AK, Rajkumar SV, Dimopoulos MA, et al. Carfilzomib, lenalidomide, and dexamethasone for relapsed multiple myeloma. *N Engl J Med*. 2015 Jan 8;372(2):142-52. doi: 10.1056/NEJMoa1411321. Epub 2014 Dec 6.
14. Dimopoulos MA, Moreau P, Palumbo A, et al. Carfilzomib and dexamethasone versus bortezomib and dexamethasone for patients with relapsed or refractory multiple myeloma (ENDEAVOR): a randomised, phase 3, open-label, multicentre study. *Lancet Oncol*. 2016 Jan;17(1):27-38. doi: 10.1016/S1470-2045(15)00464-7. Epub 2015 Dec 5.
15. Papadopoulos KP, Siegel DS, Vesole DH, et al. Phase I study of 30-minute infusion of carfilzomib as single agent or in combination with low-dose dexamethasone in patients with relapsed and/or refractory multiple myeloma. *J Clin Oncol*. 2015 Mar 1;33(7):732-9. doi: 10.1200/JCO.2013.52.3522. Epub 2014 Sep 15.
16. Siegel DS, Martin T, Wang M, et al. A phase 2 study of single-agent carfilzomib (PX-171-003-A1) in patients with relapsed and refractory multiple myeloma. *Blood*. 2012 Oct 4;120(14):2817-25. doi: 10.1182/blood-2012-05-425934. Epub 2012 Jul 25.
17. Vij R, Wang M, Kaufman JL, et al. An open-label, single-arm, phase 2 (PX-171-004) study of single-agent carfilzomib in bortezomib-naive patients with relapsed and/or refractory multiple myeloma. *Blood*. 2012 Jun 14;119(24):5661-70. doi: 10.1182/blood-2012-03-414359. Epub 2012 May 3.

18. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Waldenström Macroglobulinemia/Lymphoplasmacytic Lymphoma Version 3.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
19. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Multiple Myeloma Version 5.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
20. Rosenbaum CA, Stephens LA, Kukreti V, et al. Phase 1/2 study of carfilzomib, pomalidomide, and dexamethasone (KPd) in patients (Pts) with relapsed/refractory multiple myeloma (RRMM): A Multiple Myeloma Research Consortium multicenter study. DOI: 10.1200/JCO.2016.34.15_suppl.8007 *Journal of Clinical Oncology* 34, no. 15_suppl (May 20, 2016) 8007-8007.
21. Meid K, Dubeau T, Severns P, et al. Long-Term Follow-up of a Prospective Clinical Trial of Carfilzomib, Rituximab and Dexamethasone (CaRD) in Waldenstrom's Macroglobulinemia. *Blood* 2017; 130:2772-2772.
22. Yong K, Brown S, Hinsley S, et al. Carfilzomib, cyclophosphamide and dexamethasone is well tolerated in patients with relapsed/refractory multiple myeloma who have received one prior regimen. 2015; 126:1840.
23. Dimopoulos M, Quach H, Mateos MV, et al. Carfilzomib, dexamethasone, and daratumumab versus carfilzomib and dexamethasone for patients with relapsed or refractory multiple myeloma (CANDOR): results from a randomised, multicentre, open-label, phase 3 study. *Lancet*. 2020;396(10245):186-197.
24. Jakubowiak AJ, Dytfeld D, Griffith KA, et al. A phase 1/2 study of carfilzomib in combination with lenalidomide and low-dose dexamethasone as a frontline treatment for multiple myeloma. *Blood*. 2012 Aug 30;120(9):1801-9.
25. Korde N, Zingone A, Kwok M, et al. Phase II Clinical and Correlative Study of Carfilzomib, Lenalidomide, and Dexamethasone (CRd) in Newly Diagnosed Multiple Myeloma (MM) Patients. *Blood*. 2012 Nov 12;120(21):732.
26. Korde N, Zingone A, Kwok ML, et al. Phase II Clinical and Correlative Study Of Carfilzomib, Lenalidomide, and Dexamethasone Followed By Lenalidomide Extended Dosing (CRD-R) Induces High Rates Of MRD Negativity In Newly Diagnosed Multiple Myeloma (MM) Patients. *Blood*. 2013 Nov 15;122(21):538.
27. Zimmerman T, Raje NS, Reece D, et al. Final Results of a Phase 2 Trial of Extended Treatment (tx) with Carfilzomib (CFZ), Lenalidomide (LEN), and Dexamethasone (KRd) Plus Autologous Stem Cell Transplantation (ASCT) in Newly Diagnosed Multiple Myeloma (NDMM). *Blood*. 2016 Dec 2;128(22):675.

28. Yong K, Hinsley S, De Tute, RM, et al. Maintenance with Carfilzomib Following Carfilzomib, Cyclophosphamide and Dexamethasone at First Relapse or Primary Refractory Multiple Myeloma (MM) on the Phase 2 Muk Five Study: Effect on Minimal Residual Disease. *Blood*. 2018 Nov 29;132(1):802.
29. Moreau P, Dimopoulos MA, Yong K, et al. Isatuximab plus carfilzomib/dexamethasone versus carfilzomib/dexamethasone in patients with relapsed/refractory multiple myeloma: IKEMA Phase III study design. *Future Oncol*. 2020 Jan;16(2):4347-4358. doi: 10.2217/for-2019-0431.
30. Manwani R, Mahmood S, Sachchithanatham S, et al. Carfilzomib is an effective upfront treatment in AL amyloidosis patients with peripheral and autonomic neuropathy. *Br J Haematol* 2019 Dec;187(5):638-641.doi: 10.1111/bjh.16122. Epub 2019 Aug 6.
31. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Systemic Light Chain Amyloidosis 1.2022. National Comprehensive Cancer Network, 2022. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed May 2022.
32. Landren O, Hulcrantz M, Diamond B, et al. Safety and Effectiveness of Weekly Carfilzomib, Lenalidomide, Dexamethasone, and Daratumumab Combination Therapy for Patients With Newly Diagnosed Multiple Myeloma: The MANHATTAN Nonrandomized Clinical Trial. *JAMA Oncol*. 2021 Jun 1;7(6):862-868. doi: 10.1001/jamaoncol.2021.0611.
33. Brinthen S, D'Agostino M, De Paoli L, et al. Phase 1/2 study of weekly carfilzomib, cyclophosphamide, dexamethasone in newly diagnosed transplant-ineligible myeloma. *Leukemia*. 2018 Apr;32(4):979-985. doi: 10.1038/leu.2017.327.
34. Gasparetto C, Lipe B, Tuchman S, et al. Once weekly selinexor, carfilzomib, and dexamethasone (SKd) in patients with relapsed/refractory multiple myeloma (MM). DOI: 10.1200/JCO.2020.38.15_suppl.8530 *Journal of Clinical Oncology* 38, no. 15_suppl (May 20, 2020) 8530-8530.

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C88.0	Waldenström macroglobulinemia
C90.00	Multiple myeloma not having achieved remission
C90.02	Multiple myeloma in relapse
C90.10	Plasma cell leukemia not having achieved remission
C90.12	Plasma cell leukemia in relapse
C90.20	Extramedullary plasmacytoma not having achieved remission
C90.22	Extramedullary plasmacytoma in relapse
C90.30	Solitary plasmacytoma not having achieved remission

ICD-10	ICD-10 Description
C90.32	Solitary plasmacytoma in relapse
E85.81	Light chain (AL) amyloidosis
E85.89	Other amyloidosis
E85.9	Amyloidosis, unspecified
Z85.79	Personal history of other malignant neoplasms of lymphoid, hematopoietic and related tissues

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC